

# Occurrence of Diseases in Holdings of Carrier Pigeons with Different Prophylactic Plans



František Zigo<sup>1\*</sup>, Zuzana Farkašová<sup>1</sup>, Silvia Ondrašovičová<sup>2</sup>, Mária Vargová<sup>3</sup> and Katarína Veszelits Laktičová<sup>3</sup>

<sup>1</sup>Department of Nutrition and Animal Husbandry, University of Veterinary Medicine and Pharmacy, Slovakia

<sup>2</sup>Department of Biology and Physiology, University of Veterinary Medicine and Pharmacy, Slovakia

<sup>3</sup>Department of the Environment Veterinary Legislation and Economy, University of Veterinary Medicine and Pharmacy, Slovakia

Submission: October 20, 2021; Published: October 29, 2021

\*Corresponding author: Zigo František, Department of Nutrition and Animal Husbandry, University of Veterinary Medicine and Pharmacy, Slovakia

## Abstract

The aim of the work was to evaluate the occurrence of diseases in two carrier pigeon holdings with different prophylactic plans at the beginning and at the end of flight season. During the flight season, the pigeons completed 14 races at a distance of 140 to 1200 km. In the first holding (A), antibiotics and corticoids were applied at regular intervals in order to maintain health, while in the second holding (B) were used products based on plant extracts and essential oils during the race season. Based on clinical and laboratory examination of faecal samples and oropharynx swabs was obtained a general overview of the prevalence of the most common diseases with regard to their clinical manifestations. The most marked increases in parasitic and infectious agents were observed at the end of the season in both holdings. Despite different prophylactic plans, the prevalence of coccidiosis and trichomoniasis was increased at the end of the racing season on both farms, when the increased incidence of nematodes *Ascaridia* spp., *Capillaria* spp. and ornithosis complex was also confirmed. By comparing the prevalence of individual diseases, a lower incidence of coccidiosis, trichomoniasis and ornithosis complex was observed in breeding A, in which the breeder applied antibiotics. The results of the study point to the phenomenon of immunodeficiency. Although the incidence of monitored diseases on farm A with antibiotic treatment was reduced, the continuous administration of various drugs, particularly antibiotics and corticosteroids, aimed at maintenance of optimum condition of pigeons may impair their homeostasis, pH, and intestinal microflora. This opens the way for pathogens such as salmonella, which was confirmed in our study.

**Keywords:** Race season; Canker; *Coccidiosis*; Ornithosis complex; *Salmonellosis*

## Introduction

Pigeons have accompanied people since time immemorial and today they are a favorite hobby of many breeders due to their attractive way of flying, endurance, loyalty to home and almost perfect orientation in a foreign environment. Increased stress during the race is an important factor that significantly affects their health. The accumulation of a large number of pigeons in one place during their transport increases the potential for transmission of infectious agents, which may later be reflected in reduced performance of birds, increased disease incidence and reduced immunodeficiency [1,2]. The breeders should keep their competitors in good health so they could perform according to expectations. The owner of the pigeons is also responsible for their health and satisfaction in terms of animal protection. Disease prevention and early diagnosis of health disorders (Figure 1) are particularly important for delivering good performance and minimizing the risk of disease spreading to other individuals [3]. One should not forget that improper feeding and insufficient

prophylaxis are common predisposition factors for the development of any disease. The responsible pigeon breeder must take into account the sports performance of his charge (distance and difficulty of the flight route) and have an idea what to expect from the next race. Based on these facts, he should then compile an adequate prophylactic plan and outline compound feed for a particular pigeon [4].

The aim of this study was to evaluate the incidence of the most common diseases of homing pigeons during the racing season and their impact on health and performance of these birds.

## Materials and Methods

### Lofts and pigeon examination

During the race season, two pigeon lofts with different prophylactic programs were monitored. In the loft A, treatment and prevention were based on the use of broad-spectrum

antibiotics based on sulphonamides, tetracycline, ampicillin, erythromycin, streptomycin and corticoids, while in loft B, natural-based preparations in the form of plant extracts, organic acids and essential oils were used. Clinical examination of the health of each pigeon at the beginning and end of the race season was performed according to [5]. Faecal samples and oropharyngeal swabs were

obtained from both lofts from 60 pigeons at the beginning and from 45 pigeons at the end of the race season due to the loss of some birds. Based on clinical signs and laboratory examination, the diagnosis of individual diseases of pigeons was obtained (Figure 2-4).




Figure 1: Disease's prevention in pigeons.




Figure 2: From left: Canker and ornithosis in pigeons.



**Figure 3:** Clinical examination with detection of ectoparasites (*Columbicola columbae*) in pigeons.



**Figure 4:** Clinical symptoms of infected pigeon with *Salmonellosis* spp. -joint form.

Note: Clinical examination of the dove showed palpable feathers, reluctance to fly, weakness, mild dehydration, swelling of the humero-ulno-radial joints with stained cloaca with yellow-green faeces. After unsuccessful treatment with chloramphenicol, an autopsy was performed with sampling for culture and pathogen confirmation.

Photo: Zigo F. (2018).

## Laboratory analysis

The flotation technique of [6] was used to detect coccidiosis and nematodes from faecal samples. Determination of trichomoniasis from oropharyngeal swabs as well as detection of ectoparasites were performed according to [7]. The diagnosis of salmonellosis was made on the basis of clinical signs, autopsy of dead individuals and culturing the samples on buffered peptone agar and on Mac Conkey medium according to [8]. Using the Chi square test, the relationships between the incidence of diseases in the monitored pigeon lofts before and after the end of the race season were statistically evaluated. The differences were considered significant at  $P < 0.05$ .

## Results and Discussion

The racing season is a challenging period for homing pigeons both in terms of their health and fitness and orientation. Pigeons come into contact with other competitors during their transport and with various etiological agents during flight, which can weaken their immunity or cause death in an uncontrollable situation [9]. At the beginning of the race season, different prophylactic programs were applied in the monitored lofts. Their evaluation showed an increased occurrence of trichomoniasis in loft B in which the breeder uses products on a natural basis in the form of plant extracts and essential oils. After the end of the last race, the incidence of coccidiosis, trichomoniasis, end parasitosis and ornithosis complex was increased in both lofts.

The comparison of the incidence of diseases between lofts at the end of the season showed a reduced incidence of trichomoniasis, coccidiosis and ornithosis complex in loft A where antibiotics were regularly administered during the race. On the contrary, an increased incidence of endoparasitosis and two clinical cases of salmonellosis were recorded in this loft and ended in the death of the positive pigeons (Table 1). According to Bergman [4], the most common diseases of pigeons include coccidiosis, trichomoniasis and respiratory infections, which is confirmed by our results from the monitored lofts. In particular, coccidiosis associated with impaired immunity affects the digestion and composition of the bacterial microflora. This protozoa parasite produces in the intestines 'oocysts', which are passed out in the faeces. The oocyst matures in the environment and will then affect other birds if ingested. Most adult birds carry low levels of these parasites. Only when large numbers of parasites are present is treatment necessary. Usually raised levels of coccidial oocysts are associated with suboptimal performance but in young birds and adults under stress an acute clinical form of the disease may be seen. The parasite affects the lining of the intestine causing diarrhea and blood may be present. Affected birds are depressed, rapidly become emaciated and may die [10]. The positive cases

recorded in our study at the end of the racing season showed a thin consistency of faeces, mild dehydration and dirty feathers around the cloaca. The accompanying phenomenon in these individuals was a higher weight loss compared to healthy pigeons. On farms with an increased incidence of coccidiosis often other endoparasites in the form of a mixed infection are detected. In Istanbul, mixed coccidial and nematode infections have been found in urban pigeons nesting in well-known mosques: *Capillaria obsignata* - 19.3% and *Ascaridia columbae* - 14.6% [11]. In our study, an increased incidence of endoparasitosis was detected at the end of the racing season, especially in loft B, where *Ascaridia* spp. and *Capillaria* spp. were detected in coccidiosis-positive pigeons in the same faecal samples. Intestinal parasites can affect birds of all ages but due to some degree of resistance in older birds the more extreme clinical signs are more commonly seen in the younger birds. Weight gain in young birds may be adversely affected and in more severe infestations diarrhea, depression and loss of condition, even emaciation, may be obvious, as may anemia. In mild infestations clinical signs may not be obvious which was also confirmed in our study where positive individuals were compared to other pigeons in their performances.

**Table 1:** The most common diseases of pigeons in monitored holdings.

Diseases	Before Race Season (n = 60)			After Race Season (n = 45)		
	Holding A %	Holding B %	P Value	Holding A %	Holding B %	P Value
Coccidiosis	8.3	3.3	NS	35.6	46.7	P<0.05
Trichomoniasis	5	11.7	P<0.05	20	44.4	P<0.05
Ectoparasitosis <sup>1</sup>	0	0	NS	8.9	4.4	NS
Endoparasitosis <sup>2</sup>	0	0	NS	4.4	20	P<0.05
Ornithosis Complex	0	0	NS	11.1	24.4	P<0.05
Salmonellosis	0	0	NS	4.4	0	NS

Note: n-number of isolated strains of total 60 investigated pigeons before race and from 45 pigeons after race from each holding, Ectoparasitosis<sup>1</sup>- caused by *Columbicola columbae*, *Ceratophylus columbae*, Endoparasitosis<sup>2</sup> - caused by *Ascaridia* spp., *Capillaria* spp. NS-nonsignificant.

Another common disease recorded at the end of the season from oropharyngeal swabs was trichomoniasis. Trichomoniasis or canker is caused by a parasitic organism called *Trichomonas colombi* and three forms are recognized that affect the pharynx (throat), navel and internal organs respectively. The majority of adult pigeons are symptomless carriers of the organism, but clinical cases may occur if the bird is under stress and in young pigeons the disease may be severe and even fatal. The disease is spread from adults to squabs in the crop milk and between pigeons by the oral route [12]. In our study, we observed an increased incidence of pharyngeal form in both holdings at the end of the season. It was visible cheesy yellow deposits on the membranes of the pharynx at the back of the mouth during the clinical examination. These deposits affect food intake and also breathing. One of the difficult to detect and treat diseases is

ornithosis complex, which was confirmed in both farms at the end of the season. Ornithosis complex is one of the combined diseases that can be caused by the microorganisms *Chlamydia psittaci*, *Pasteurella* spp. or mycoplasma.

Clinical signs vary from poor performance to an acute disease causing a marked conjunctivitis, decreased appetite, respiratory disease, diarrhea and death. The acute form is particularly prevalent in young birds and older birds under stress. In some cases, the conjunctivitis can lead to secondary bacterial infection of the eye resulting in blindness. Recovered birds can transmit the disease and subsequently spread it to other pigeons [13]. The positive cases of individuals recorded by us showed a mild form of conjunctivitis with shortness of breath, especially after returning from the race.

## Conclusion

The monitoring of diseases at the beginning and end of the race season in two lofts pigeons and using different prophylactic plans provided an overview of the health status of housed birds. The prevalence of monitored diseases in both lofts was the highest at the end of the racing season. Comparison of lofts showed a reduced incidence of coccidiosis, trichomoniasis and ornithosis complex in loft A, in which the breeder administered antibiotics in regular cycles to maintain pigeon's health. Although the incidence of monitored diseases in loft A was reduced it should be borne in mind that breeders who constantly apply various drugs, especially antibiotics and corticoids, to their pigeons to maintain their "optimal shape", reduce the competitors 'lifespan' and disrupt their immune system. It is the long-term application of antibiotics and corticoids that causes the phenomenon of immunodeficiency, in which homeostasis and pH are disrupted, the intestinal microflora is declassified, thus opening the way for pathogenic agents such as salmonellosis, which was confirmed also in our study.

## Acknowledgment

This study was supported by the Slovak project KEGA 009UVLF-4/2021: Innovation and implementation of new knowledge of scientific research and breeding practice to improve the teaching of foreign students in the subject of Animal husbandry.

## References

1. Duncan JR, Prasse KW, Mahaffey EA (1994) Veterinary laboratory medicine Clinical Pathology. Iowa State University Press USA.
2. Hand O (2004) The Effects of Stress on Racing Pigeons.
3. Balicka Ramisz A, Pilarczyk B (2014) Occurrence of coccidia infection in pigeons in amateur husbandry Diagnosis and prevention. An of Par 60: 93-97.
4. Bergmann R (2013) Sports performances at the pigeon (Czech). Letu zdar 23(1): 22.
5. Kimpe A, Decostere A, Martel A, Haesebrouck F, Devriese LA (2002) Prevalence of antimicrobial resistance among pigeon isolates of *Streptococcus gallolyticus*, *Escherichia coli* and *Salmonella enterica* serotype Typhimurium. Avian Pathol 31: 393-397.
6. Dranzoa C, Ocaido M, Katete P (1999) The ecto gastrointestinal and haemo parasites of live pigeons (*Columba livia*) in Kampala Uganda. Av Pat 28: 119.
7. Letková V, Goldová M, Cszimárová G (1997) Laboratory diagnostics in veterinary parasitology (Slovak) Košice Datahelp p.71.
8. Stenzel T, Koncicki A (2007) Occurrence of parasitic invasions in domestic pigeons (*Columba livia domestica*) in the Northern Poland. Pol J of Vet Sci 10: 275-278.
9. Zigo F, Takáč L, Chripková M (2016) Infectious diseases of carrier pigeons and antimicrobial resistance in isolated bacteria. Int J of Sci Res 5: 2277-2817.
10. Rupipper DJ (1998) Diseases that affect race performance of homing pigeons. Bacterial fungal and parasitic diseases 12(3): 138-148.
11. Koroglu E, Şimşek S (2001) The prevalence of *Eimeria* species in pigeons (*Columba livia*) in Elazığ. Fırat Üniversitesi Sağlık Bilimleri Dergisi Vet 15: 401-403.
12. Bogdan B (2017) Trichomonoz 22.
13. De Weerd HJM (2012) Ornithosis complex in racing pigeons. The racing pigeon 22.
14. Dik B, Halajian A (2013) Chewing Lice (*Phthiraptera*) of Several Species of Wild Birds in Iran with New Records. Journal of Arthropod Borne Diseases 7: 83-89.



This work is licensed under Creative Commons Attribution 4.0 License  
DOI: [10.19080/AAPS.2021.01.555571](https://doi.org/10.19080/AAPS.2021.01.555571)

### Your next submission with Juniper Publishers will reach you the below assets

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats  
( Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

### Track the below URL for one-step submission

<https://juniperpublishers.com/online-submission.php>