

A Study on Major Ectoparasites and Associated Skin Diseases of Ruminants in and around Bishoftu Town, Central Ethiopia



Samson Amanuel¹ and Temesgen Kassa*²

¹College of Veterinary Medicine and Agriculture, Addis Ababa University, Ethiopia

²Holeta Animal Health Research National Program, Ethiopian Institute of Agricultural Research, Ethiopia

Submission: February 17, 2023; **Published:** April 03, 2023

***Corresponding author:** Temesgen Kassa, Holeta Animal Health Research National Program, Ethiopian Institute of Agricultural Research, Ethiopia

Abstract

A cross-sectional study was conducted from November 2018 to April 2019 to evaluate the occurrence of ectoparasites and associated skin diseases in ruminants in and around Bishoftu town, central Ethiopia. A total of 437 ruminants (216 cattle, 141 sheep, and 80 goats) brought to the open-air Veterinary Teaching Hospital (VTH) of the College of Veterinary Medicine, Addis Ababa University, were sampled using simple random sampling. Accordingly, the occurrence of ectoparasites and associated skin diseases was found to be 59% (258), and out of these animals, 195 (44.6%) were infected with ectoparasites, of which 111 (25.4%) were bovine, 27 (6.2%) were caprine, and 57 (13%) were ovine. The major ectoparasite species identified were *Amblyomma variegatum* (13%), *A. gemma* (0.7%), *A. cohaerens* (1.8%), *A. nymph* (3.2%), *H. truncatum* (2.7%), *Boophilus decoloratus* (6.2%), *Rhipicephalus pulchellus* (1.8%), *R. eversi* (0.7%), *Damalimia ovis* (3%), *Linognathus vituli* (5.5%), *Psoroptes* (0.7%), *Demodex bovis* (0.5%), *Ctenocephalides canis* (0.7%) and *Ct. felis* (5.03). On the other hand, the associated skin diseases infesting animals were: 63 (14.4%), lumpy skin disease (0.9%), sheep pox (5.5%), goat pox (5%), dermatophilosis (0.2%), contagious ecthyma in caprine (2.1%), and contagious ecthyma in ovine (0.7%). The findings of the present study suggest that ectoparasites and associated skin diseases in and around Bishoftu are still widespread and economically important constraints for the productivity and use of ruminants. Hence, stringent adherence to the control of ectoparasites and associated skin diseases is required as a responsibility of the concerned government body based on professional research results in the study area.

Keywords: Associated skin diseases; Ectoparasite; Prevalence; Ruminants; Bishoftu; Ethiopia

Introduction

Ethiopia is fundamentally an agrarian country whose livelihood depends on plant cultivation and animal husbandry. It has a numerous livestock population with an estimated 70 million herds of cattle, 30.7 million sheep, 30.2 million goats, and 56.53 million poultry [1]. This sector provides various functions in Ethiopia as sources of meat, milk, manure, fertilizer, drought power, and employment [2]. In addition, it contributes 19% of the GDP and 16-19% of the foreign exchange earnings of the country [3]. Livestock is also a major source of foreign currency through the export of chilled meat, live animals, hides, and skins [4]. Because of this huge potential, Ethiopia is capable of supplying 16-18 million skins and hides per year for the tanneries within the country [5].

Ectoparasites cause serious economic loss in ruminants through the mortality of animals, decreased production, downgrading, and rejection of skin and hide [6,7]. Moreover, they

are the most important vectors of protozoan, bacterial, viral, and rickettsial diseases [8]. Ectoparasite diseases such as sarcoptic and psoroptic mange, tick, flea, and louse infestation have frequently been reported in Ethiopia, and these are among the threats that result in serious economic loss to the tanning industry and the country as a whole [9].

Among the ectoparasites infesting cattle, sheep, and goats, ticks are very significant and harmful because of their blood-sucking habits and direct debilitating effect [10]. In Ethiopia, there are 47 species of ticks found on livestock, and most of them are important as vectors and disease-causing agents and also have a mechanical damaging (tick bite) effect on skin and hide production [11].

Lice are one of the common parasites of domestic ruminants. The most common and clinically significant type of lice is the chewing louse. Chewing lice may cause less individual damage

than sucking lice, but because they are present in greater numbers, they can cause extremely severe damage. Poor control may be associated with a failure to detect or identify lice infestation in its initial stages, and by the time clinical diagnosis is achieved, the entire herd may be infested [12].

Mites are an important ectoparasite of livestock with great veterinary and medical significance. Due to their behavior, mites may have a direct or indirect effect on their host. These could be direct (blood loss, skin inflammation, purities, etc.) or indirect (cause disturbance and self-wounding) when present in high density [10]. The mites damage the skin, which is usually accompanied by irritation, rubbing, and scratching. On a larger scale, mite infections have a great impact on the local and international trade of animals [13].

Fleas are the most common ectoparasite-causing dermatitis in livestock. The feeding activity of this ectoparasite results in significant blood loss, secondary infestation, pruritus, excoriation, and, in some cases, premature death. A heavy infestation in young animals may cause severe anemia and death [10,14]. From the flea species, *Ctenocephalides felis*, and *Ctenocephalides canis* are frequently recognized in ruminants [15].

Various skin diseases affecting ruminants are frequently reported from different parts of Ethiopia. These different skin diseases are liable for considerable economic losses, especially to the skin and hide export industry, due to various defects, 65% of which occur in the pre-slaughter conditions and are most directly related to the skin and hide rejection because of inferior quality. Apart from quality degradation, skin and hide diseases induce associated economic losses due to reduction of wool quality, meat and milk yield, losses as a result of culling and occasional mortalities, and related costs of treatment and prevention of the diseases. Namely, dermatophilosis, dermatophytosis, photosensitization, lumpy skin disease, contagious ecthyma, and sheep and goat pox have been frequently reported in Ethiopia, which is badly affecting the product, health, and economy of the country [16,17]. Even though several studies have been conducted in the study area, ectoparasites are still a serious problem that causes significant economic losses. Therefore, this study aimed to identify the major ectoparasite prevailing in and around Bishoftu, Ethiopia, and to evaluate the prevalence of ectoparasites and associated skin diseases in ruminants.

Material and Methods

Study area

The study was carried out in and around Bishoftu. Bishoftu is a town in Adia district, in the east Shoa zone of the Oromia regional state, in central Ethiopia. It is located 47 kilometers south of Addis Abebe on the declivity of Ethiopia's Great Rift Valley. It is situated at 9° N latitude and 40° S longitude, at an altitude of about 1850 meters above sea level. The area receives an average annual rainfall of 600mm (400-800mm). The rainy season is

bimodal, with the short rainy season occurring from March to May and the long rainy season beginning from June to September. In addition, the dry season extends from October to February. The mean annual minimum and maximum temperatures are 10.91°C and 29.45°C, respectively [18]. The livestock production system in the area is both intensive and extensive.

Study animals

The animals used in this study were domestic ruminants: cattle, sheep, and goats. Local and cross-bred animals of both sexes, as well as young and adult age groups, are among those included. According to [19], lambs and children aged 6 months and under were considered young, while those aged 6 months and older were considered adults. In addition, body condition scores were grouped into poor, medium, and good according to [20]. All the above-mentioned animals presented to the open-air veterinary teaching hospital (VTH) of the Addis Ababa University College of Veterinary Medicine, which is found in Bishoftu town. Those animals were either clinically sick and came to get medication or to obtain other related services such as pregnancy diagnosis, vaccination, and castration.

Study design

The study design implemented was cross-sectional. A total of 437 ruminants (216 cattle, 141 sheep, and 80 goats) were examined during the study period from November 2018 to April 2019 to identify the major ectoparasites prevailing in the study area and associated skin diseases. Cattle, sheep and goats presented each day in VTH were sampled using simple random sampling.

Sample collection and examination

Sample collection was carried out after the animal was physically restrained, followed by visual inspection and palpation of the head, neck, dewlap, brisket, axillae, belly, prepuce, scrotum, udder, groins, and tail regions. These were performed to search for ectoparasites and lesions on the external body of the animal. Then, a detailed history of the animal, such as address, species, sex, age, body condition, site collection, and laboratory results, was written on a special field registration format prepared for this study.

The ectoparasites encountered during the study period (ticks, lice, and fleas) were collected manually from their sites of attachment. The removal of ticks was performed with great care with the aid of forceps so as to avoid detaching the mouth parts of the ticks, which helps to identify the genus easily. Coat brushing techniques were used for lice collection. Fleas were collected from the infested body parts using saturated gauze soaked in 70% ethanol. Identification of their genus or species was made based on differentiating structures that included the width and length of the head and the combs and spines on the head or legs. The removed ectoparasite from the animal's body was dropped into a labeled universal bottle containing 70% ethanol for later identification. Forceps were used to remove the collected ectoparasites from the

bottle and place them in a Petri dish. Meanwhile, the petridish was placed under a stereomicroscope and examined according to the morphological descriptions given by [21,22]. For the suspected mange mite infestations, skin scrapings were taken by clipping the hair around the periphery of active lesions in the affected area with a scalpel blade [22]. The skin was squeezed firmly and scraped deeply until blood was drawn. The scrapped materials were transferred to petridish and taken for laboratory examination. A few drops of 10% potassium hydroxide were added to the specimen, which was allowed to stand for 30 minutes before being appreciated under a light microscope at 40 x magnifications. The mange mites were identified according to [23].

For the suspected cases of dermatophilosis, scabs, pus, and exudates were collected, and Gram stain smears were made according to the procedures described by [24]. Diagnoses of contagious ecthyma and sheep and goat pox are made based on flock history and clinical signs such as lesions around the oral cavity, body, and tail region [19].

Sample Size Determination

As previous studies on species identification of ectoparasites in the study area had not been conducted, the expected prevalence was assumed to be 50%. Therefore, the sample size was calculated at a 50% prevalence rate with a desired precision of 5% and a 95% confidence interval. The desired sample size for the study was calculated using the formula given by [25].

$$n = 1.96^2 \frac{P_{exp}(1 - P_{exp})}{d^2}$$

Table 1: Occurrence of Ectoparasite.

| Factors | Variables | Tick(%) | Fleas(%) | Lice(%) | Mite(%) | X ² -value | P-value |
|---------|-----------|-----------|----------|----------|---------|-----------------------|---------|
| Species | Bovine | 85(65.9) | 2(8) | 20(54.1) | 4(100) | 42.59 | 0 |
| | Caprine | 17(13.2) | 6(24) | 4(10.8) | 0 | | |
| | Ovine | 27(20.9) | 17(68) | 13(35.1) | 0 | | |
| Sex | Male | 100(77.5) | 11(44) | 14(37.8) | 3(75) | 32.81 | 0 |
| | Female | 29(22.5) | 14(56) | 23(62.2) | 1(25) | | |
| Breed | Cross | 5(3.9) | 0 | 0 | 0 | 2.5 | 0.561 |
| | Local | 124(96.1) | 25(100) | 37(100) | 4(100) | | |
| Age | Adult | 114(88.4) | 12(48) | 18(48.6) | 3(75) | 41.06 | 0 |
| | Young | 15(11.6) | 13(52) | 19(54.5) | 1(25) | | |
| BCS | Good | 23(17.8) | 2(8) | 1(2.7) | 0 | 46.85 | 0 |
| | Medium | 71(55) | 13(52) | 6(16.2) | 2(50) | | |
| | Poor | 35(27.1) | 10(40) | 30(81.1) | 2(50) | | |

Species of ectoparasite

Amblyomma variegatum, *Linognathus vituli*, *Psoroptes*, and

Where, P_{exp} = expected prevalence; d = absolute precision; n = sample size.

Accordingly, a total of 384 cattle, sheep, and goats needed to be sampled. However, the sample size was raised to 437 cattle, sheep, and goats to increase the precision of the study.

Data analysis

The entire collection of raw data was entered into a Microsoft Excel spreadsheet and coded. For statistical analyses, R version 4.01 software packages were used. A percentage was used to calculate the prevalence. Additionally, chi-square was used to measure the degree of association between risk factors and the prevalence of ectoparasites and associated skin diseases. In the analysis, a difference was taken as significant at a p-value less than 0.05, and the confidence level was held at 95%.

Results

The occurrence of ectoparasite

Out of 384 animals sampled, 258 (59%) were infested by ectoparasites and skin diseases. From 216 bovine, 80 caprine, and 141 ovines, 116 (26.5%), 58 (13.3%), and 84 (19.2%) were infected, respectively. From the total infected animals, 195 (44.6%) were infected with ectoparasites, of which 111 (25.4%) bovine, 27 (6.2%) caprine, and 57 (13%) ovine were infested with ectoparasites. Bovine was highly infested with ticks, and local breeds were the most affected with ectoparasites (Table 1).

Ctenocephalides felis were the frequently occurring problems of tick, louse, mite and flea species with respective prevalences of 13%, 5.5%, 0.7%, and 5.03% (Table 2).

Table 2: Species of Ectoparasite.

| Ectoparasite Species | Number Positive | Prevalence (%) |
|---------------------------------|-----------------|----------------|
| Ticks | | |
| <i>Amblyomma variegatum</i> | 57 | 13 |
| <i>Amblyomma gemma</i> | 3 | 0.7 |
| <i>Amblyomma cohaerens</i> | 8 | 1.8 |
| <i>Amblyomma nymph</i> | 14 | 3.2 |
| <i>Hyalomma truncatum</i> | 12 | 2.7 |
| <i>Boophilus decoloratus</i> | 27 | 6.2 |
| <i>Rhipicephalu spulchellus</i> | 8 | 1.8 |
| <i>Rhipicephaluseversi</i> | 3 | 0.7 |
| Lice | | |
| <i>Damalinia ovis</i> | 13 | 3 |
| <i>Linognatus vituli</i> | 24 | 5.5 |
| Mites | | |
| <i>Psoroptes</i> | 3 | 0.7 |
| <i>Demodex bovis</i> | 2 | 0.5 |
| Fleas | | |
| <i>Ctenocephalides canis</i> | 3 | 0.7 |
| <i>Ctenocephalides felis</i> | 22 | 5.03 |

Overall prevalence of associated skin diseases

According to our study, 63 (14.4%) cases were caused by associated skin diseases. The result showed that sheep pox was

the most common, with a prevalence of 5.5%, and dermatophilosis (0.2%) was the least prevalent problem in the study animals (Table 3).

Table 3: Commonly occurring associated skin diseases.

| Skin Problems | Positive Animals | Prevalence (%) |
|-------------------------------|------------------|----------------|
| Lumpy skin disease | 4 | 0.9 |
| Sheep pox | 24 | 5.5 |
| Goat pox | 22 | 5 |
| Dermatophilosis | 1 | 0.2 |
| Contagious ecthyma in ovine | 3 | 0.7 |
| Contagious ecthyma in caprine | 9 | 2.1 |

Species wise occurrence of ectoparasites and associated skin diseases

Ectoparasite species *A. gemma*, *A. cohaerens*, *R. eversi*, *Psoroptes*, *D. bovis*, and *C. canis* were isolated only from bovines. Bovine species were not infested with *Damalina ovis*. Caprine species examined for ectoparasites were infested with *A. variegatum*, *A. nymphs*, *B. decoloratus*, *R. pulchellus*, *Ct. felis*, and

L. vituli. Ovine ectoparasites identified were: *A. variegatum*, *A. nymph*, *H. truncatum*, *B. decoloratus*, *R. pulchellus*, *R. eversi*, *D. ovis*, and *Ct. felis* (Table 4).

Occurrence of ectoparasites and associated skin diseases according to body condition

Most ecto-parasites and other skin diseases were commonly identified from animals with poor body conditions (Table 5).

Table 4: Occurrence of ectoparasites and associated skin diseases according to species.

| Ectoparasite Species | Total Positive | Animal Species | | | X ² - Value | P-Value |
|---------------------------------|----------------|----------------|-------------|-----------|------------------------|---------|
| | | Bovine (%) | Caprine (%) | Ovine (%) | | |
| Ectoparasites | | | | | | |
| <i>A. variegatum</i> | 57 | 40(70.2) | 1(1.8) | 16(25.1) | 15.876 | 0.001 |
| <i>A. gemma</i> | 3 | 3(100) | 0 | 0 | 3.091 | 0.251 |
| <i>A. cohaerens</i> | 8 | 8(100) | 0 | 0 | 8.338 | 0.017 |
| <i>A. nymph</i> | 14 | 8(57.1) | 3(21.4) | 3(21.4) | 0.77 | 0.723 |
| <i>H. truncatum</i> | 12 | 9(75) | 0 | 3(25) | 4.093 | 0.33 |
| <i>B. decoloratus</i> | 27 | 12(44.4) | 10(37) | 5(18.5) | 7.345 | 0.024 |
| <i>R. pulchellus</i> | 8 | 2(25) | 3(37.5) | 3(37.5) | 2.693 | 0.263 |
| <i>R. eversi</i> | 3 | 3(100) | 0 | 0 | 3.091 | 0.251 |
| <i>D. ovis</i> | 13 | 0 | 0 | 13(100) | 28.128 | 0 |
| <i>L. vituli</i> | 24 | 20(83.3) | 4(16.7) | 0 | 14.137 | 0.001 |
| <i>Psoroptes</i> | 3 | 3(100) | 0 | 0 | 3.091 | 0.251 |
| <i>D. bovis</i> | 2 | 2(100) | 0 | 0 | 2.056 | 0.515 |
| <i>Ct. canis</i> | 3 | 3(100) | 0 | 0 | 1.819 | 0.459 |
| <i>Ct. felis</i> | 22 | 1(5) | 6(27) | 15(68) | 2.104 | 0.508 |
| Associated Skin Diseases | | | | | | |
| Lumpy skin disease | 4 | 4(100) | 0 | 0 | 4.13 | 0.0146 |
| Sheep pox | 24 | - | - | 24(100) | 53.311 | 0 |
| Goat pox | 22 | - | 22(100) | - | 103.37 | 0 |
| Dermatophilosis | 1 | 1(100) | 0 | 0 | 1.025 | 0.999 |
| Contagious ecthyma | 12 | 0 | 9(75) | 3(25) | 27.964 | 0 |

Table 5: Occurrence of Ectoparasite and associated skin diseases according to body condition.

| Ectoparasite Species | No. Positive | Body Condition | | | X ² -Value | P-Value |
|-----------------------|--------------|----------------|--------------|--------------|-----------------------|---------|
| | | Good No. | Medium No. | Poor No. | | |
| | | Positive (%) | Positive (%) | Positive (%) | | |
| Ectoparasites | | | | | | |
| <i>A. variegatum</i> | 57 | 18(31.6) | 28 (49.1) | 11(19.3) | 27.24 | 0 |
| <i>A. gemma</i> | 3 | 0 | 3(100) | 0 | 2.345 | 0.294 |
| <i>A. cohaerens</i> | 8 | 1(12.5) | 4(50) | 3(37.5) | 0.135 | 0.111 |
| <i>A. nymph</i> | 14 | 0 | 9(64.3) | 5(37.5) | 1.972 | 4.439 |
| <i>H. truncatum</i> | 12 | 0 | 8(66.7) | 4(33.3) | 1.652 | 0.562 |
| <i>B. decoloratus</i> | 27 | 12(44.4) | 10(37) | 5(18) | 7.345 | 0.024 |
| <i>R. pulchellus</i> | 8 | 2(25) | 3(37.5) | 3(37.5) | 2.693 | 0.263 |
| <i>R. eversi</i> | 3 | 3(100) | 0 | 0 | 3.091 | 0.251 |
| <i>D. ovis</i> | 13 | 0 | 12(15.4) | 11(84.6) | 16.68 | 0.001 |
| <i>L. vituli</i> | 24 | 1(4.2) | 4(16.7) | 19(79.2) | 25.57 | 0 |
| <i>Psoroptes</i> | 3 | 0 | 2(66.7) | 1(33.3) | 0.4 | 0.111 |
| <i>D. bovis</i> | 2 | 0 | 1(50) | 1(50) | 0.44 | 0.112 |
| <i>Ct. canis</i> | 3 | 1(33.3) | 0 | 2(66.7) | 1.81 | 0.045 |

| | | | | | | |
|---------------------------------|----|--------|----------|----------|-------|-------|
| <i>Ct. felis</i> | 22 | 2(9.1) | 12(54.6) | 8(36.36) | 2.1 | 0.005 |
| Associated Skin Diseases | | | | | | |
| LSD | 4 | 0 | 1(25) | 3(75) | 3.449 | 0.179 |
| Sheep pox | 24 | 0 | 7(29.2) | 17(70.8) | 17.93 | 0 |
| Goat pox | 22 | 0 | 1(25) | 3(75) | 6.698 | 0.033 |
| Dermatophilosis | 1 | 0 | 1(100) | 0 | 0.778 | 0.132 |
| Contagious ecthyma | 12 | 0 | 5(41.7) | 7(58.3) | 4.479 | 0.102 |

Discussion

The study revealed that skin diseases caused by ectoparasites in and around Bishoftu town were causing serious problems in the area. Ticks, lice, fleas, psoroptic mange, and demodectic mange were the most common ectoparasites, while dermatophytosis and viral diseases such as lumpy skin disease, contagious ecthyma, and sheep and goat pox skin diseases were encountered in the study area.

The occurrence of ectoparasites in cattle, sheep, and goats was 25.4%, 13%, and 6.2%, respectively. This finding was lower than the findings of [26], who reported 73.3% [17] who reported 40.2%, and [27], who reported 27.3%, while slightly higher than the findings reported by [28], who reported 15.41%. This difference may be due to the agro-climatic conditions, the health care of the animals, or a lack of regular deworming habits.

On the other hand, in the present study, 13% of the sheep were infested with one or more ectoparasites, which was lower than the prevalence reported by different authors from other regions of Ethiopia: 55.7% by [17], 81.5% by [29], 99.38% by [30] and 78.4% by [7]. Variations in geographical locations, seasons, and management practices in the different study areas might have contributed to the disparity in prevalence.

In this study, the major ectoparasites identified were ticks, lice, mites, and fleas, with an overall prevalence of 30.1%, 8.5%, 1.2%, and 0.9%, respectively. It was found that tick infestation (30.1%) was the outstanding ectoparasite in cattle, and the genera identified were *Boophilus* (6.2%), *Rhipicephalus* (2.5%), *Amblyomma* (18.7%), and *Hyalomma* (2.7%). The genera found were similar to those found by [17] and [26]. The dominance of tick infestation among others is in agreement with the work of [27]. The relatively higher prevalence of tick infestation in this study might be due to the humidity and tropical climate that favor the survival and reproduction of ticks.

In the present study, the identified genera of ticks in cattle, goats, and sheep were *Amblyomma* (56.5%), *Boophilus* (118.6%), *Hyalomma* (8.3%), and *Rhipicephalus* (7.6%). Of these, a higher proportion of *Amblyomma* (56.5%) and *Boophilus* (18.6%) were observed in cattle. Moreover, two genera of lice; *Linognathus* and *Damalina*, were identified in cattle and sheep, respectively. *Ct.*

felis in cattle, sheep, and goats and *Ct. canis* in cattle were also identified. Similar genera of ticks, lice, and mites were identified in different locations of the country by different authors [17,31].

The overall prevalence of major skin diseases in ruminants encountered in the current study was 14.4%. This is comparable with the report of [17], which found 15.41%. According to [32], dermatophilosis is a treat for livestock production in Ethiopia; 0.2% of cattle were obtained in the study area. The finding was lower than the findings of [33], who reported 5.22%, and [34], who reported 8.7%. The low prevalence of dermatophilosis may be attributed to agro-climate changes, the season of the study conducted, and the management system of the animal, but in small ruminants' no prevalence was registered in the study period.

Lumpy skin disease was only found in cattle, accounting for 0.9% of the prevalence in the presented study area, which was lower than the previous study of 0.68% conducted at the Adama veterinary clinic by [17] and also lower than the study conducted in Wolliso (Southwest Oromia) by [35], which showed a prevalence rate of 27.91%. This is assumed to be a result of the study period, which was not the season of multiplication of flies, which act as mechanical vectors for the virus. Because, in the Ethiopian context, the availability of flies during the spring season exacerbates the viral infection rate of lumpy skin disease as a mechanical vector.

Contagious ecthyma (orf) had a prevalence of 0.7% in sheep and 2.1% in goats at the time of the study, while poxvirus disease had a prevalence of 5.5% in sheep and 5% in goats. These findings were lower than the findings of [17] who reported contagious ecthyma in 4.91% of sheep and 9.21% of goats and the poxvirus disease in 8.55% of goats and 8.87% of goats in the Adama veterinary clinic. Hence, the prevalence carried out at Bishoftu Veterinary Teaching Hospital was less than that of the study conducted at Adama Veterinary Clinic. This may be due to the difference in the climatic conditions of the area. Since contagious ecthyma and sheep and goat pox occur mostly in drought periods.

Conclusion and Recommendation

The presented study concerning ectoparasites and associated skin diseases was undertaken on ruminants, namely, cattle, sheep, and goats. In the study area, ticks, lice, fleas, dermatophilosis, demodectic and psoroptic mange, and viral diseases such as

lumpy skin disease, pox virus, and contagious ecthyma were discovered, resulting in the condemnation or downgrading of the quality of hides and skins in tanneries, a reduction in production and productivity, and, in some cases, death. Those resulted in the greatest economic losses to ruminant production and were also subject to secondary infection. The higher prevalence of ectoparasites in ruminants could be due to the government giving less attention to preventing the diseases in terms of campaign immunization before exposure if available and supporting the livestock sector in disease prevention. Unable to use effective acaricide, lack of proper management by the owner, insufficient awareness creation, and education of the owner by veterinarians paved the way for the problem to persist.

Based on aforementioned idea the following recommendations are forwarded.

i. Continuous research must be conducted using scientifically sound data to determine the incidence and distribution of various ectoparasites and associated skin diseases to identify which species are veterinary significant in that area, and the responsible body should take immediate action.

ii. Ruminant owners should receive education regarding how to dip with conventional acaricides, maintain a clean ruminant barn, and isolate an infected host.

iii. To perform effectively, strict adherence to the control of ectoparasites and associated skin diseases is required as a responsibility of the concerned government body based on professional research results on the study area.

Author's Contribution

Samson Amanuel has recorded all the laboratory and survey data and compiled and written the write-up of the paper. Temesgen Kassa contributes through data analysis and edition.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

1. CSA (2021) Agricultural sample survey. Report on Livestock and Livestock Characteristics statistical bulletin Volume II, Statistical Bulletin 589, Central Statistical Authority, Addis Ababa, Ethiopia.
2. Solomon A (2003) Livestock marketing in Ethiopia: a review of structure, performance, and development initiatives.
3. Alemnesh B, Getachew T, Tariku J (2018) Assesment of quality and marketing of hide and skin in Adamitulu Jidokombolcha and Bora Woreda in East Shewa Zone of Oromia Regional State, Ethiopia. *Int J Livest Prod* 9(10): 269-279.
4. Sheferaw D, Degefu H, Banteyirgu D (2010) Epidemiological study of small ruminant mange mites in three agro-ecological zones of Wolaita, Southern Ethiopia. *Ethiop Vet J* 14(1): 31-38.
5. Kabir MHB, Mondal MMH, Eliyas M, Mannan MA, Hashem MA, et al. (2011) An epidemiological survey on investigation of tick infestation in cattle at Chittagong District, Bangladesh. *African Journal of Microbiology Research* 5(4): 346-352.
6. Tiki B, Addis M (2011) Distribution of ixodid ticks on cattle in and around Holeta town, Ethiopia. *Global Veterinaria* 7(6): 527-531.
7. Fentahun T, Woldemariam F, Chanie M, Berhan M (2012) Prevalence of ectoparasites on small ruminants in and around Gondar Town. *American-Eurasian Journal of Scientific Research* 7(3): 106-111.
8. Radostits OM, Gay C, Hinchcliff KW, Constable PD (2007) A textbook of the diseases of cattle, sheep, goats, pigs and horses. *Vet. Med.* 10th (edn). Bailliere, Tindall, London, UK, pp. 1576-1580.
9. Yacob HT (2014) Ectoparasitism: threat to Ethiopian small ruminant population and tanning industry. *Journal of Veterinary Medicine and Animal Health* 6(1): 25-33.
10. Taylor MA, Coop RL, Wall RL (2015) *Veterinary parasitology*. John Wiley & Sons.
11. Kassa B (2005) Standard veterinary laboratory diagnostic manual. Vol. III. MOA Addis Ababa, Ethiopia.
12. Milnes AS, Green LE (1999) Prevalence of lice on dairy cattle in England and the bordering counties of Wales. *Vet Rec* 145(13): 357-362.
13. Maxie MG, Jubb KVF, Kennedy PC, Palmer N (2007) Judd, Kennedy, and Palmer's pathology of domestic animals. Elsevier Saunders.
14. Agarwal DP, Gupta AR (2010) Management of ectoparasites of livestock. *Dep. Med. QUAT-Bhubaneswar, Orissa, India*.
15. Yeruham I, Rosen S, Perl S (1997) An apparent flea-allergy dermatitis in kids and lambs. *Zentralbl Veterinarmed A* 44(7): 391-397.
16. Woldemeskel M (2000) Dermatophilosis: a threat to livestock production in Ethiopia. *Dtsch Tierarztl Wochenschr* 107(4): 144-146.
17. Yacob HT, Nesanet B, Dinka A (2008) Part II: Prevalences of major skin diseases in cattle, sheep and goats at Adama Veterinary Clinic, Oromia regional state, Ethiopia. *Revue de Médecine Vétérinaire* 159(8): 455-461.
18. NMSA (2013) Rainfall and temperature data of Bishoftu; National Metrological Service Agency, Addis Ababa, Ethiopia.
19. Aiello SE, Mays A (1998) *The Merck Veterinary Manual*. Whitehouse Station, NJ: Merck & Co. Inc 8, New York, United states, pp. 2187-2197.
20. Nicholson MJ, Butterworth MH (1986) A guide to condition scoring of zebu cattle.
21. Soulsby EJJ (1982) Helminths. Arthropods Protozoa Domesticated Animals *Transactions of The Royal Society of Tropical Medicine and Hygiene* 78(3): 291.
22. Urquhart GM, Armour J, Duncan JL, Dunn AM, Jennings FW (1996) *Veterinary parasitology* 2nd (edn), Black Well Science, Univ. Glas. Scotland, UK, pp. 181-188.
23. Beyecha K, Kumsa B, Beyene D (2014) Ectoparasites of goats in three agroecologies in central Oromia, Ethiopia. *Comparative Clinical Pathology* 23: 21-28.
24. Carter GR, Cole JR (2012) Diagnostic procedure in veterinary bacteriology and mycology. *Can Vet J* 33(6): 360.
25. Thrusfield M (2005) *Veterinary Epidemiology* 3rd (edn), Blackwell Publishing, UK.
26. Abebayehu T, Endris F, Berhanu M, Rahmeto A, Solomon M, et al. (2011) Study on the prevalence of ectoparasite infestation of ruminants in and around Kombolcha and damage to fresh goat pelts and wet blue (pickled) skin at Kombolcha tannery, Northeastern Ethiopia. *Ethiopian Veterinary Journal* 15(2): 87-101.
27. Onu SH, Shiferaw TZ (2013) Prevalence of ectoparasite infestations of cattle in Bench Maji zone, southwest Ethiopia. *Veterinary World* 6(6): 291-294.

28. Yacob HT, Yalew TA, Dinka AA (2008b) Part I: ectoparasite prevalences in sheep and in goats in and around Wolaita soddo, Southern Ethiopia. *Revue de Médecine Vétérinaire* 159(8): 450-454.
29. Chanie M, Negash T, Sirak A (2010) Ectoparasites are the major causes of various types of skin lesions in small ruminants in Ethiopia. *Trop Anim Health Prod* 42(6): 1103-1109.
30. Abebe R, Tatek M, Megersa B, Sheferaw D (2011) Prevalence of small ruminant ectoparasites and associated risk factors in selected districts of Tigray Region, Ethiopia. *Global Veterinaria* 7(5): 433-437.
31. Dawit T, Mulugeta A, Tilaye D, Mengistie T (2012) Ectoparasites of small ruminants presented at Bahir dar veterinary clinic, Northwest Ethiopia. *African J Agric Res* 7(33): 4669-4674.
32. Woldemeskel M, Taye G (2002) Prevalence of bovine dermatophilosis in a tropical highland region of Ethiopia. *Trop Anim Health Prod* 34(3): 189-194.
33. Kassaye E, Moser I, Woldemeskel M (2003) Epidemiological study on clinical bovine dermatophilosis in northern Ethiopia. *Dtsch Tierarztl Wochenschr* 110(10): 422-425.
34. Awad WS, Nadra-Elwgoud MI, El-Sayed AA (2008) Diagnosis and treatment of bovine, ovine and equine dermatophilosis. *Journal of Applied Sciences Research* 4(4): 367-374.
35. Beshahwured S (1991) Outbreak of lumpy skin disease in and around Wolliso. FVM, AAU, Debre-Zeit DVM Thesis.



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/AAPS.2023.02.555583](https://doi.org/10.19080/AAPS.2023.02.555583)

**Your next submission with Juniper Publishers
will reach you the below assets**

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission

<https://juniperpublishers.com/online-submission.php>