



Comprehensive Review of Dental Anomalies in Children with Cleft Lip and Palate: Prevalence, Etiology, and Management Strategies



Sara Mohamed Hamid*

Lecturer of Pediatric Dentistry, University of Khartoum, Sudan

Submission: March 27, 2025; Published: April 10, 2025

*Corresponding author: Sara Mohamed Hamid, Lecturer of Pediatric Dentistry, University of Khartoum, Sudan

Abstract

Abstract: Cleft lip and palate (CLP) are a common congenital malformation affecting the maxillofacial region, with prevalence varying globally. The etiology involves genetic and environmental factors leading to unsuccessful fusion during embryonic development. Dental anomalies are prevalent among CLP patients, including hypodontia, supernumerary teeth, tooth shape and size abnormalities, impaction, and malposition. These anomalies significantly impact oral function, aesthetics, and overall well-being. Early detection and multidisciplinary intervention are crucial for optimal management. Further research is needed to understand the complex etiology and improve treatment outcomes for CLP patients

Keywords: Cleft lip and palate; Dental anomalies; Prevalence; Etiology; Hypodontia; Teeth malformation

Abbreviations: CLP: Cleft lip and palate; CBCT: Cone-Beam Computed Tomography; CL: Cleft lip; P: Palate

Introduction

Cleft lip and/or palate (CL and/or P) is one of the most common congenital malformations in the maxillofacial region [1]. The prevalence throughout the world varies from 1/500 to 1/2500 live births. This variation largely depends on racial origin, ethnic background, and geographic location [2]. The aetiology of oral clefts is multifactorial in nature, with genetic and environmental factors contributing to its presence [3]. Clefts in the lip and palate result from the unsuccessful fusion of the embryonic medial nasal processes and maxillary processes and the medial nasal processes and palatine processes, respectively [4]. These fusions normally take place during the seventh to eleventh week of gestational age [5]. It can occur in isolation (non-syndromic) or be part of a wider series of birth anomalies or syndromes (syndromic) [1]. Cleft lip and/or palate malformations are divided into four main categories: palatal cleft lip and palate clefts, lip clefts, and lip and alveolar clefts [6]. According to the definition, palatal clefts do not extend to the maxillary alveolus [6]. Submucosal clefts are covered by mucosa and are also classified as a type of palatal cleft. Palatal clefts may involve both the soft and hard palates or only the soft palate [4]. Lip and palate clefts may extend bilaterally or unilaterally through the alveolar ridge and the lip to the hard

and soft palates [4]. Clefts of the palate are more common in girls, while clefts of the lip, with or without palatal involvement, are more common in boys. It is interesting to note that the left side is affected more often than the right [7]. The affinity of the unilateral cleft to the left side is not well understood [8].

Clinical Challenges

Children with cleft lip and palate face several problems, which include infant feeding problems due to an improper oral seal, swallowing and nasal regurgitation, multiple surgeries, visible facial scarring, nasolabial deformities, speech and hearing problems, and problems with appearance [9]. This may subject affected children to cruel teasing by their peers in school [9].

Dental Anomalies

Subjects affected with clefts commonly show various dental anomalies involving tooth number shape, size, structure, and position [10,11]. The degree of these dental anomalies varies according to gender, ethnicity, and type of the cleft [12]. These anomalies adversely impact the dentition, resulting in aesthetic problems, impairment of mastication, and improper phonation. Moreover, they can also make dental procedures more complicated

[13]. Children with cleft and their parents prioritize the surgical correction of the clefts and neglect their dental health, resulting in a higher prevalence of tooth decay, missing teeth, and overall poorer oral health [14]. Understanding dental anomalies in children with cleft lip and palate is essential for providing multidisciplinary care that addresses both the cleft and associated dental issues. It facilitates early intervention, improves oral function and psychosocial well-being, prevents secondary complications, and promotes long-term oral health and quality of life.

The objective of this paper is to provide a comprehensive review on the prevalence, types, etiology, and management strategies of dental anomalies observed in children with cleft lip and palate. The occurrence of dental anomalies in children with CLP is higher compared to the general population [15,16]. The association between these anomalies and the presence of clefts remains inadequately understood, with both genetic and environmental factors have been suggested to clarify this correlation [17,18]. This association may also arise from the close proximity of their anatomical structures as well as the concurrent timing of dental development and cleft formation. Notably, the occurrence of clefting coincides with odontogenesis, suggesting a potential interplay between these processes [19]. Some studies have confirmed that certain genes may contribute to both orofacial clefts and congenital dental anomalies [18,20,21]. Among the identified gene candidates implicated in the occurrence of clefts and congenital defects are *MSX1*, *PAX9*, and *IRF6* [22]. Furthermore, the effect of genetic risk factors may cause structural deficiencies in the embryonic oral tissues, leading to abnormalities in dental structure, shape, and number, followed by crowding, ectopic eruption, and malposition, thereby complicating access to oral hygiene [21]. The environmental risk factor that influences the occurrence of dental anomalies in the cleft population is early surgical repair of the cleft, which disturbs the tooth development process. These repair procedures are typically conducted during the initial years of life, coinciding with the critical phase of dental formation. [23].

Patients with various types of clefts show varying occurrences of dental anomalies. The prevalence is higher in individuals with cleft lip and cleft palate compared to other cleft types. Among these, bilateral clefts of the lip and palate have the highest frequency of anomalies, followed by left unilateral clefts of the lip and palate [24,25]. These anomalies are more common in permanent teeth, although it can also occur in primary dentition and it is more seen in the ipsilateral side of the cleft [26,27]. Moreover, the severity of these anomalies appears to be related to the severity of the cleft, i.e. the primary width of the cleft is positively correlated to the degree of central incisor rotation and enamel hypoplasia [28,29]. These anomalies may include anomalies of the number of teeth (hypodontia, supernumerary teeth), abnormalities of shape, dental eruption disorders, and disorders related to enamel mineralization [30]. The most prevalent dental anomalies observed in individuals with clefts are the absence of maxillary

lateral incisors, the presence of supernumerary teeth, and the absence of lower incisors [31]. In Northern Finland, cleft children were found to have a higher occurrence of dental anomalies (47%) compared to the general population, where the prevalence was 11.7%. Agenesis emerged as the most frequently observed dental anomaly in this group [32]. Akcam and colleagues examined the occurrence of various dental anomalies in the maxillary dental arch in different cleft groups and found that a significant proportion (96.7%) of individuals with a cleft had at least 1 dental anomaly. The most common anomaly observed was agenesis in the anterior region on the cleft side [23].

In a case-control study, a higher prevalence of enamel defect was found in children with cleft when compared with a control group. Furthermore, these defects were more commonly observed on the cleft side of the maxilla, with the central incisor being the tooth most frequently affected in this area [33]. A case-control investigation was carried out among Colombian children aged 5 to 12 years, comprising 210 subjects with non-syndromic cleft lip and palate and an equal number of healthy controls. The findings revealed that dental anomalies were most prevalent in cleft-affected children, particularly in those with bilateral cleft lip and palate, followed by left unilateral cleft lip and palate. Notably, these anomalies were predominantly located within the cleft area. Microdontia of the lateral maxillary incisors exhibited the highest prevalence, succeeded by rotations of the central maxillary incisors, agenesis of the lateral maxillary incisors, and supernumerary teeth [24].

Common observed dental anomalies

Hypodontia

The prevalence of hypodontia in children with cleft has been reported to range from 28% to 66% [26,34]. Permanent teeth are more often affected (52.7%) than primary (16.2%). The prevalence of hypodontia increases strongly with the severity of cleft [35]. Multiple hypodontia was found more frequently in the subjects with bilateral cleft lip and palate and those with unilateral cleft lip and palate [35]. Jamilian et al., in their study, did not reveal any significant difference between the genders in the prevalence of hypodontia, which differs from the healthy population because the female gender is prone to hypodontia [36]. The teeth most commonly affected by hypodontia are the maxillary lateral incisor, followed by the upper and lower second premolars [34,37], with the maxillary second premolar being the more frequently missing tooth [25,35]. The location of hypodontia can be inside and outside the cleft region, but it's more frequent on the cleft side of maxillary dentition [25,38]. When agenesis is found outside, this can suggest a genetic background and usually affects the contralateral incisor, or less often, the second premolar in the maxilla or in the mandible [39,40]. Other authors claimed the hypodontia may be due to local conditional effects of cleft, such as surgery for cleft palate would increase the risk of losing the tooth germs of permanent teeth such as the maxillary second premolars and/or

lateral incisors. Korolenkova et al. found primary periosteoplasty and reduced blood supply associated with palatal defects as a reason of agenesis of maxillary central and lateral incisors [41]. These findings suggest that the hypodontia in cleft patients is influenced by environmental factors [35]. Furthermore, primary bone grafting in cleft patients lowers the prevalence of hypodontia [42,43].

Supernumerary teeth

The prevalence of supernumerary teeth in cleft patients ranges from 4.6 to 42.0%, which is higher than in the general population [26,36]. It is more commonly found in the lateral incisor region adjacent to the cleft [44]. The prevalence of a supernumerary lateral incisor in patients with a left ranged from 5.1% to 22.1% [27]. In the study of Pradhan et al. (2020) conducted in Nepal, supernumerary teeth were mostly the maxillary laterals, both inside and outside the cleft. Aetiology of Supernumerary teeth has not been fully recognized, it may be explained by fragmentation of the dental lamina during cleft formation [27] or lengthening of the pre-canine section of the oral epithelium caused by cleft and thus an extension of the dental lamina, which can develop into a supernumerary tooth [25,45]. The division of the lateral incisor's tooth bud situated across the clefted nasopalatal sulcus may also lead to supernumerary teeth [25,45]. The clinical solution is usually the extraction of the additional tooth, but sometimes there are difficulties in differentiating the normal tooth from an additional tooth. A Cone-Beam Computed Tomography (CBCT) examination is obligatory for good clinical assessment and treatment planning in the case of supernumerary teeth [27].

Tooth shape and size anomalies

In patients with cleft the upper incisors are often affected with shape anomalies. The lateral incisor in the cleft area is often peg-shaped or hypoplastic [46-48]. In a sample of 90 patients (aged 4-20 years) affected by cleft, Rullo et al. found the upper lateral incisor microdontia revealed 5.6% of the examined individuals [46]. In a study by Tan et al., 12.5% of 60 examined patients had macrodontia [44]. An interesting finding is that posterior teeth can be bigger in size than in the healthy population, which suggests a multiple-teeth-size disorder in patients with cleft [47]. Jamilian et al. reported asymmetries in teeth dimensions comparing cleft and non-cleft side, where Maxillary central and lateral incisors were larger on the non-cleft side in the mesiodistal dimension compared with the cleft side. Upper central incisors and first molars are significantly larger mesiodistally on the non-cleft side [47]. In a retrospective study conducted by K uchler et al, taurodontism, which result failure or late invagination of Hertwig's epithelial root sheath, and lack of shift of the root furcation was found in 15.2% of the examined cleft patients [49]. Tooth shape and size anomalies cause asymmetry in the dental arch and poor aesthetic appearance. It demands orthodontic alignment followed by prosthetic restoration of hypoplastic or deformed teeth.

Tooth Impaction

Tooth impaction in patients with cleft can affect different teeth: upper incisors, canines, and premolars. Canines are the most commonly affected teeth [36]. The frequency of impaction varied highly in the literature, ranging from 0% to 58% [50]. Jamilian et al. in the Iranian population revealed the maxillary canines were the most commonly impacted teeth in both unilateral and bilateral cleft lip and palate patients [36]. In the literature both genetic and anatomical factors are described as etiological factors of tooth impaction in cleft patients. Narrowed clefted maxilla and lack of space may also cause impaction. In the cleft are a secondary bone grafting should improve eruption of an upper lateral incisor and canine; therefore, lack of exact treatment can be the reason of upper canines impaction. A Polish study by Pastuszek et al. did not confirm any relationship between bone grafting, maxillary expansion or extraction of non-resorbed primary canine, and prevalence of upper canine impaction [51]. In non-cleft patients, there is a positive correlation between the occurrence of hypodontia or reduced size of maxillary lateral incisors and canine impaction, but this was not confirmed in the population with cleft [51]. Patients who suffer from tooth impaction may demand complicated surgical and orthodontic procedures to achieve the tooth and align the dental arch.

Teeth Malposition

In children, the maxillary teeth are more frequently in an abnormal position. A narrow and short upper arch means a lack of space, which ultimately leads to crowding [44]. Rotations, ectopic eruptions, and transpositions are the most common. Tan et al. found the rotations of central incisors is a common problem in the area of the cleft with a prevalence of 86.7% and a significantly greater frequency of rotations was found in females [44]. Transposition of maxillary canine and first premolars was found in 5.5% of bilateral, 8% of right, and 3.3% of left unilateral clefts in the study of Eslami et al. [12]. Common teeth malposition in cleft patients is palatal eruption of upper lateral incisors and upper second premolars due to lack of space and class III tendency [52]. All described teeth displacement demands comprehensive orthodontic fixed appliance therapy.

Conclusion

This review highlights the prevalence and significance of dental anomalies in children with cleft lip and palate (CLP). These anomalies, including hypodontia, supernumerary teeth, tooth shape and size abnormalities, impaction, and malposition, are more common and severe in CLP patients compared to the general population. Clinically, early detection and intervention for dental anomalies in CLP patients are vital for optimizing oral function and aesthetics. Multidisciplinary care involving various dental specialists is necessary. Addressing dental anomalies in CLP children is paramount for enhancing their oral health, psychosocial well-being, and overall quality of life. Research should focus on

understanding the complex aetiology of these anomalies, tracking dental development longitudinally, and exploring innovative treatment modalities to improve outcomes.

References

1. Owens JR, Jones JW, Harris F (1985) Epidemiology of facial clefting. *Arch Dis Child* 60(6): 521-524.
2. Molina-Solana R (2013) Current concepts on the effect of environmental factors on cleft lip and palate. *Int J Oral Maxillofac Surg* 42(2): 177-184.
3. Yazdy MM (2007) Priorities for future public health research in orofacial clefts. *Cleft Palate Craniofac J* 44(4): 351-357.
4. Poswillo D (1988) The aetiology and pathogenesis of craniofacial deformity. *Development* 103 Suppl: 207-212.
5. Rice DP (2005) Craniofacial anomalies: from development to molecular pathogenesis. *Curr Mol Med* 5(7): 699-722.
6. Eppley BL (2005) The spectrum of orofacial clefting. *Plast Reconstr Surg* 115(7): 101e-114e.
7. Kohli SS, Kohli VS (2012) A comprehensive review of the genetic basis of cleft lip and palate. *J Oral Maxillofac Pathol* 16(1): 64-72.
8. Daskalogiannakis J (1998) Unilateral cleft lip with or without cleft palate and handedness: is there an association? *Cleft Palate Craniofac J* 35(1): 46-51.
9. Mossey P, Little J (2009) Addressing the challenges of cleft lip and palate research in India. *Indian J Plast Surg* 42Suppl: S9-S18.
10. Ranta R, Stegars T, Rintala AE (1983) Correlations of hypodontia in children with isolated cleft palate. *Cleft Palate J* 20(2): 163-165.
11. da Silva Dalben G (2008) Conjoined twins with mirror-image cleft lip and palate: case report in Brazil. *Cleft Palate Craniofac J* 45(3): 315-318.
12. Eslami N (2013) Prevalence of dental anomalies in patients with cleft lip and palate. *J Craniofac Surg* 24(5): 1695-1698.
13. Hardin-Jones MA, Jones DL (2005) Speech production of preschoolers with cleft palate. *Cleft Palate Craniofac J* 42(1): 7-13.
14. Al-Wahadni A, Alhaja EA, Al-Omari MA (2005) Oral disease status of a sample of Jordanian people ages 10 to 28 with cleft lip and palate. *Cleft Palate Craniofac J* 42(3): 304-308.
15. Germec Cakan D (2018) Dental Anomalies in Different Types of Cleft Lip and Palate: Is There Any Relation? *J Craniofac Surg* 29(5): 1316-1321.
16. Lourenco Ribeiro L (2003) Dental anomalies of the permanent lateral incisors and prevalence of hypodontia outside the cleft area in complete unilateral cleft lip and palate. *Cleft Palate Craniofac J* 40(2): 172-175.
17. Tannure PN (2012) Prevalence of dental anomalies in nonsyndromic individuals with cleft lip and palate: a systematic review and meta-analysis. *Cleft Palate Craniofac J* 49(2): 194-200.
18. Johnson DB (1967) Some observations on certain developmental dento-alveolar anomalies and the stigmata of cleft. *Dent Pract Dent Rec* 17(12): 435-43.
19. Mangione F (2018) Cleft palate with/without cleft lip in French children: radiographic evaluation of prevalence, location and coexistence of dental anomalies inside and outside cleft region. *Clin Oral Investig* 22(2): 689-695.
20. Vieira AR (2008) Candidate gene/loci studies in cleft lip/palate and dental anomalies finds novel susceptibility genes for clefts. *Genet Med* 10(9): 668-674.
21. Howe BJ (2017) Dental Decay Phenotype in Nonsyndromic Orofacial Clefting. *J Dent Res* 96(10): 1106-1114.
22. Phan M (2016) Tooth agenesis and orofacial clefting: genetic brothers in arms? *Hum Genet* 135(12): 1299-1327.
23. Akcam MO (2010) Dental anomalies in individuals with cleft lip and/or palate. *Eur J Orthod* 32(2): 207-213.
24. Yezioro-Rubinsky S (2020) Dental Anomalies in Permanent Teeth Associated with Nonsyndromic Cleft Lip and Palate in a Group of Colombian Children. *Cleft Palate Craniofac J* 57(1): 73-79.
25. Ranta R (1986) A review of tooth formation in children with cleft lip/palate. *Am J Orthod Dentofacial Orthop* 90(1): 11-18.
26. Howe BJ (2015) Spectrum of Dental Phenotypes in Nonsyndromic Orofacial Clefting. *J Dent Res* 94(7): 905-912.
27. Lasota A (2022) The Prevalence and Morphology of Supernumerary Teeth in Children With Nonsyndromic Cleft Lip and Palate. *Cleft Palate Craniofac J* 59(7): 867-872.
28. Schroeder DC, Green LJ (1975) Frequency of dental trait anomalies in cleft, sibling, and noncleft groups. *J Dent Res* 54(4): 802-807.
29. Jabbari F (2016) Correlations between initial cleft size and dental anomalies in unilateral cleft lip and palate patients after alveolar bone grafting. *Ups J Med Sci* 121(1): 33-37.
30. Ribeiro LL (2002) Dental development of permanent lateral incisor in complete unilateral cleft lip and palate. *Cleft Palate Craniofac J* 39(2): 193-196.
31. Schwartz JP (2014) Prevalence of dental anomalies of number in different subphenotypes of isolated cleft palate. *Dental Press J Orthod* 19(1): 55-59.
32. Lehtonen V (2015) Dental anomalies associated with cleft lip and palate in Northern Finland. *Eur J Paediatr Dent* 16(4): 327-332.
33. Shen CA, Guo R, Li W (2019) Enamel defects in permanent teeth of patients with cleft lip and palate: a cross-sectional study. *J Int Med Res* 47(5): 2084-2096.
34. Al Jamal GA, Hazza'a AM, Rawashdeh MA (2010) Prevalence of dental anomalies in a population of cleft lip and palate patients. *Cleft Palate Craniofac J* 47(4): 413-420.
35. Suzuki A (2017) A Longitudinal Study of the Presence of Dental Anomalies in the Primary and Permanent Dentitions of Cleft Lip and/or Palate Patients. *Cleft Palate Craniofac J* 54(3): 309-320.
36. Jamilian A (2015) Hypodontia and supernumerary and impacted teeth in children with various types of clefts. *Am J Orthod Dentofacial Orthop* 147(2): 221-225.
37. Harris EF, Hullings JG (1990) Delayed dental development in children with isolated cleft lip and palate. *Arch Oral Biol* 35(6): 469-473.
38. Matern O (2012) Left-sided predominance of hypodontia irrespective of cleft sidedness in a French population. *Cleft Palate Craniofac J* 49(3): e1-5.
39. Bartzela TN (2013) Tooth agenesis patterns in unilateral cleft lip and palate in humans. *Arch Oral Biol* 58(6): 596-602.
40. Mikulewicz M (2014) Prevalence of second premolar hypodontia in the Polish cleft lip and palate population. *Med Sci Monit* 20: 355-360.
41. Korolenkova MV, Starikova NV, Udalova NV (2019) The role of external aetiological factors in dental anomalies in non-syndromic cleft lip and palate patients. *Eur Arch Paediatr Dent* 20(2): 105-111.
42. Hellquist R (1979) Dental abnormalities in patients with alveolar clefts, operated upon with or without primary periosteoplasty. *Eur J Orthod* 1(3): 169-180.

43. Helms JA, Speide TM, Denis KL (1987) Effect of timing on long-term clinical success of alveolar cleft bone grafts. *Am J Orthod Dentofacial Orthop* 92(3): 232-240.
44. Tan ELY (2018) Secondary Dentition Characteristics in Children with Nonsyndromic Unilateral Cleft Lip and Palate: A Retrospective Study. *Cleft Palate Craniofac J* 55(4): 582-589.
45. Kim NY, Baek SH (2006) Cleft sidedness and congenitally missing or malformed permanent maxillary lateral incisors in Korean patients with unilateral cleft lip and alveolus or unilateral cleft lip and palate. *Am J Orthod Dentofacial Orthop* 130(6): 752-758.
46. Rullo R (2015) Prevalence of dental anomalies in children with cleft lip and unilateral and bilateral cleft lip and palate. *Eur J Paediatr Dent* 16(3): 229-232.
47. Antonarakis GS, Tsiouli K, Christou P (2013) Mesiodistal tooth size in non-syndromic unilateral cleft lip and palate patients: a meta-analysis. *Clin Oral Investig* 17(2): 365-377.
48. Walker SC (2009) Abnormal tooth size and morphology in subjects with cleft lip and/or palate in the north of England. *Eur J Orthod* 31(1): 68-75.
49. Kuchler EC (2011) Side of dental anomalies and taurodontism as potential clinical markers for cleft subphenotypes. *Cleft Palate Craniofac J* 48(1): 103-108.
50. Westerlund A (2014) What factors are associated with impacted canines in cleft patients? *J Oral Maxillofac Surg* 72(11): 2109-2014.
51. Pastuszak PI (2020) Dunin-Wilczynska, and A. Lasota, Frequency of Additional Congenital Dental Anomalies in Children with Cleft Lip, Alveolar and Palate. *J Clin Med* 9(12): 3813.
52. Marzouk T (2021) Association between Dental Anomalies and Orofacial Clefts: A Meta-analysis. *JDR Clin Trans Res* 6(4): 368-381.



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/ADOH.2025.18.555984](https://doi.org/10.19080/ADOH.2025.18.555984)

**Your next submission with Juniper Publishers
will reach you the below assets**

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission
<https://juniperpublishers.com/online-submission.php>