



Recent Advances in Synthetic Applications of Polyvinylpyrrolidone Supported Reagents and Catalysts



Masoud Mokhtary*

Department of Chemistry, Rasht Branch, Islamic Azad University, Iran

Submission: August 23, 2018; Published: November 05, 2018

*Corresponding author: Masoud Mokhtary, Department of Chemistry, Rasht Branch, Islamic Azad University, Iran

Abstract

This review summarized recent progresses in the application of polyvinylpyrrolidone supported reagents and catalysts in organic synthesis.

Keywords: Polyvinylpyrrolidone; Supported reagents; Supported catalysts; Multi-component reactions


Introduction

The use of solid-supported reagents and catalysts in solution-phase chemistry has emerged as a leading strategy that exploits the advantages of both solid- and solution-phase synthesis. The approach essentially combines the benefits of product isolation and purification in solid-phase synthesis with the high-speed development and flexible choice of chemistry from the vast repertoire of solution phase organic reactions. The organic molecules synthesis using polymer-supported reagents and catalysts is highly attractive because the work-up involves only simple filtration and evaporation of the solvent [1]. Polyvinylpyrrolidone (PVP) is an amorphous polymer having broad applications in biomedical field due to its special properties such as low toxicity and good solubility in water and most organic solvents, good adhesion characteristics, and great physiological compatibility [2]. Also, PVP has good biocompatibility and has been applied for many years as a biomaterial or additive to drug compositions, e.g. as a blood plasma expander [3].

Polyvinylpyrrolidone (PVPP, crospovidone, or crospolividone) is a highly cross-linked polyvinylpyrrolidone (PVP). The cross-linked form of polyvinylpyrrolidone is insoluble in water, though it still absorbs water and swells very rapidly generating a swelling force. This property makes PVPP useful as a disintegrant in pharmaceutical tablets [2]. Polyvinylpyrrolidone shows a strong binding affinity to small molecules. Furthermore, its iodine complex, povidon-iodine, is widely used as an anti-infective agent in clinical treatments [4].

Polyvinylpyrrolidone Supported Reagents


A range of polyvinylpyrrolidone-supported reagents has been developed for applications in organic synthesis. In general, these reagents are employed in stoichiometric excess to drive the reaction to completion. Simple filtration removes the spent resin from the reaction solution and, thus, eliminates the need for any time-consuming chromatographic work-up.



Scheme 1: Ring opening of epoxides and ring opening dimerization of thiiranes by PVP-I₂.

Iranpoor and coworkers [5] have prepared iodine supported on polyvinylpyrrolidone (betadine) as catalysts for ring opening of epoxides and as reagent for ring opening dimerization of thiiranes in alcohols, water and acetic acid (Scheme 1). In this

report, the reaction of *R*-(+)-styrene oxide with I₂ supported on PVP in methanol was found to be very stereospecific and the product isolated in 93% ee.




R = alkyl, aryl

R' = CH₂OH, COR, COAr, NO₂, CH, COOEt

Scheme 2: Bromination of alkenes using PVP-Br₂.

In another research, Lakouraj et al. [6] described the preparation of polyvinylpyrrolidone-bromine complex (PVP-Br₂) as a mild and convenient reagent for selective bromination of alkenes (Scheme 2) [6].


Selective oxidation of benzyl alcohol in the presence of 2-phenylethanol was also achieved at room temperature in the presence of PVP-Br₂ (Scheme 3).




Scheme 3: Selective oxidation of benzyl alcohol in the presence of 2-phenylethanol by PVP-Br₂.

In the next research, Lakouraj and Mokhtary have developed a convenient method for deprotection and direct oxidative deprotection of silylethers to the corresponding hydroxy and carbonyl compounds using polyvinylpyrrolidone-

bromine complex (PVPP-Br₂) (Scheme 4) [7]. Selective oxidative deprotection of benzylic silyl ethers in the presence of primary aliphatic alcohols was also achieved at room temperature.



Scheme 4: Deprotection and direct oxidative deprotection of silylethers by PVPP-Br₂.




R = OH, OMe, N(Me)₂, N(Et)₂

X = Electron releasing groups

Scheme 5: bromination of electron-rich aromatic compounds by PVPP-Br₂.

Also, PVPP-Br₂ has been used for bromination of electron-rich aromatic compounds [8]. The reaction proceeded smoothly with phenols and *N, N*-alkylated amines to afford the corresponding


mono brominated product in good yields at ambient temperature (Scheme 5).




Scheme 6: Synthesis of benzylic bromides by PVPP-Br₂.

Furthermore, Mokhtary and Lakouraj synthesized benzylic bromides in high yields by the reaction of the corresponding alcohols with cross-linked polyvinylpyrrolidone-bromine complex (PVPP-Br₂) in the presence of hexamethyldisilane in chloroform at reflux condition (Scheme 6) [9]. Selective conversion of benzyl alcohol to benzyl bromide in the presence of 2-phenylethanol was also achieved.

Surya Prakash et al. have prepared solid polyvinylpyrrolidone-hydrogen peroxide complex and used as solid hydroxylating reagent [10]. This solid hydrogen peroxide is found to be much safer, convenient and efficient reagent system for the ipso-hydroxylation of arylboronic acids to the corresponding phenols in high yields at a faster rate (Scheme 7). The versatility of the reagent has been further expanded for the one-pot synthesis of halophenols.



Scheme 7: Ipso-hydroxylation of arylboronic acids by PVPP-H₂O₂.




Scheme 8: Oxygenation of sulfides to sulfoxides, and oxidation of aldehydes to carboxylic acids by PVPP-POCl₃ in the presence of hydrogen peroxide.


Lakouraj et al., [11] have demonstrated polyvinylpyrrolidone-phosphorous oxychloride as a versatile polymeric Vilsmeier reagent that exhibits excellent selectivity for oxygenation of sulfides to sulfoxides, and oxidation of aldehydes to carboxylic acids in the presence of hydrogen peroxide under mild reaction conditions (Scheme 8) [11]. This polymeric Vilsmeier reagent was

found to retain its activity after months and is stable in a glass bottle at room temperature.

Iranpoor et al. [12] have reported dinitrogen tetraoxide supported on polyvinylpyrrolidone (PVP-N₂O₄) as a nitrosating and coupling agent for thiols and selective oxidation of sulfides to sulfoxides and disulfids to thiosulfonates (Scheme 9) [12].




Scheme 9: Coupling of thiols and selective oxidation of sulfides to sulfoxides and disulfides to thiosulfonates by PVP-N₂O₄.



Scheme 10: Dehydration of a variety of aldoximes by PVP-SOCl₂.

Tamami and Kiasat have reported [13] a polyvinylpyrrolidone-thionyl chloride complex by the reaction of thionyl chloride with two equivalents of polyvinylpyrrolidone in dichloromethane at 0°C.

The polymer-bound complex I was used for the rapid dehydration of a variety of aldoximes to produce the corresponding nitriles in high yields (Scheme 10).



Scheme 11: Ring opening of epoxides to afford β -chlorohydrins using PVP-SOCl₂.

Also, Tamami et al. [14] have reported rapid ring opening of epoxides to afford β -chlorohydrins with cross-linked polyvinylpyrrolidone/thionyl chloride complex (PVP-SOCl₂), under mild reaction condition in high yields (Scheme 11).

Polyvinylpyrrolidone Supported Lewis Acid Catalysts


Replacement of conventional, toxic and unstable Lewis acidic catalysts with eco-friendly reusable solid acid catalysts is an essential requirement in the development of green chemistry. For example, boron trifluoride is widely used in organic syntheses as a Lewis acid. However, boron trifluoride is highly water sensitive, irritant, and has to be used in a carefully dried apparatus. Moreover, all work must be carried out in an efficient fume hood, and its recovery from the reaction mixture results in a main

source of waste, which on an industrial scale is environmentally unacceptable.


Mokhtary et al. have reported several papers on the application of the polyvinylpyrrolidone-bound boron trifluoride (PVPP-BF₃) as a stable polymeric Lewis acid catalyst in some organic reactions such as synthesis of amides by the reaction of nitriles with benzhydrol and tertiary alcohols or *tert*-butyl acetate via Ritter reaction (Scheme 12) [15, 16], the acylation of alcohols, phenols and trimethylsilyl ethers with acetic anhydride (Scheme 13) [17], the synthesis of coumarins via Pechmann condensations of phenols with ethyl acetoacetate (Scheme 14) [18], the synthesis of 14-aryl-14*H*-dibenzo [a,j] xanthenes and bis(naphthalen-2-yl-sulfane) derivatives (Scheme 15) [19], the synthesis of 1,8-dioxooctahydroxanthenes and 1,8-dioxodecahydroacridines via condensation of aromatic aldehydes and dimedone in

acetonitrile at room temperature, and aromatic aldehydes, dimedone, and aromatic amines in acetonitrile at 80 °C respectively (Scheme 16) [20], the oxidation of aldehydes to carboxylic acids and oxidative esterification of benzaldehydes in the presence of 35% hydrogen peroxide (Scheme 17) [21] and the


oxidation of sulfides to sulfones in the presence of 35% hydrogen peroxide at room temperature (Scheme 18) [22]. Excellent yields, easy work-up and reusability and stability of the catalyst are some advantages of these methods.




Scheme 12: Synthesis of amides by the reaction of nitriles with benzhydrol and tertiary alcohols or tert-butyl acetate using PVPP-BF₃.




Scheme 13: Acylation of alcohols, phenols and trimethylsilyl ethers using PVPP-BF₃.



Scheme 14: Condensations of phenols with ethyl acetoacetate by PVPP-BF₃.




Scheme 15: The synthesis of 14-aryl-14H-dibenzo[a,j]xanthenes and bis(naphthalen-2-yl-sulfane) derivatives by PVPP-BF₃.




Scheme 16: synthesis of 1,8-dioxooctahydroxanthenes and 1,8-dioxodecahydroacridines using PVPP-BF₃.



Scheme 17: Oxidation of aldehydes to carboxylic acids and oxidative esterification of benzaldehydes by PVPP-BF₃/H₂O₂.




Scheme 18: Oxidation of sulfides to sulfones in the presence of PVPP-BF₃/H₂O₂.



Scheme 19: Ring-opening of epoxides by various alcohols and regioselective conversion of epoxides to β -azidoalcohols by PVP-GaCl₃.

Pourali and et al. prepared cross-linked polyvinylpyrrolidone supported GaCl₃ for efficient and regioselective ring-opening reaction of epoxides by various alcohols under solvent-free conditions at room temperature (Scheme 19) [23]. Furthermore, regioselective conversion of epoxides to β -azidoalcohols was

accomplished by sodium azide in MeOH in the presence of GaCl₃/PVP at room temperature. GaCl₃/PVPP is a non-hygroscopic and recoverable catalyst and is easily separated from reaction mixture by a simple filtration and reused repeatedly.




Scheme 20: Synthesis of chromenylphenylpropanone derivatives using PVPP-SbCl₃.

In the next research, an efficient synthesis of chromenylphenylpropanone derivatives as warfarine-like analogues was developed by the Michael addition of 4-hydroxycoumarin to α , β -unsaturated compounds in the presence of polyvinylpyrrolidone supported antimony (III) chloride (PVPP-SbCl₃) as a new polymeric Lewis acid catalyst in chloroform at reflux conditions (Scheme 20) [24].

accomplished by sodium azide in MeOH in the presence of GaCl₃/PVP at room temperature. GaCl₃/PVPP is a non-hygroscopic and recoverable catalyst and is easily separated from reaction mixture by a simple filtration and reused repeatedly.

Polyvinylpyrrolidone Supported Brønsted Acidic Catalysts



Scheme 21: Trimethylsilylation and formylation of alcohols by PVPP-HBr₃.


Polymer-supported Brønsted acidic catalysts have gained considerable importance due to their low cost, high efficiency, easy

work-up, and reusability. Ghorbani-Choghamarani et al. reported trimethylsilylation and formylation of alcohols in the presence of


polyvinylpolypyrrolidonium tribromide in acetonitrile at room temperature (Scheme 21) [25].

Khaksar et al. have prepared polyvinylpolypyrrolidone-supported triflic acid (PVPP-OTf) as an environmentally friendly

and efficient catalyst for the synthesis of bis-indolyl methane derivatives by the reaction of indole or *N*-methyl indole with aldehydes (Scheme 22) [26].



Scheme 22: synthesis of bis-indolyl methane derivatives by PVPP-OTf.




Scheme 23: Synthesis of quinoxaline derivatives using PVPP-OTf.

Furthermore, PVPP-OTf was found to be useful as a recyclable heterogeneous catalyst for the rapid synthesis of quinoxaline derivatives (Scheme 23) [27].

Also, polyvinylpolypyrrolidone-supported triflic acid has been


used as a recyclable catalyst for synthesis of a series of 7-hydroxy-10-aryl-10H-indeno[1,2-*b*]chromen-11-one derivatives, 13-phenyl indeno[1,2-*b*]naphtho[1,2-*e*]pyran-12(13H)-one and 12-phenyl-12H-indeno[1,2-*b*]naphtho[3,2-*e*]pyran-5,11, 13-triones (Scheme 24) [28].




Scheme 24: Synthesis of 7-hydroxy-10-aryl-10H-indeno[1,2-*b*]chromen-11-one derivatives, 13-Phenyl indeno[1,2-*b*]naphtho[1,2-*e*]pyran-12(13H)-one and 12-phenyl-12H-indeno[1,2-*b*]naphtho[3,2-*e*]pyran-5,11, 13-triones by PVPP-OTf.

Nitration of the substituted phenols (Scheme 25) [29] and oxidation of urazoles and bis-urazoles to the corresponding triazolinediones (Scheme 26) [30] were reported by Nikoorazm et al. and Ghorbani-Choghamarani et al. respectively in dichlorometh-


ane at room temperature using supported nitric acid on polyvinylpyrrolidone as an efficient, environmentally friendly, mild catalyst.



Scheme 25: Nitration of the substituted phenols by PVP-HNO₃.



Scheme 26: Oxidation of urazoles and bis-urazoles using PVP-HNO₃.




Scheme 27: Chemo selective oxidation of sulfides to sulfoxides using PVP-HNO₃.


Furthermore, Zolfigol et al. was described chemo selective oxidation of sulfides to sulfoxides using PVP-HNO₃ in the presence of a catalytic amount of KBr or NaBr (Scheme 27) [31].

The synthesis of xanthenes derivatives including 1,8-diox-


ooctahydroxanthenes, 14-aryl-14H-dibenzo[a,j] xanthenes, and 12-aryl-8,9,10,12-tetrahydrobenzo[a]xanthen-11-ones reported by Shirini et al, using *O*-sulfonated poly(4-vinylpyrrolidonium) chloride [PVP-SO₃H]Cl as a polymeric solid acid catalyst (Scheme 28) [32].



Scheme 28: Synthesis of xanthenes derivatives including 1,8-dioxooctahydroxanthenes, 14-aryl-14H-dibenzo[a,j] xanthenes, and 12-aryl-8,9,10,12-tetrahydrobenzo[a]xanthen-11-ones by [PVP-SO₃H]Cl.




Scheme 29: Synthesis of hexahydroquinolines by [PVPP-SO₃H]Cl.




Scheme 30: Synthesis of dihydropyrimidinones and octahydroquinazolin-2,5-diones by [PVPP-SO₃H]Cl.

Also, polyvinylpyrrolidone supported chlorosulfonic acid ([PVPP-SO₃H]Cl) was evaluated by Mokhtary et al, as a recoverable catalyst for the one-pot synthesis of hexahydroquinolines (Scheme 29) [33], dihydropyrimidinones and octahydroquinazolin-2,5-diones (Scheme 30) [34].

In another research, Abedini et al. have described a green approach for the promotion of the synthesis of Hantzsch products using polyvinylpyrrolidinium perchlorate ([PVPH]ClO₄) as a new modified polymeric catalyst (Scheme 31) [35].




Scheme 31: Synthesis of Hantzsch products using [PVPH]ClO₄.



Scheme 32: Synthesis of 2-amino-3-cyano-tetrahydro-4H-benzopyran derivatives by PVP-H]H₂PO₄.

In the next research, Shirini et al., have introduced polyvinylpyrrolidonium hydrogen phosphate ([PVP-H] H_2PO_4) as a heterogeneous, and reusable catalyst for the synthesis of 2-amino-3-cyano-5-oxo-5,6,7,8-tetrahydro-4*H*-benzopyrans (Scheme 32) [36].

In another study, sulfuric acid-modified polyvinylpyrrolidone ([PVP- SO_3H] H_2SO_4) was prepared by Safaei et al. as an efficient reusable polymeric catalyst for the one-pot multi component synthesis of acridinedione derivatives as an important class of heterocyclic compounds (Scheme 33) [37].




Scheme 33: Synthesis of acridinedione derivatives by [PVP- SO_3H] H_2SO_4 .


Polyvinylpyrrolidone Supported Pd Catalyst

A series of poly(*N*-vinyl-2-pyrrolidone) immobilized Pd nanoparticles (PVP-Pd) with varying particle size have prepared by Li et al., [38] using the stepwise growth reaction. The effect of Pd particle size on the Suzuki reaction between phenylboronic acid and iodobenzene was investigated by the use of four Pd catalysts (Scheme 34). The catalytic activity of the Pd nanoparticles expressed in terms of the initial turnover frequency (moles of the biphenyl product per mole of total surface Pd atoms per min) was

found to be in the order of Pd (3.9nm) > Pd (3.0nm) ~ Pd (5.2nm) > Pd (6.6nm), indicating that surface Pd atoms do not all have the same reactivity in this reaction. The general trend of increased catalytic activity with the decrease in the particle size suggests that the low-coordination number vertex and edge atoms on the particle surface are active sites for the Suzuki reaction. The lower catalytic activity for the smallest Pd nanoparticles may be due to stronger adsorption of the reaction intermediates on the particle surface, in which the strongly adsorbed species act as a poison to the reaction thereby decreasing the rate of the reaction.



Scheme 34: Reaction between phenylboronic acid and iodobenzene by PVP-Pd NPs.



Scheme 35: Synthetic strategy for the preparation of PVP-grafted silica with $PdCl_2$ catalyst.

In another work, Tamami et al., [39] was prepared a catalytic system based on palladium nanoparticles supported on poly(*N*-vinylpyrrolidone) grafted silica (Scheme 35).

out to obtain the heterogeneous catalytic system. Transmission electron microscopy (TEM) image showed that palladium dispersed through the support in nanometer size (Figure 1).

The complexation of PVP-grafted silica with $PdCl_2$ was carried





Figure 1: Transmission electron microscopy (TEM) image of palladium supported on polyN-vinylpyrrolidone)-grafted silica [39].



Scheme 36: Cross-coupling reactions of aryl iodides, bromides and also chlorides with olefinic compounds by PVP-grafted silica with PdCl₂ catalyst.

This catalytic system exhibited excellent activity in cross-coupling reactions of aryl iodides, bromides and also chlorides with olefinic compounds in Heck-Mizoraki reactions in short reaction time and high yields (Scheme 36).

Conclusion

Polymer-supported reagents and catalysts have emerged as important tools for the rapid preparation of chemical compounds in solution-phase. Clean methodologies, easy preparation of the catalysts, simple work-up procedures, good to high yields, environmentally friendly and reusable catalysts are some advantages of polyvinylpyrrolidone supported catalysts and reagents in organic multi-step synthesis. Further progresses in the development of new PVP-bound reagents, and catalysts will continue to attract innovative application of this strategy in multi-component synthesis.

Acknowledgements

Financial support by Rasht Branch, Islamic Azad University is gratefully acknowledged.

References

- Bhattacharyya S (2000) Polymer-supported reagents and catalysts: recent advances in synthetic applications. *Comb Chem High Throughput Screen* 3(2): 65-92.
- Bühler V (2005) In: *Polyvinylpyrrolidone excipients for pharmaceuticals: povidone, crospovidone and copovidone*. Springer, Berlin Heidelberg New York, USA, pp. 1-254.
- Teodorescu M, Bercea M (2015) Poly(vinylpyrrolidone)-a versatile polymer for biomedical and beyond medical applications. *Polym Plast Technol Eng* 54(9): 923-943.
- Liu C, Cui S, Wang Z, Zhang X (2005) Single-chain mechanical property of poly(N-vinyl-2-pyrrolidone) and interaction with small molecules. *J Phys Chem B* 109:14807-14812.
- Iranpoor N, Tamami B, Niknam Kh (1997) Iodine and iodine supported on polyvinylpyrrolidone as catalysts and reagents for alcoholysis, hydrolysis and acetolysis of epoxides and thiiranes. *Can J Chem* 75(12): 1913-1919.
- Lakouraj MM, Tajbakhsh M, Mokhtary M (2005) Poly(vinylpyrrolidone)-bromine complex; a mild and efficient reagent for selective bromination of alkenes and oxidation of alcohols. *J Chem Res* 2005(8): 481-483.
- Lakouraj M M, Mokhtary M (2007) Polyvinylpyrrolidone-bromine complex, mild and efficient polymeric reagent for selective deprotection and oxidative deprotection of silylethers. *Lett Org Chem* 4(1): 64-67.
- Mokhtary M, Lakouraj M M (2011) Polyvinylpyrrolidone-bromine complex: mild and efficient polymeric reagent for bromination of activated aromatic compounds. *Chin Chem Lett* 22(1): 13-17.
- Mokhtary M, Lakouraj M M (2012) Polyvinylpyrrolidone-bromine complex: an efficient polymeric reagent for selective preparation of benzyl bromides in the presence of hexamethyldisilane. *Bull Chem Soc Ethiop* 26(2): 305-309.

10. Surya Prakash GK, Chacko S, Panja Ch, Thomas TE, Gurung L, et al. (2009) Regioselective synthesis of phenols and halophenols from arylboronic acids using solid poly(N-vinylpyrrolidone)/hydrogen peroxide and poly(4-vinylpyridine)/hydrogen peroxide complexes. *Adv Synth Catal* 351: 1567-1574.
11. LakourajMM, AghajaniB, MokhtaryM (2010) Polyvinylpolypyrrolidone-supported hydrogen peroxide for selective oxidation of aldehydes to carboxylic acids and sulfides to sulfoxides. *Phosphorus Sulfur Silicon Relat Elem* 185(12): 2393-2401.
12. Iranpoor N, Firouzabadi H, Pourali A R (2002) Dinitrogen tetroxide supported on polyvinylpyrrolidone (PVP-N₂O₄): a new nitrosating and coupling agent for thiols and selective oxidant for sulfides and disulfides. *Tetrahedron* 58(25): 5179-5184.
13. Tamami B, Kiasat A R (1999) Dehydration of aldoximes to nitriles using a polymeric reagent. *J Chem Res(S)* 444-445.
14. Tamami B, Ghazi I, Mahdavi H (2002) polyvinylpyrrolidone/thionylchloride complex as a new polymeric reagent for facile conversion of epoxides to β-chlorohydrins. *Synth Commun* 32(24): 3725-3731.
15. Lakouraj MM, Mokhtary M (2009) Polyvinylpolypyrrolidone-supported boron trifluoride: a high-loaded, polymer-supported Lewis acid for the Ritter reaction. *Monatsh Chem* 140: 53-56.
16. Mokhtary M, Najafzadeh F (2012) Polyvinylpolypyrrolidone-supported boron trifluoride; highly efficient catalyst for the synthesis of N-tert-butyl amides. *E-J Chem* 9(2): 576-582.
17. Mokhtary M, Qandalee M, Najafzadeh F (2012) Polyvinylpolypyrrolidone-bound boron trifluoride: A highly efficient catalyst for acylation of alcohols, phenols and trimethylsilyl ethers by acetic anhydride. *CR Chim* 15(5): 389-393.
18. Mokhtary M, Najafzadeh F (2012) Polyvinylpolypyrrolidone-bound boron trifluoride (PVPP-BF₃); a mild and efficient catalyst for synthesis of 4-methyl coumarins via the Pechmann reaction. *C R Chim* 15(6): 530-532.
19. Mokhtary M, Refahati S (2013) Polyvinylpolypyrrolidone-supported boron trifluoride (PVPP-BF₃): Mild and efficient catalyst for the synthesis of 14-aryl-14H-dibenzo[a,j] xanthenes and bis(naphthalen-2-yl-sulfane) derivatives. *Dyes Pigm* 99(2): 378-381.
20. Mokhtary M, Mirfarjood Langroudi S A (2014) Polyvinylpolypyrrolidone-supported boron trifluoride: a mild and efficient catalyst for the synthesis of 1,8-dioxooctahydroxanthenes and 1,8-dioxodecahydroacridines. *Monatsh Chem* 145: 1489-1494.
21. Mokhtary M, Rastegar Niaki M (2013) Polyvinylpolypyrrolidone-supported boron trifluoride (PVPP-BF₃); highly efficient catalyst for oxidation of aldehydes to carboxylic acids and esters by H₂O₂. *Iran J Chem Chem Eng* 32(1): 43-48.
22. Mokhtary M, Lakouraj M M, Rastegar Niaki M (2012) Polyvinylpolypyrrolidone-supported boron trifluoride (PVPP-BF₃); highly efficient catalyst for chemoselective oxygenation of sulfides to sulfones by H₂O₂. *Phosphorus Sulfur Silicon Relat Elem* 187(3): 321-326.
23. Pourali A R, Ghayeni S, Afghahi F (2013) Efficient and regioselective ring-opening of epoxides with alcohols and sodium azide by using catalytic amounts of GaCl₃/polyvinylpyrrolidone. *Bull Korean Chem Soc* 34(6): 1741-1744.
24. Nabatchi Ahmadi V, Mokhtary M (2018) Polyvinylpolypyrrolidone supported antimony(III) chloride (PVPP-SbCl₃): an efficient catalyst for the synthesis of chromenylphenylpropanones. *Iran Catal* 8(1): 53-58.
25. Ghorbani-Choghamarani A, Goudarziafshar H, Zamani P (2011) Polyvinylpoly pyrrolidonium tribromide as new and metal-free catalyst for the formylation and trimethylsilylation of hydroxyl group. *Chin Chem Lett* 22(10): 1207-1210.
26. Khaksar S, Tajbakhsh M, Gholami M (2014) Polyvinylpolypyrrolidone-supported triflic acid (PVPP-OTf) as a new, efficient, and recyclable heterogeneous catalyst for the synthesis of bis-indolyl methane derivatives. *C R Chim* 17(1): 30-34.
27. Khaksar S, Tajbakhsh M, Gholami M, Rostamnezhad F (2014) A highly efficient procedure for the synthesis of quinoxaline derivatives using polyvinylpolypyrrolidone supported triflic acid catalyst (PVPP-OTf). *Chin Chem Lett* 25(9): 1287-1290.
28. Sheik Mansoor S, Ghashang M (2015) Synthesis of a novel series of 7-hydroxy-10-aryl-10H-indeno[1,2-b]chromen-11-ones, indeno[1,2-b]naphtho[1,2-e]pyran-12(13H)-one, and indeno [1,2-b]naphtho [3,2-e]pyran-5,11,13-trione catalyzed by reusable polyvinylpolypyrrolidone-supported triflic acid. *Res Chem Intermed* 41: 9085-9100.
29. Nikoorazm M, Ghorbani-Choghamarani A, Goudarziafshar H (2011) Supported nitric acid on polyvinylpolypyrrolidone as an efficient and mild nitrating agent for the selective preparation of nitrophenols. *Sci Iran* 18: 1353-1355.
30. Ghorbani-Choghamarani A, Chenani Z, Mallakpour S (2009) Supported Nitric Acid on Silica Gel and Polyvinyl Pyrrolidone (PVP) as an Efficient Oxidizing Agent for the Oxidation of Urzoles and Bis-urzoles. *Synth Commun* 39: 4264-4270.
31. Zolfigol M A, Amani K, Ghorbani-Choghamarani A, Hajjami M, Ayazi-Nasrabadi R, et al. (2008) Chemo and homoselective catalytic oxidation of sulfides to sulfoxides with supported nitric acid on silica gel and poly vinyl pyrrolidone (PVP) catalyzed by KBr and/or NaBr. *Catal Commun* 9(8): 1739-1744.
32. Shirini F, Najafi Moghadam P, Moayedi S, Seddighi M (2014) Introduction of O-sulfonated poly(4-vinylpyrrolidonium) chloride as a polymeric and reusable catalyst for the synthesis of xanthenes derivatives. *RSC Adv* 4: 38581-38588.
33. Yosefzadeh M, Mokhtary M (2016) Polyvinylpolypyrrolidone supported chlorosulfonic acid: An efficient catalyst for the one-pot synthesis of hexahydroquinolines. *Iran J Catal* 6(2): 153-159.
34. Goudarzvand Chegini M, Mokhtary M (2017) Polyvinylpolypyrrolidone-supported chlorosulfonic acid: an efficient catalyst for one-pot synthesis of dihydropyrimidinones and octahydroquinazolin-2,5-diones. *Polycycl Aromat Comp* 37(1): 63-72.
35. Abedini M, Shirini F, Mousapour M (2016) Poly(vinylpyrrolidonium) perchlorate as a new and efficient catalyst for the promotion of the synthesis of polyhydroquinoline derivatives via Hantzsch condensation. *Res Chem Intermed* 42(3): 2303-2315.
36. Shirini F, Goli-Jolodar O, Akbari M, Seddighi M (2016) Preparation, characterization, and use of poly(vinylpyrrolidonium) hydrogen phosphate ([PVP-H]H₂PO₄) as a new heterogeneous catalyst for efficient synthesis of 2-amino-tetrahydro-4H-pyran. *Res Chem Intermed* 42: 4733-4749.
37. Safaei M A, Safaei M, Shekouhy M (2015) Sulfuric acid-modified poly(vinylpyrrolidone) ((PVP-SO₃H)HSO₄): a new highly efficient, bio-degradable and reusable polymeric catalyst for the synthesis of acridinedione derivatives. *RSC Adv* 5: 6797-6806.
38. Li Y, Boone E, El-Sayed M A (2002) Size effects of PVP-Pd nanoparticles on the catalytic Suzuki reactions in aqueous solution. *Langmuir* 18: 4921-4925.
39. Tamami B, Allahyari H, Ghasemi S, Farjadian F (2011) Palladium nanoparticles supported on poly(N-vinylpyrrolidone)-grafted silica as new recyclable catalyst for Heck cross-coupling reactions. *J Organometallic Chem* 696(2): 594-599.



This work is licensed under Creative Commons Attribution 4.0 License

**Your next submission with Juniper Publishers
will reach you the below assets**

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission

<https://juniperpublishers.com/online-submission.php>