



Proceeding

Volume3 Suppl 1 – March 2017
DOI:10.19080/AJPN.2017.03.555677

Acad J Ped Neonatol

Copyright © All rights are reserved by Mohamed A Talat

The Role of Osteopontin in the Pathogenesis and Complication of Type 1 Diabetes Mellitus in Children and Adolescents

Mohamed A Talat*

Department of Pediatrics, Zagazig University, Egypt

Submission: March 05, 2017; **Published:** March 28, 2017

***Corresponding author:** Mohamed A Talat, Department of Pediatrics, Zagazig University, Egypt

Back ground

Type 1 diabetes mellitus (T1DM) is the one of the most common autoimmune disorders in childhood. Osteopontin has a role in the development and progression of several autoimmune diseases and promotes adipose tissue inflammation, dysfunction and insulin resistance.

Objective

To investigate serum osteopontin in pediatric patients with T1DM and to ascertain if it predicts diabetes complications.

Methods

This was a case-control study at Endocrinology unit, Children's Hospital, Zagazig University, Egypt, from October 2014 to December 2015. Sixty patients with T1DM and 60 healthy subjects were enrolled. All the children had detailed medical history, clinical examination, ophthalmoscopy; laboratory estimation for fasting blood glucose, HbA1c, urine albumin/creatinine ratio and serum osteopontin.

Results

Patients with T1DM had significantly higher serum osteopontin levels compared with controls (mean±S.D: 13.7±3.4

µg/L vs 8.9±2.9 µg/L, P<0.001), Serum osteopontin concentrations were higher in patients with micro albuminuria than patients with normal albumin excretion rate and control group. Those who had retinal disease had higher osteopontin concentrations than those without (16.8±2 vs. 12.4±3mg/L; P=0.005). Serum osteopontin levels in diabetic patients correlated with higher systolic and diastolic blood pressure, body mass index, lower HDL, retinopathy and micro albuminuria. No correlation was found between osteopontin levels and HbA1c, insulin dose, co-medications, and diabetes duration in T1DM patients. The association between high osteopontin levels and T1DM was independent from all confounders.

Conclusion

Increased osteopontin levels are associated with T1DM in pediatric patients. Osteopontin may have a role in the prediction of microvascular diabetes complications.



This work is licensed under Creative Commons Attribution 4.0 License
DOI:10.19080/AJPN.2017.03.555677

Your next submission with Juniper Publishers will reach you the below assets

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission

<https://juniperpublishers.com/online-submission.php>