



Case Report

Volume 13 Issue 1 - October 2023
DOI: 10.19080/AJPN.2023.13.555908

Acad J Ped Neonatol

Copyright © All rights are reserved by Aniruddha Basak

A Case of Omphalocele Associated with Teratoma in a New Born Baby: A Very Rare Case Report



Aniruddha Basak*, Sathi Dhar and Niladri Shekar Dey

Assistant Professor, Department of General Surgery, Tripura medical college and DR. BRAM teaching hospital, India

Submission: August 31, 2023; **Published:** October 12, 2023

***Corresponding author:** Aniruddha Basak, Assistant Professor, Department of General Surgery, Tripura medical college and DR. BRAM teaching hospital, India

Abstract

An omphalocele is a congenital midline abdominal wall defect at the level of the umbilical cord insertion into the fetal abdomen that results in herniation of intra-abdominal contents into a sac comprising of an outer layer of amnion and an inner layer of peritoneum. The most common eviscerated contents are liver and small intestine though virtually any organ can be seen within the omphalocele sac including colon, pancreas, spleen, omentum, stomach, heart or genitalia. We report an unusual case of omphalocele associated with mature teratoma in Tripura medical college and DR. BRAM teaching hospital. Only five cases of teratoma arising in a case of omphalocele are reported in the literature reviewed.

Keywords: Omphalocele; Teratoma; Children

Introduction

Omphalocele is a congenital malformation of the anterior abdominal wall, which has a central localization and is characterized by the presence of a hernial sac with contents inside. Most often, the contents of the hernial sac are the liver, and loops of the small or large intestine. The incidence of omphalocele varies from 1 in 4,000 to 1 in 10,000 [1-3]. In comparison with other malformations of the anterior abdominal wall, omphalocele is often combined with other congenital malformations of the cardiovascular, respiratory or gastrointestinal tracts. Approximately 40% of patients with omphalocele have an abnormal karyotype [4]. Teratoma is a germ cell tumor, which is most often localized in the sacrococcygeal region (89%). Teratomas can be cystic, cystic-solid, or mixed. They can also be found in other areas. Congenital teratomas are most common benign formations, although in recent years there have been reports of early malignancy of the formation. Surgical treatment for small and medium-sized omphalocele without viscerio-abdominal imbalance consists of a one-stage reconstruction of the anterior abdominal wall. With a large defect and an insufficient volume of the abdominal cavity, treatment consists in a two-stage immersion of the organs as the volume of the abdominal cavity increases. We describe a clinical

case of successful surgical treatment of umbilical cord teratoma in a case of omphalocele major containing liver, small and large intestine as content in Tripura medical college and DR. BRAM teaching hospital.

A Case Report

A full-term neonate of uneventful antenatal history presented with two large swellings in umbilical region. There was no other complaint. On physical examination the neonate was pink, active and weighed 2.5 kg. Systemic examination was unremarkable. On local examination, there was a large swelling measuring 8 × 7 cm covered partly by skin and amnion both. A clinical diagnosis of exomphalos major was made. Another swelling measuring 5 cm x 4 cm covered fully by skin with tuft of hair in it. The surface of the swelling was irregular, and the consistency is variegated. A clinical diagnosis of umbilical cord teratoma was made. Apart from routine investigations, X-ray chest and echocardiography were done all of which were within normal limits. Surgical treatment was performed 15 hours after birth following stabilization of the new born's condition and correction of water electrolyte balance and impaired coagulation. The tumour was surgically removed without disturbing the omphalocele major component. The tumor-

like formation was on the external surface of the omphalocele, 5 cm × 4 cm in size with irregular margins along with tuft of hairs and no invasion into adjacent tissues. Inside, a structure of bony-cartilaginous consistency was palpated. Since there was insufficient volume of the abdominal cavity, one-stage plastic

surgery of the anterior abdominal wall was not performed. The postoperative period was uneventful. On the third post-operative day there were no signs of inflammation and dehiscence of the surgical wound and it eventually healed by primary intention. The child was discharged on the 11th day after surgery (Figure 1-5).



Figure 1: pre-operative picture showing a case of omphalocele major with umbilical teratoma.



Figure 2: post operative picture showing wide local excision of teratoma with treatment of omphalocele major with 2% mercurochrome solution.

Discussion

The medical literature describes rare cases of congenital umbilical cord teratomas [5]. In our clinical case, the child had a combination of a mature teratoma with a congenital malformation

of the anterior abdominal wall- omphalocele. We share our experience of successful surgical treatment of a newborn with this extremely rare combination. One-third of newborns with omphalocele detected in the antenatal period have related developmental abnormalities [1]. The frequency of malformation

ranges from 1:5,000-1:10,000 live-born children [6]. Congenital teratomas of the umbilical cord are extremely rare and only 15 cases have been reported in the literature. Five cases included a combination of umbilical cord teratoma and omphalocele [7-9], with two early abortions due to chromosomal abnormalities. Histologically, teratomas are often constituted of embryonic cells of all three germ layers. In this case, omphalocele and teratoma are two different pathologies in the same baby at the same time. Since there was significant viscero-abdominal disproportion, surgical treatment was performed to deal with Umbilical Teratoma by removal of the teratoma with simultaneous regular dressing of the omphalocele with 2% mercurochrome solution. Congenital

omphalocele needs to be differentiated from gastroschisis, which is also a congenital malformation of the anterior abdominal wall. Unlike omphalocele, the defect in gastroschisis is most often located to the right of the normally formed umbilical cord, and the hernial sac is absent in most cases. Despite the pronounced changes in the internal organs in gastroschisis and the presence of viscer abdominal imbalance, the prognosis following treatment is most often favourable. In conclusion, combination of the omphalocele and the teratoma that we have described is extremely rare and difficult to diagnose antenatally. This malformation is correctable and does not constitute an indication for the termination of pregnancy.




Figure 3: HPE showing hair follicle and neural tissue.




Figure 4: HPE showing bone and marrow.




Figure 5: HPE showing GI mucosa.

References

1. Shawkat E, Gillham J, Craigie RJ (2012) Rare combination of exomphalos with umbilical cord teratoma. *Ultrasound Obstet Gynecol* 40(4): 481.
2. Hamilton CA, Harris JE, Ellison MC, et al. (2011) Cystic teratoma. In: *Medscape Reference WebMD, LLC Drugs, Diseases & Procedures* p. 1-8.
3. Harish K, Ashok AC, Alva NK (2003) Low grade fibromyxoid sarcoma of the falciform ligament: A case report. *BMC Surg* 3: 7.
4. Kreczy A, Alge A, Menardi G, Gassner I, Gschwendtner A, et al. (1994) Teratoma of the umbilical cord Case report with review of the literature. *Arch Pathol Lab Med* 118(9): 934-937.
5. Hargitai B, Csabai L, Ban Z, Hetenyi I, Szucs I, et al. (2005) Rare case of exomphalos complicated with umbilical cord teratoma in a fetus with trisomy. *Fetal Diagn Ther* 20(6): 528-533.

6. Gross RE, Clatworthy HW, Meeker IA (1951) Sacrococcygeal teratoma in infants and children: report of 40 cases. Surg Gynecol Obstet 92(3): 341-354.
7. Sant PK, Sanjay K, Manish C, Gajender S (2012) Teratoma arising in omphalocele sac. case report of a rare entity. Ann Diagn Paediatr Pathol 16: 493-497.
8. Available online at: Teratoma_arising_in_omphalocele_sac_Case_report_of_a_rare_entity.
9. https://www.researchgate.net/publication/233961016_Teratoma_arising_in_omphalocele_sac_Case_report_of_a_rare_entity



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/AJPN.2023.13.555906](https://doi.org/10.19080/AJPN.2023.13.555906)

**Your next submission with Juniper Publishers
will reach you the below assets**

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission

<https://juniperpublishers.com/online-submission.php>