



Mini Review

Volume 9 Issue 2 – March 2018  
DOI: 10.19080/ARGH.2018.09.555756

Adv Res Gastroentero Hepatol

Copyright © All rights are reserved by Jayesh V Trivedi

# Coarctation of Aorta



Jayesh V Trivedi\*

Gujarat Adani Institute of Medical Sciences (GAIMS), G K General Hospital, India

Submission: January 12, 2018; Published: March 22, 2018

\*Corresponding author: Jayesh V Trivedi, Gujarat Adani Institute of Medical Sciences (GAIMS), G K General Hospital, India,  
Email: drjvtrivedi@rediffmail.com

## Mini Review

Coarctation of aorta occurs in 5 to 8 % of patients with congenital heart diseases patient .It is one of the cause of secondary Hypertension in young patients. It may be isolated or may be accompanied by congenital cardiac diseases like Ventricular septal disease or Bicuspid Aortic valve. Sometimes it is associated with Left ventricular heart failure. It may be associated with Turner's syndrome.Coarctation of aorta is defined as constricted aortic segment that comprises localised medial thickening with some infolding of medial and superimposed neonatal tissues. The Coarctation may be discrete or long segment of aorta may be involved.

- a) Preductal Coarctation
- b) Ductal Coarctation
- c) Postductal Coarctation

The classical coarctation of aorta is in thoracic aorta after the origin of left subclavian artery. Rarely it may be seen in lower thoracic aorta or abdominal aorta. Usually all coarctations are

juxtaductal. Post stenotic dilatation of aorta is commonly seen and it occurs due to jet effect of the blood flow distal to the occlusion. As it is congenital disorder other associated disorders like Patent Ductus Arteriosus, Transposition of great vessels, Tricuspid atresia and Hypoplastic left ventricle syndrome is to be ruled out [1-5].

## Signs and Symptoms

Common in boys than girls. Girls with Turner syndrome are likely to have coarctation. Usually children are asymptomatic but may present as Left ventricular failure. Intermittent claudication occur at a later age. Hypotension or absent pulsation in lowerlimb and hypertension in upperlimb is common. Infective endocarditis can take place. Mitral prolapse is associated with this. Differential cyanosis is also one clinical presentation. Usually children are asymptomatic and in adults at about 40 years this is diagnosed. However some patients do get chest pain, cold feet, leg cramps with exercise, poor growth in lower half of body, epistaxis [6].

## Diagnosis




Figure 1: MRI picture of one of our patient.

Pulse in the lower limb will be weaker than upper limb and radio femoral delay can be felt. On auscultation murmur will be audible on chest or back. Cardiac catheterisation and aortography helps in localising the extent of coarctation but they are invasive tests [7] (Figure 1).

Post stenotic dilatation will have inverted 3 appearance on chest x-ray. With prominent lower intercostal grooves on ribs due to dilated intercostal arteries. Echocardiography is helpful in children. Magnetic resonance angiography is the most accurate test to diagnose Coarctation of aorta. Contrast MRI and Phase Contrast MRI are special investigation to diagnose the severity of the disease.

Rarely it is being diagnosed at the time of pregnancy in female while treating secondary pregnancy associated hypertension. Circle of willis abnormalities in the brain is also associated with Coarctation of Aorta. This results in to stroke in adults. This may be the cause of habitual abortion in females [8].

### Conclusion

Fainting and chest pain in paediatric practice is being ignored but in such case suspicious of coarctation should be done. One

should always form a habit of palpating pulse in all four limbs. Secondary hypertension should be investigated. Whenever congenital heart disease is diagnosed associated coarctation of aorta should be excluded. Patients having coarctation of aorta should receive proper antibiotics for the prevention of infective endocarditis.

Early diagnosis will help in proper development of child and helps in preventing the complications.

### References

1. Coarctation of the aorta - Mayo Clinic
2. [www.mayoclinic.org/diseases-conditions/coarctation-of-the-aorta/.../con-20031772](http://www.mayoclinic.org/diseases-conditions/coarctation-of-the-aorta/.../con-20031772)
3. Coarctation of the Aorta (CoA) - American Heart Association
4. [www.heart.org/.../Coarctation-of-the-Aorta-CoA\\_UCM\\_307022\\_Article.jsp](http://www.heart.org/.../Coarctation-of-the-Aorta-CoA_UCM_307022_Article.jsp)
5. Coarctation of the aorta: MedlinePlus Medical Encyclopedia
6. <https://medlineplus.gov> Medical Encyclopedia
7. Co-arctation of the Aorta | Congenital Heart Defects UK - [chd-uk.co.uk](http://chd-uk.co.uk)
8. [www.chd-uk.co.uk/types-of-chd-and-operations/co-arctation-of-the-aorta/](http://www.chd-uk.co.uk/types-of-chd-and-operations/co-arctation-of-the-aorta/)



This work is licensed under Creative Commons Attribution 4.0 License  
DOI: [10.19080/ARGH.2018.09.555756](https://doi.org/10.19080/ARGH.2018.09.555756)

### Your next submission with JuniperPublishers will reach you the below assets

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats  
( Pdf, E-pub, Full Text, audio)
- Unceasing customer service

Track the below URL for one-step submission  
<https://juniperpublishers.com/online-submission.php>