



Research Article

Volume 14 Issue 5 - March 2018
DOI: 10.19080/ARTOAJ.2018.14.555936

Agri Res & Tech: Open Access J

Copyright © All rights are reserved by Nichanun Kernasa

Efficiency of Entomopathogenic Fungi to Sugarcane White Leafhopper, *Matsumuratettix hiroglyphicus* (Matsumura) (Hemiptera: Cicadellidae)



Nichanun Kernasa^{1,2,3*}, Parichat Jamrutsri², Siriya Kumpiro^{1,4}, Roongnapa Korpaditskul³ and Sophon Uraichuen^{1,2}

¹Department of Entomology, Kasetsart University, Thailand

²National Biological Control Research Center, Kasetsart University, Thailand

³Center of Excellence for Sugarcane, Kasetsart University, Thailand

⁴Department of Entomology and Plant Pathology, Chiang Mai University, Thailand

Submission: March 02, 2018; **Published:** March 28, 2018

*Corresponding author: Nichanun Kernasa, Department of Entomology, Faculty of Agriculture Kamphaengsaen, Kasetsart University, Kamphaeng Saen campus, Nakhon Pathom, 73140, Thailand, Email: agropk@ku.ac.th

Abstract

Sugarcane White Leafhopper, *Matsumuratettix hiroglyphicus* (Matsumura) (Hemiptera: Cicadellidae) is an economic important insect pest of sugarcane in Thailand. The three entomopathogenic fungi, *Metarhizium anisopliae* isolated from the sugarcane longhorn beetle, *Beauveria brassiana* isolated from the brown plant hopper and *Purpureocillium lilacium* (*Paecilomyces lilacinus*) isolated from *M. hiroglyphicus* were tested as biological control agents of *M. hiroglyphicus*. The results indicated that *M. anisopliae* isolated from *D. buqueti* showed the best pathogenicity to *M. hiroglyphicus* adults with $LC_{50} 7.39 \times 10^6$

Introduction

Sugarcane White Leafhopper, *Matsumuratettix hiroglyphicus* (Matsumura) (Hemiptera: Cicadellidae) is an economic important insect pest of sugarcane in Thailand. It represents as the reservoir of phytoplasma that cause sugarcane white leaf disease [1]. In recently, sugarcane white leaf disease spread out from the Northeastern region to the lower north region and the central region of Thailand. These causes sharply decrease in sugarcane yields. So that, entomopathogenic fungi become the discriminatory technique to control *M. hiroglyphicus*. This paper aims to evaluate the efficiency of three species of entomopathogenic fungi to *M. hiroglyphicus* adults in the laboratory.

Materials and Methods

Insect culture

The adults of *M. hiroglyphicus* were collect lively by setting the light trap in sugarcane field that having the white leaf disease at Bueng Samakkhi district, Kamphaeng Phet province. Then, transferred them to the National Biological Control Research Center, Central Regional Center, Kasetsart University, Kamphaeng Saen campus, Nakhon Pathom' s laboratory for rearing. They were reared on the one month sugarcane that placed in the

rounded plastic cages until we prompted to do the experiment and bioassay.

Study on pathogenicity of three species of entomopathogenic fungi to *M. hiroglyphicus* adults

The experiment consisted of three species of entomopathogenic fungi that were *Metarhizium anisopliae* isolated from the sugarcane longhorn beetle, *Beauveria brassiana* isolated from the brown plant hopper and *Purpureocillium lilacium* (*Paecilomyces lilacinus*) isolated from *M. hiroglyphicus* compared with control, distill water mixed with 0.05% Triton X 100. Each treatment consisted of five replications that were five adults per one Petri-dish. The trial done by dropping 1 μ l of 10^8 conidia/ml of each fungus on each adult. They were placed in 25 ± 2 °C and $70 \pm 2\%$ RH. The data were checked for ten days, the adults were checked that the spores of each fungus grow cover their bodies by the necked eyes. The data were collected and calculated by statistic tool.

Bioassay of *M. anisopliae* to *M. hiroglyphicus* adults

Metarhizium anisopliae was the better candidate of entomopathogenic fungus for controlling *M. hiroglyphicus* adults. We used five concentrations of *M. anisopliae*; 10^5 , 10^6 ; 10^7 ; 10^8 and

10⁹ conidia/ml compared with distill water mixed with 0.05% Triton X 100. Each treatment consisted of five replications that were five adults per one Petri-dish. The trial done by dropping 1µl of each concentration of each fungus on individual adult. They were placed in 25±2 °C and 70±2% RH. The data were checked for ten days, the adults were checked that the spores of each fungus grow cover their bodies by the necked eyes. The data were collected and calculated LC₅₀ by probit analysis.

Results and Discussion

Pathogenicity of three species of entomopathogenic fungi to *M. hiroglyphicus* adults

Table 1: Percent mortality (%) and LT₅₀ (days) of *M. hiroglyphicus* adults caused by *M. anisopliae*, *B. brassiana* and *P. lilacinum*.

Treatment	Percent Mortality (%)	LT50 (days)
<i>Beauveria bassiana</i>	76.00b	8.5
<i>Metarhizium anisopliae</i>	100.00a	3.71
<i>P. lilacinum</i>	68.00b	9.24
Control	0.00c	-

Percent mortality within a column followed by the same letter are not significantly different based on DMRT (P ≤ 0.05).




Figure 1: Percent mortality (%) of *M. hiroglyphicus* adults caused by *M. anisopliae*, *B. brassiana*, *P. lilacinum* within ten days.

The results of pathogenicity of the three fungi revealed that they can infect *M. hiroglyphicus* adults. Percent mortalities of *M. hiroglyphicus* adults were significantly different between the three fungi there were 100, 76, 68 and 0 percent by *M. anisopliae*, *B. brassiana*, *P. lilacinum* and control, respectively. The LT₅₀ indicated that *M. anisopliae* showed the rapid mortality was 3.71 days followed by *B. brassiana*, *P. lilacinum* that were 8.50 and 9.24 days, respectively (Table 1) and (Figure 1). Vestergaard et al. [2] treated *M. anisopliae* to adult *Frankliniella occidentalis* with resulted in at least 94% mortality at 7 days post-inoculation. Annamalai et al. (2016) reported that *B. brassiana* showed percent mortality of 78.48% for the concentrations of 1.23×10⁸ spores/mL to *Thrips tabaci*. Jone et al. [3] reported that

M. anisopliae strains were more virulent, with lower LT₅₀ values, than were the *B. brassiana* strains [4].

Bioassay of *M. anisopliae* to *M. hiroglyphicus* adults

The result from pathogenicity indicated that *M. anisopliae* showed the highest percent mortality and lowest LT₅₀ so that, we chosen *M. anisopliae* to do the bioassay to *M. hiroglyphicus* adults. The bioassay consisted of five treatments compared with control. The treatments were conidial suspensions 1×10⁵, 1×10⁶, 1×10⁷, 1×10⁸, 1×10⁹ conidia/ml and control. The results revealed that *M. anisopliae* showed 7.39×10⁶ of the LT₅₀ (Table 2). The probit analysis showed that R² was 0.9638 (Figure 2).




Figure 2: Probit analysis of five conidial suspensions of *M. anisopliae* to *M. hiroglyphicus* adults.

Table 2: LT₅₀ of *M. anisopliae* to *M. hiroglyphicus* adults.

Conidia Suspensions (Conidia/ml)	LC ₅₀ (Conidia/ml)	95% Fiducial CI	
		Lower	Upper
1x10 ⁵	7.39x10 ⁶	2.29x10 ⁶	2.38x10 ⁷
1x10 ⁶			
1x10 ⁷			
1x10 ⁸			
1x10 ⁹			
Control			

According to the results revealed that *M. anisopliae* isolated from *D. buqueti* showed the best pathocinity to *M. hiroglyphicus* adults with LT₅₀ 7.39×10⁶. This fungus will be a promising biological control agent to control *M. hiroglyphicus* in sugarcane plantations.

References

- Hanboonsong Y, Choosai C, Panyim S, Damak S (2002) Transovarial transmission of sugarcane white leaf phytoplasma in the insect vector *Matsumuratettix hiroglyphicus* (Matsumura). *Insect Mol Biol* 11(1): 97-103.
- Vestergaard S, Gillespie AT, Butt TM, Schreiter G, Eilenberg J (1995) Pathogenicity of the Hyphomycete Fungi *Verticillium lecanii* and *Metarhizium anisopliae* to the Western Flower Thrips, *Frankliniella occidentalis*. *Biocontrol Science and Technology* 5(2): 185-192.

3. Jones WE, Grace JK, Tamashiro M (1996) Virulence of Seven Isolates of *Beauveria bassiana* and *Metarhizium anisopliae* to *Coptotermes formosanus* (Isoptera: Rhinotermitidae). *Environmental Entomology* 25(2): 481-487.
4. Annamalai M, Kaushik HD, Selvaraj K (2016) Bioefficacy of *Beauveria bassiana* (Balsamo) Vuillemin and *Lecanicillium lecanii* Zimmerman against *Thrips tabaci* Lindeman. *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences* 86(2): 505-511.



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/ARTOAJ.2018.14.555936](https://doi.org/10.19080/ARTOAJ.2018.14.555936)

Your next submission with Juniper Publishers will reach you the below assets

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission

<https://juniperpublishers.com/online-submission.php>