

Bacterial Survey of Diabetic Ulcers in Diabetic Patients in Hospitals at Aba, Nigeria



Akunneh-Wariso C¹ and Aduema W^{2*}

¹Department of Human Physiology, Abia State University, Nigeria

²Department of Human Physiology, PAMO, University of Medical Sciences, Nigeria

Submission: April 27, 2019; **Published:** July 18, 2019

***Corresponding author:** Aduema Wadioni, Department of Human Physiology, PAMO University of Medical Sciences, Port Harcourt, Rivers State, Nigeria

Abstract

Diabetic ulcers are usually seen as infected ulcers and run a chronic course. Selected documented records of diabetic patients with diabetic ulcers were taken from the laboratory sections of different hospitals and clinics in Aba. From these records, a total of 110 diabetic ulcer cases were used in this study irrespective of their sexes and ages. Trauma related ulcers were ruled out. Microscopy and culture of specimen or wound swabs from these ulcers after 48 hours at 36 °C incubation showed that the most common bacteria in these ulcers is *Escherichia Coli* (*E. coli*) followed by *staphylococcus* species which is 22%. These bacteria species were seen in most of these ulcers. This study showed that most diabetic ulcers are mostly or commonly infected by *Escherichia species* and *staphylococcus species*.

Keywords: Diabetic ulcer; Patient; E. Coli; Bacteria; Diabetes mellitus; Carbohydrate; Fat; Protein metabolism; Diabetic nephropathy; Diabetic neuropathy; Diabetic retinopathy; Non-enzymatic glycosylation; Antioxidant

Introduction

Diabetes mellitus is one of the commonest diseases in tropical areas. Diabetes mellitus may be described as a syndrome characterized by hyperglycemia and disturbance of carbohydrate, fat and protein metabolism resulting from defects in insulin secretion, insulin action or both [1]. Aside complications like diabetic nephropathy, neuropathy and retinopathy, diabetic ulcers are seen as the most troubling problem in the hospitals and clinics [2]. Most times, it involves the attention of both surgeon and internal physicians. It has been reported that diabetic complications are associated with overproduction of Reactive Oxygen Species (ROS) [3]. This complication is considered the leading cause of death among these patients. Oxidation plays a major role in diabetes [4]. The increase in free radical release accompanied by decrease in antioxidants is a major cause of diabetes [5]. In diabetes mellitus, there are usually alterations in the endogenous free radical scavenging defenses which leads to ineffective scavenging of reactive oxygen species resulting to oxidative damage [6]. Experimental diabetes induced by streptozotocin, selectively destroys the β -cells of pancreas by generating excess reactive oxygen species and produces kidney lesions that are similar to human diabetic nephropathy [7].

Studies have shown that individuals with diabetes have a higher incidence of kidney function abnormalities, as well as for matation of free radical due to glucose oxidation, non-enzymatic

glycosylation of proteins and subsequent oxidative degradation of glycated proteins, leading to a decline in antioxidant defense mechanism and damage of cellular organelles and enzymes and development of insulin resistance which ultimately, culminate in renal complications [8,9]. Thus, an early control of DM is recommended as one of main strategy to prevent these complications and increase the life span of these patients. Knowledge of the common pathogens associated with diabetic ulcers may hasten quick recovery of the patients as definitive therapy will be applied after confirmatory results of wound swab culture.

Materials and Methods

A retrospective study of confirmed 110 diabetic patients with diabetic ulcers managed in different hospitals and clinics at Aba over 9 months were taken from the laboratory sections of the hospitals and clinics irrespective of their sexes and ages. Trauma related ulcers were ruled out. Specimens or wound swabs from different ulcers were taken before commencement of treatment under sterile conditions for microscopy, culture and sensitivity (MSC) at 30°C incubation for 48 hours. The results of these ulcer and wound swabs were shown below.

It should be noted that some of these patients had been on treatment either from patent medicine dealers or other clinics without records.

Ethical Consideration

Ethical approval was obtained from the State ministry of Health, Aba State; Nigeria. Consent from patients from whom data were obtained”.

Results: Table 1 & Figure 1

Table 1: Bacterial Survey of Diabetic Ulcers in Diabetic Patients.

Bacterial spp.	Frequency	Percentage (%)
Escherichia Coli (E coli)	40	36
Staphylococcus spp.	24	22
Pseudomonas spp.	20	18
Proteus spp.	10	9
No significant bacterial growth	16	15
Total	110	100




Figure 1: Bacterial Survey of Diabetic Ulcers in Diabetic Patients.

Discussion

Aside complications like diabetic nephropathy, diabetic ulcers are seen as the most troubling problem in the hospital and clinics [2]. Thus, early control is recommended as one of the main strategies to prevent these complications. Diabetic ulcer if not checked leads to the destruction of body tissues and impair patient’s quality of life. Most diabetic ulcers can be prevented

with good care and screening of people for risk factors [10]. It has also been implicated as one of the causes of Charot arthropathy [11,12], which involves progressive damage to the bones, joints and tissues of the body [12,13].

Therefore, the aim of this study was to carry out a bacterial survey of diabetic ulcers among diabetic patient in Aba, hospital in Abia State, Nigeria. Documented medical records of 110 confirmed diabetic patients admitted and managed in different hospitals and clinics at Aba were taken. Specimens or wound swabs were taken before commencement of treatment, though some patients would have being in several antibiotics without records. Our findings showed that Out of 110 diabetic patients with diabetic ulcers, specimens or wound swabs from these

patients showed that 36.0% of cases were infected with *Escherichia coli*, 22.0% cases with *Staphylococcus* spp. This result also showed that 18.0% and 9.0% of cases were infected with *pseudomonas* spp and *proteus. spp* respectively. Cultures of specimens or wound swabs of 15.0% of cases showed no significant growth. This indicated that these patients had being on several antibiotics in other places with no documentation or records.

Conclusion

In conclusion, our bacterial survey of diabetic ulcers in diabetic patients revealed that the most predominant bacterial in diabetic ulcers in our center is *Escherichia coli (E. coli)* followed by *Staphylococcus* spp. Therefore, the knowledge of this bacterial survey will be useful in management of diabetic ulcers.

References

1. Falase AO, Akinkugbe OO (2006) A Compendium of Clinical Medicine.
2. Giorgino F, Lavida L, Cavallo PP, Solnica B, Fuller J, et al. (2004) Factors associated with progression to macroalbuminuria in microalbuminuric type 1 diabetic patient: The EURODIAB prospective complications study. *Diabetologia* 47(6): 1020-1028.
3. Palanduz S, Ademoglu E, Gokkusu C (2001) Plasma antioxidants and type 2 diabetes mellitus. *Res Commun Mol Pathol Pharmacol* 109(5-6): 309-318.
4. Lieber CS (1997) Ethanol metabolism, cirrhosis and alcoholism. *Clin Chim Acta* 258(1): 59-84.
5. Mohamed AK, Bierhaus A, Schiekofers S, Tritschler H, Ziegler R, et al. (1999) The role of oxidative stress and NF (B) activation in late diabetic complications. *Biofactors* 10(2-3): 175-179.
6. Oberley LW (1988) Free radicals and diabetes. *Free Radic Biol Med* 5(2): 113-124.
7. Boukhris MM, Bouaziz I, Feki H, Jemai A, Feki EL, et al. (2012) Hypoglycemic and antioxidant effects of leaf essential oil of *Pelargonium graveolens* L'Hér. in alloxan induced diabetic rats. *Lipids Health Dis* 11: 81.
8. Kangralkar VA, Shivraj DP, Bandivadekar RM (2010) Oxidative stress and diabetes: a review. *International Journal of Pharmaceutics* 1(1): 38-45.
9. Arora S (2010) Renal function in diabetic nephropathy. *World J Diabetes* 1(2): 48-56.
10. Zhang P, Lu J, Jing Y, Tang S, Zhu D, et al. (2017) Global epidemiology of diabetic foot ulceration: a systemic review and meta-analysis (f). *Ann Med* 49(2): 106-116.
11. (2017) American Diabetes Association. Standard of Medical Care in diabetes-2017. Abridged for primary care providers. *Clinical Diabetes* 35: 5-26.
12. Rogers LC, Frykberg RG, Sanders IJ (2016) The diabetic Charcot foot: recognition, evaluation and management. In: Armstrong DG, Lavery LA (Eds.), *Clinical care of the diabetic foot*, (3rd edn.), p. 99.
13. Pop-Busiu, R, Boulton AJM, Feldman EL, Bril V, Freeman R (2017) Diabetic neuropathy: A position statement by the American Diabetic Association. *Diabetes care* 40(1): 136-154.



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/CRDOJ.2019.11.555814](https://doi.org/10.19080/CRDOJ.2019.11.555814)

**Your next submission with Juniper Publishers
will reach you the below assets**

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission

<https://juniperpublishers.com/online-submission.php>