

Aneurysmal Subarachnoid Hemorrhage



José Manuel Ortega Zufiría*, Pilar Jerez Fernández and Jorge Zamorano Fernández

Neurosurgery Department, Hospital Universitario de Getafe, Spain

Submission: January 21, 2018; Published: March 26, 2018

*Corresponding author: José Manuel Ortega Zufiría, Neurosurgery Department, Hospital Universitario de Getafe, Ctra. de Toledo km. 12,5, 28905, Getafe, Madrid, Spain, Email: fuencarral108@hotmail.com

Abstract

Introduction: Aneurysmal subarachnoid hemorrhage (ASH) is a very serious phenomenon associated with high rates of morbidity and mortality. After the initial hemorrhage, 50% of patients die, and 30-40% of patients re-blood during the first month. The percentage of deaths due to re-bleeding is between 60% and 75%. Re-bleeding prevention with both endovascular embolization and surgery, decrease mortality and improve the quality of life of survivors. Nowadays, endovascular embolization has become the method of choice to prevent re-bleeding, especially in Europe, largely replacing surgery.

Objective: The aim of this study was to describe a series of patients with spontaneous subarachnoid hemorrhage treated at the University Hospital of Getafe between 1995 and 2015 (n=343; 185 surgically and 158 endovascular embolization), analyze the diagnosis and treatment, establish main prognosis factors and compare the results of both treatments (clip and coil).

Material and method: We have made a retrospective study, by reviewing medical history, analyzing epidemiological data.

Results: We studied a total of 343 patients, 185 surgery and 158 embolization. It has been shown that embolization, the treatment of choice today, has decreased the number of complications in these years.

Discussion: The long period of time analyzed, 20 years, does not allow a homogeneous study of the sample due to changes in the diagnostic and therapeutic criteria, but allows obtaining interesting results and conclusions.

Conclusion: It has been established that the performance of brain Computed Tomography (CT) and arteriography, and ICU admission, is the best medical management in these patients. In this study, clinical income situation, the amount of blood in CT and the patient's age, have proved the most decisive factors in the final outcome. Endovascular embolization is a safe method of treating ruptured aneurysms.

Keywords: Subarachnoid hemorrhage; Intracranial aneurysm; Therapeutics; Prognosis

Introduction

Aneurysmal Subarachnoid hemorrhage (ASH) is a serious clinical entity, and represents an important cause of mortality and morbidity in industrialized countries, which constitutes a very important effort, both at a human and economic level [1]. ASH is a very serious phenomenon, severe associated with high morbidity and mortality rates. After the initial hemorrhage, 50% of the patients die, and between 30-40% of the patients undergo episode of re-bleeding during the first month [2]. The percentage of deaths due to re-bleeding varies in the different series analyzed between 60% and 75% [1-4]. Currently, the knowledge of the intracranial pathophysiology has advanced a lot, but there are still wide gaps in the understanding of the pathogenesis that limit the development of an optimal therapeutic scheme. In addition, the clinical model of ASH is multifactorial, both in terms of clinical and pathological treatment, so it is very difficult to determine the effect of a pathological or therapeutic variable on the evolution or the final prognosis of patients.

The progress experienced in the management and treatment of patients suffering from ASH allowed a marked improvement in prognosis, achieving a reduction in the mortality rate by up to 30%, and not increasing the number of patients remaining in a persistent vegetative state or developing severe disability, according to the criteria of the Glasgow Evolutionary Scale, although this is still close to 40%. Currently, it is necessary to determine the most influential factors in the final evolution in order to establish prognostic models in order to be able to elaborate effective treatment and rehabilitation plans [5-10]. Considering the foregoing, this work intends to fulfill the following objectives:

1. To gather a wide series of patients who have suffered from ASH and to study their epidemiological profile and clinical presentation form.
2. Design a practical scheme of clinical performance in these patients, as well as describe the risk factors that

determine the probability of developing complications in the group of patients suffering from ASH, studying the main factors that influence clinical evolution.

3. Compare conventional surgical treatment ("clip") with endovascular embolization therapy ("coil").
4. Analyze the different variables that can act on the final forecast, determining which have greater significance and design a prediction model of the initial and late prognosis.

Results

Regarding descriptive epidemiology, 494 clinical histories were reviewed, 151 patients were excluded because they did not meet inclusion criteria or met exclusion criteria, so finally in this study the data of 343 patients with aneurysmal ASH. Of these, 185 have been treated by surgery and 158 with endovascular embolization. The presence of multiple aneurysms was observed in 15.73% of patients (clipping=12.94% vs embolized=21.79%). Of the total of 343 patients, there were 191 women (55.65%) and 152 men (44.35%). The age range ranges from 14 to 82 years with an average age of 50 ± 13 years. The average annual incidence estimated in the study was 1.84 ± 0.63 per 100,000 inhabitants per year (considering the subsidiary health area of this center as 850,000 inhabitants).

No statistically significant differences were found in relation to the percentage of hospital deaths among the patients treated at our center, although there is a trend towards a higher percentage of deaths among surgically treated patients (clipping=8.97% vs embolized=2.94%, $p=0.055$). All the figures analyzed improve with the passage of time in the group of embolized patients, remaining stable in surgical patients.

Discussion

In the present study we have two groups of patients, which could be grouped into those operated surgically and those patients with embolized aneurysms. According to the data obtained, both treatment groups are homogeneous, uniform and comparable, although significant differences can be observed in the aneurysmal location. The ample period of time analyzed, of 20 years, does not allow a homogeneous study of the sample due to the changes experienced in the diagnostic and therapeutic criteria, but it allows obtaining interesting results and conclusions. For example, with the passage of time, the indications for performing cerebral arteriography of control or the timing of the intervention have varied.

Considering the treatment, the clinical guidelines accepted at each moment were followed. We assume that the neurological status of admission is comparable in the two groups, having been determined by the Glasgow scale, despite the relative subjectivity of the observer. The data found in the study on age at which hemorrhage occurs, distribution by sex and presence of multiple aneurysms are superimposable to those published in the scientific literature.

The incidence of ASH collected is lower than expected with respect to data published in both the national and international literature.³ However, it is interesting to see that it has been reported that in certain centers with a high volume of ASH treatment, longer hospital stays are shown [11-16]. These studies argue that this phenomenon could be due to the fact that they group patients of greater age complexity, or because they are less "efficient" centers. According to the opinion of several authors, the possible cause of the long stay is not due to any of the previous theories but to the fact that the collection of all re-admissions penalizes the average global stay compared to other studies, as in our work.

The average stay in ICU collected in the diverse literature analyzed is 1.8 and 1.7 days for those operated and embolized respectively. In the present study, the previously published trend in which embolized patients usually have a shorter stay in the ICU is reversed. Unfortunately, we do not have a sufficiently large sample of patients operated on to be able to affirm this trend.

The logistic regression models allow us to know, introducing the analyzed variables, the "theoretical" final evolutionary prognosis of the patient, and according to this, decide which is the ideal treatment for each patient and the moment of it, always following the principles of "optimism" "Logical that make that all valid therapeutic possibilities are administered, even if the patient is in a bad clinical and neurological situation.¹³⁻¹⁵

Conclusion

- i. Subarachnoid hemorrhage is more frequent in women (1.2/ 1), predominantly in patients whose ages range between 45 and 55 years. The incidence in our environment can be established in 1.5 cases per 100,000 inhabitants per year, below the figures collected in previous studies.
- ii. In the clinical management of subarachnoid hemorrhage, patients can be classified into groups, according to the risk factors they present, according to the neurological situation at admission, the medical history and the images found on CT.
- iii. All patients must be examined with CT, immediately after being received in the hospital, as well as proceed to their admission to the ICU. The measurement of the volume of blood in the CT is useful, being demonstrated that to greater volume of blood, worse prognosis.
- iv. The Glasgow scale for the Coma and the Glasgow Evolutionary scale are well related to the clinical situation and the final prognosis of the patient with ASH.
- v. Endovascular treatment improves the prognosis and overall survival of patients suffering from aneurysmal rupture, being a safe procedure at the present time.
- vi. The most influential parameters in the final prognosis in patients suffering from ASH are, in this order, the

neurological situation at admission, the amount of blood in the skull CT and age.

References

1. Alshafai N, Cusimano MD, Falenchuk O (2013) Global differences in the present and future management of cerebral aneurysms. *World Neurosurg* 80(6): 17-22.
2. Ayling OG, Ibrahim GM, Drake B, Torner JC, Macdonald RL (2015) Operative complications and differences in outcome after clipping and coiling of ruptured intracranial aneurysms. *J Neurosurg* 123(3): 621-628.
3. Rincon F, Rossenwasser RH, Dumont A (2013) The epidemiology of admissions of nontraumatic subarachnoid hemorrhage in the United States. *Neurosurgery* 74(2): 227-229.
4. Darsaut TE, Jack AS, Kerr RS, Raymond J (2013) International Subarachnoid Aneurysm Trial - ISAT part II: study protocol for a randomized controlled trial. *Trials* 29(14): 156.
5. Grasso G, Alafaci C, Macdonald RL (2017) Management of aneurysmal subarachnoid hemorrhage: State of the art and future perspectives. *Surg Neurol Int* 19: 8-11.
6. Jaja BN, Lingsma H, Schweizer TA, Thorpe KE, Steyerberg EW, et al. (2015) Prognostic value of premorbid hypertension and neurological status in aneurysmal subarachnoid hemorrhage: pooled analyses of individual patient data in the SAHIT repository. *J Neurosurg* 122(3): 644-652.
7. Konczalla J, Seifert V, Beck J, Güresir E, Vatter H, Raabe A, Marquardt G. Outcome after Hunt and Hess Grade V subarachnoid hemorrhage: a comparison of pre-coiling era (1980-1995) versus post-ISAT era (2005-2014). *J Neurosurg* 128(1): 100-110.
8. Korja M, Kivisaari R, Rezai Jahromi B, Lehto H (2017) Natural History of Ruptured but Untreated Intracranial Aneurysms. *Stroke* 48(4): 1081-1084.
9. Lo BW, Fukuda H, Nishimura Y, Farrokhyar F, Thabane L, et al. (2015) Systematic review of clinical prediction tools and prognostic factors in aneurysmal subarachnoid hemorrhage. *Surg Neurol Int* 11(6): 135.
10. Molyneux AJ, Birks J, Clarke A, Sneade M, Kerr RS (2015) The durability of endovascular coiling versus neurosurgical clipping of ruptured cerebral aneurysms: 18 year follow-up of the UK cohort of the International Subarachnoid Aneurysm Trial (ISAT). *Lancet* 385(9969): 691-697.
11. Pegoli M, Mandrekar J, Rabinstein AA, Lanzino G (2015) Predictors of excellent functional outcome in aneurysmal subarachnoid hemorrhage. *J Neurosurg* 122(2): 14-18.
12. Song JP, Ni W, Gu YX, Zhu W, Chen L, et al. (2017) Epidemiological Features of Nontraumatic Spontaneous Subarachnoid Hemorrhage in China: A Nationwide Hospital-based multicenter Study. *Chin Med J (Engl)* 130(7): 776-781.
13. Spetzler RF, McDougall CG, Zabramski JM, Albuquerque FC, Hills NK, et al. (2015) The Barrow Ruptured Aneurysm Trial: 6-year results. *J Neurosurg* 123(3): 609-617.
14. Tarnutzer AA, Lee SH, Robinson KA, Wang Z, Edlow JA, et al. (2017) ED misdiagnosis of cerebrovascular events in the era of modern neuroimaging: A meta-analysis. *Neurology* 88(15): 1468-1477.
15. Wesali S, Persson HC, Cederin B, Sunnerhagen KS (2017) Improved survival after non-traumatic subarachnoid haemorrhage with structured care pathways and modern intensive care. *Clin Neurol Neurosurg* 31(138): 52-58.
16. Xia ZW, Cheng Q (2017) Strategy for poor grade aneurysmal subarachnoid haemorrhage. *Eur J Neurol* 24(4): e23.



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/CTBEB.2018.13.555861](https://doi.org/10.19080/CTBEB.2018.13.555861)

Your next submission with Juniper Publishers will reach you the below assets

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission
<https://juniperpublishers.com/online-submission.php>