

Features of Relationships between Temperamental Characteristics in Male and Female Adolescents



Liudmyla D Popova*, Irina M Vasylyeva and Oxana A Nakonechnaya

Kharkiv National Medical University, Ukraine

Submission: February 14, 2017; **Published:** March 27, 2017

***Corresponding author:** Liudmyla D Popova, Kharkiv National Medical University, Ukraine, Tel: +380984285463; Email: popova_ld@ukr.net

Abstract

Aggression, including aggression of adolescents, is a major problem in modern society, but the mechanisms of its development are not clear up today. Researches of mechanisms of aggression development in women are virtually absent. We studied the relationships between temperamental characteristics in male and female adolescents. Aggressiveness index, physical, verbal aggressions, irritability, anxiety, extraversion and neuroticism levels were estimated using Buss-Durkee Hostility Inventory, Spielberger State-Trait Anxiety Inventory, Eysenck Personality Inventory. Higher neuroticism, anxiety levels and irritability were found in female adolescents compared with male adolescents, that perhaps due to a greater activity of the hypothalamic pituitary adrenal axis in women. Significant positive correlations between neuroticism and anxiety, and high negative correlations between anxiety and verbal aggression were revealed in male adolescents. Significant positive correlations between neuroticism and physical aggression, neuroticism and irritability, irritability and verbal aggression, verbal aggression and extraversion were revealed in female adolescents. Significant positive correlations between neuroticism and physical aggression, neuroticism and irritability, irritability and verbal aggression in female adolescents make it possible to suggest that female adolescents are more prone to impulsive aggression than male adolescents.

Keywords: Aggression; Extraversion; Anxiety; Neuroticism; Male and Female; Adolescents

Introduction

Aggression, including aggression of adolescents, is a major problem in modern society, but the mechanisms of its development are not clear up today. Researches of mechanisms of aggression development in women are virtually absent. Aggression is more pronounced in men than in women, so the study of mechanisms of aggression development, mainly conducted on men and male animals. Researches on human confined to conducting psychological tests, blood tests and Magnetic Resonance Imaging of brain. Most studies were performed on males of different animals. It is shown that male sex hormones facilitate the manifestation of aggressive behavior in adults [1]. In the embryonic period they are involved in the formation of neural networks [2]. Testosterone, which is synthesized by fetal gonads, diffuses to the brain of males, where it is locally aromatized in estradiol and then initiates a process of masculinization [3]. In puberty it triggers a second period of structural reorganization and plasticity in the brain [4]. In adults, testosterone is involved in the modulation of neural networks that regulate aggression [5]. Many mediators are involved in aggression development [6-10]. Specific neurotransmitter systems involved in mammalian aggression include serotonin, dopamine, norepinephrine, GABA, and neuropeptides such as

vasopressin and oxytocin [11]. Their investigation on human's brain is impossible, but temperamental characteristics reflect the features of neuromediator status of person. The study of relationships between temperamental characteristics in males and females of different age groups is important for elucidation of aggression mechanisms development. The aim of our work was the study of relationships between temperamental characteristics in male and female adolescents.

Methods

The study involved 34 adolescents within the age range 15-16 years (17 males and 17 females). The work was done in accordance with order № 319 of the Ministry of Education and Science of Ukraine of 18.03.2015 and carried out with the ethical standards.

Aggressiveness index, physical, verbal aggressions, irritability, anxiety, extraversion and neuroticism levels were estimated using Buss-Durkee Hostility Inventory, Spielberger State-Trait Anxiety Inventory, Eysenck Personality Inventory. Buss-Durkee Hostility Inventory is used to study aggression. Eysenck Personality Inventory provides the estimation of extraversion-introversion and emotional stability-instability (neuroticism). The levels of extraversion and neuroticism

were assessed in points. Aggressiveness index, physical, verbal aggressions, irritability were estimated in a percentage of the maximum level. Eysenck Personality Inventory provides to estimate the sincerity of answers. If the answers have not been sincere, they were not taken into account. Statistical analysis of the results was carried out by methods of nonparametric statistics using the package "Statistica 6.0", Mann-Whitney test and correlation analysis according to Spearman.

Results




Figure 1: Neuroticism level in male and female adolescents (in points) (Me, [25%; 75%], minimum and maximum values; $p=0.00225$).




Figure 2: Anxiety level in male and female adolescents (in points) (Me, [25%; 75%], minimum and maximum values; $p=0.012858$).




Figure 3: Irritability level in male and female adolescents (in points) (Me, [25%; 75%], minimum and maximum values; $p=0.00828$).

To identify gender-specific relationships between temperamental characteristics we investigated temperamental characteristics in male and female adolescents. According to the results, they did not differ in the levels of extraversion, physical and verbal aggressions, and aggression index. But higher levels of neuroticism, anxiety and irritability were found in female adolescents compared with male adolescents (Figure 1-3). Significant positive correlations between neuroticism and anxiety, and high negative correlations between anxiety and verbal aggression were revealed in male adolescents (Table 1).

Table 1: Correlation coefficients between temperamental characteristics in male adolescents.

	Neuroticism	Anxiety	Verbal Aggression
Neuroticism	-	+0.55*	-0.22
Anxiety	+0.55*	-	-0.62*
Verbal aggression	-0.22	-0.62*	-

*Correlation coefficient is statistically significant, $p < 0.05$

Table 2: Correlation coefficients between temperamental characteristics in female adolescents.

	Extraversion	Neuroticism	Physical Aggression	Verbal Aggression	Irritability
Extraversion	-	+0.23	+0.14	+0.57*	+0.22
Neuroticism	+0.23	-	+0.58*	+0.48	+0.77*
Physical aggression	+0.14	+0.58*	-	+0.28	+0.21
Verbal aggression	+0.57*	+0.48	+0.28	-	+0.54*
Irritability	+0.22	+0.77*	+0.21	+0.54*	-

*Correlation coefficient is statistically significant, $p < 0.05$

Significant positive correlations between neuroticism and physical aggression, neuroticism and irritability, irritability and verbal aggression, verbal aggression and extraversion were revealed in female adolescents (Table 2).

Discussion

According to our results male and female adolescents did not differ by aggression index, physical and verbal aggressions. Usually adolescent are characterized by immature and impulsive behavior and neurobehavioral excitement due to maturation of adolescent brain [10]. Glutamatergic neurotransmission predominates, whereas gamma-aminobutyric acid transmission is in formation process [10].

Higher levels of neuroticism, anxiety and irritability were found in female adolescents compared with male adolescents. Elevated levels of neuroticism and trait anxiety in female adolescents may be associated with higher levels of cortisol in women than in men [12]. This is related to testosterone organizing role in the formation of hypothalamic pituitary adrenal axis [12] and masculinization of the brain during fetal development [3]. Organizational effects of testosterone occur pre- and perinatally. Testosterone provides masculinizing and defeminizing neural circuits in males, and the absence of testosterone leads to female neural phenotype [4].

A greater reactivity in the adrenal response to adrenocorticotrophic hormone is observed in women [13]. Basal corticotropin-releasing hormone expression is lower in males, than in females [14]. Corticotropin-releasing factor regulates instinctive forms of emotional behavior (fear, anxiety, frustration and deliverance from them) [15]. Exposure to stress is associated with an increase in firing of locus coeruleus and with increased release and turnover of norepinephrine in brain regions which receive noradrenergic innervations [16]. Almost half of noradrenergic fibers begins in locus coeruleus [17]. The locus coeruleus has dense excitatory projections to cortex, hippocampus, amygdale, thalamus, hypothalamus, serotonergic neurons of the dorsal raphe nucleus [16]. These features of noradrenergic neuronal system determine its adaptability to rapid and global modulation of brain functions in response to environmental change as it comes under stress. Increased activity of locus coeruleus is associated with behavioral manifestations of fear [18]. Disorders induced by stress, such as depression and anxiety are more common among women than among men [19]. The increased anxiety in female adolescents may be also due to features of connections between amygdale and hippocampus. A stronger coupling between the left hippocampus and the left amygdale is observed in women than in men. A higher coupling of hippocampus and amygdale in women may act as a neural contributing factor to sex-specific prevalence rates in anxiety and panic disorders [20].

Neuroticism is characterized by increased emotionality, impulsivity, a tendency to irritability. Irritability belongs to aggression-related behavioral states. This state is characterized

by reduced control over a temper and usually results in verbal or behavioral outbursts [21]. Significant positive correlations between neuroticism and physical aggression, neuroticism and irritability, irritability and verbal aggression in female adolescents make it possible to suggest that female adolescents are more prone to impulsive aggression than male adolescents.

Conclusion

1. Higher neuroticism, anxiety levels and irritability index were found in female adolescents compared with male adolescents, that perhaps due to a greater activity of the hypothalamic pituitary adrenal axis in women.
2. Significant positive correlations between neuroticism and physical aggression, neuroticism and irritability, irritability and verbal aggression in female adolescents make it possible to suggest that female adolescents are more prone to impulsive aggression than male adolescents.

References

1. Chichinadze KN, Dominiadze TR, Matitaischvili T, Chichinadze NK, Lazaraschvili AG (2010) Is the blood plasma testosterone level linked with aggressive behaviour of prisoner men? *Bul Exp Biol Med* 149(1): 11-13.
2. Wingfield JC, Lynn S, Soma KK (2001) Avoiding the 'costs' of testosterone: ecological bases of hormone-behavior interactions. *Brain Behav* 57(5): 239-251.
3. Roselli CE, Liu M, Hurn PD (2009) Brain aromatization. Classical roles and new perspectives. *Semin. Reprod. Med* 27(3): 207-217.
4. Blakemore SJ, Burnett S, Dahl RE (2010) The role of puberty in the developing adolescent brain. *Hum Brain Mapp* 31(6): 926-933.
5. Neumann ID, Veenema AH, Beiderbeck DI (2010) Aggression and anxiety: social context and neurobiological links. *Front Behav Neurosci* 4: 1-12.
6. Vegt BJ, Lieuwes N, Cremers TI, de Boer SF, Koolhaas JM (2003) Cerebrospinal fluid monoamine and metabolite concentrations and aggression in rats. *Horm Behav* 44(3): 199-208.
7. Gaalen MM, Brueggeman RJ, Bronius PF, Schoffelman AN, Vanderschuren LJ (2006) Behavioral disinhibition requires dopamine receptor activation. *Psychopharmacology (Berl)* 187(1): 73-85.
8. Pavlov KA, Chistiakov DA, Chekhonin VP (2012) Genetic determinants of aggression and impulsivity in humans. *J Appl Genet* 53(1): 61-82.
9. Popova NK (2008) From gene to aggressive behavior: the role of brain serotonin. *Neurosci Behav Physiol* 38(5): 471-475.
10. Arain M, Haque M, Johal L, Mathur P, Nel W, et al. (2013) Maturation of the adolescent brain. *Neuropsychiatr Dis Treat* 9: 449-461.
11. Yanowitch R, Coccaro EF (2011) The neurochemistry of human aggression. *Adv Genet* 75: 151-169.
12. Swaab DF, Bao AM, Lucassen PJ (2005) The stress system in the human brain in depression and neurodegeneration. *Ageing Res Rev* 4(2): 141-194.
13. Arnetz L, Rajamand Ekberg N, Brismar K, Alvarsson M (2015) Gender difference in adrenal sensitivity to ACTH is abolished in type 2 diabetes. *Endocr Connect* 4(2): 92-99.
14. Raison CL, Miller AH (2003) When not enough is too much: the role of insufficient glucocorticoid signaling in the pathophysiology of stress-related disorders. *Am J Psychiatry* 160(9): 1554-1565.

15. Shabanov PD, Lebedev AA (2012) Neurochemical mechanisms of the nucleus accumbens realizing the reinforcing effects of self-stimulation on the lateral hypothalamus. *Med Acad Journal* 12(2): 68-76.
16. Bremner JD, Krystal JH, Southwick SM, Charney DS (1996) Noradrenergic mechanisms in stress and anxiety: Preclinical studies. *Synapse* 23(1): 28-38.
17. Prokopova I (2010) Noradrenalin and behavior. *Cesk Fysio* 59(2): 51-58.
18. Samuels ER, Zabadi ES (2008) Functional neuroanatomy of the noradrenergic locus coeruleus: its roles in the regulation of arousal and autonomic function. Part I: Principles of functional organization. *Curr Neuropharmacol* 6(3): 235-253.
19. Weiser JM, Goel N, Sandau SU, Bale TL, Handa RG (2008) Androgen regulation of corticotropin-releasing hormone receptor 2 (CRHR2) mRNA expression and receptor binding in the rat brain. *Exp Neurol* 214(1): 62-68.
20. Kogler L, Müller VI, Seidel EM, Boubela R, Kalcher K, et al. (2016) Sex differences in the functional connectivity of the amygdalae in association with cortisol. *Neuroimage* 134: 410-423.
21. Picciotto MR, Lewis AS, van Schalkwyk GI, Mineur YS (2015) Mood and anxiety regulation by nicotinic acetylcholine receptors: A potential pathway to modulate aggression and related behavioral states. *Neuropharmacology* 96(Pt B): 235-243.



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/CTCMI.2017.01.555560](https://doi.org/10.19080/CTCMI.2017.01.555560)

Your next submission with Juniper Publishers will reach you the below assets

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission

<https://juniperpublishers.com/online-submission.php>