

Research Article

Volume 3 Issue 2 - June 2019
DOI: 10.19080/CTCMI.2019.03.555610

Curr Trends Clin Med Imaging

Copyright © All rights are reserved by Huynh Quang Huy

Computed Tomography finding of Intraabdominal Ruptured Liver Abscess



Huynh Quang Huy^{1*}, Le Cong Tri² and Pham Hong Duc³

¹Radiology Department, Pham Ngoc Thach University of Medicine and HCMC Oncology Hospital, Vietnam

²Surgery Department, Cho Ray hospital, Vietnam

³Diagnostic Imaging Department, Ha Noi University of Medicine

Received date: May 22, 2019; **Published date:** June 11, 2019

***Corresponding author:** Huynh Quang Huy, Radiology Department, Pham Ngoc Thach University of Medicine and HCMC Oncology Hospital, District 10, Ho Chi Minh city, Vietnam

Abstract

Objectives: The purpose of this study is to describe the computed tomography of intra abdominal ruptured liver abscess.

Patients and Methods: The computed tomography findings of 32 patients who had a confirmed intraabdominal ruptured liver abscess were analyzed retrospectively at Cho Ray hospital between 2014 and 2018.

Results: 32 patients with a median age of 53.3 ± 15.3 years (range, 24–85 years). There were more men than women in the study, with a male-to-female ratio of 1.9:1. Most of the patients were suffering from abdominal pain (96.9%), malaise and anorexia (96.9%), fever (78.1%), peritonitis (78.1%). The mean size of the liver abscess was 8.4 cm (range, 4.0–14 cm). A single abscess was found in 11 patients, and multiple abscesses were seen in 21 patients. Unilobar involvement was seen in 29 patients, with the right lobe affected more commonly (68.8%, 22 of 32). The abscesses were completely liquefied in 25 patients and gas in the abscess cavity in 7 patients. Free intraperitoneal fluid was seen in all patients.

Conclusion: A knowledge of intraabdominal ruptured liver abscess is important for an early diagnosis and appropriate management.

Keywords: Liver; Abscess; Ruptured; Computed Tomography; Clinical

Introduction

Liver abscess is a common condition in tropical countries and is associated with significant morbidity and mortality. Traditionally, there are two major classifications of hepatic abscess; pyogenic and amoebic [1, 2]. There are various complications associated with hepatic abscesses, of which, rupture of the abscess is the most common [3-5]. Intraperitoneal rupture of liver abscess is a rare but potentially fatal disease, often involving the elderly, who are commonly of poor surgical risk with background of significant medical illness [6,7]. Accurate preoperative diagnosis is difficult and often necessitates exploratory laparotomy for peritonitis [8].

Computed tomography (CT) is an ideal tool for diagnosing hepatic abscesses, and the sensitivity of CT for detecting hepatic abscesses is as high as 97% [9]. On CT, an hepatic abscess appears as a single or multiloculated mass with low attenuation [9]. Early diagnosis and prompt therapy are essential to reduce the morbidity and mortality associated with a pyogenic hepatic abscess. The mortality rate is low when the abscess is confined to the liver; however, the mortality rate is increased when

the hepatic abscess extends into the chest, peritoneal cavity, or pericardial cavity [10]. Therefore, the early detection of complications associated with hepatic abscesses is important. In hepatic abscess, various complications have been described [11-13], and the rate of complications was reported to be 10.3% [10]. Reports of complications in pyogenic hepaticabscess are rare, however, and the imaging appearance has been studied in only a small number of patients [13,14]. This article presents the CT and clinical findings of our patients with intraabdominal ruptured liver abscess.

Patients and Methods

This is a retrospective observational study of patients with intraabdominal ruptured liver abscess presenting to Cho Ray hospital between 2014 and 2018. The study was approved by the Hospital Ethical Board. All diagnosed cases of ruptured liver abscess based on radiology and laparoscopic investigation were included in the study. Details of demographics (age, sex), clinical features, computed tomography were recorded. Data will be analyzed using SPSS software.

All patients underwent contrast enhanced CT of the liver before laparoscopic drainage of the liver abscess. In some patients, CT studies were also performed during follow-up to assess the size of the abscess cavity, monitor complications. However, we only reviewed the contrast-enhanced CT images obtained before drainage of the liver abscess, for the purpose of this study. The CT examinations were performed using Somatom Sensation 64, Siemens, Germany. The scanning parameters varied during the study period and with different scanners: collimation ranged from 1.25 mm to 7 mm; pitch ranged from 0.75 to 1.5; section thickness ranged from 1 mm to 5 mm. Liver examinations were conducted using 100 ml of intravenous non-ionic iodinated contrast medium (Ultravist, Schering, Berlin, Germany) and dosage was calculated based on the patients' weight and administered via a power injector at a rate of 3 mL/sec. Axial sections of 3-5 mm thickness were reconstructed, reported, and archived.

The scans were reviewed by two radiologists who reached agreement between them. The following features were recorded: (a) lobe involvement (unilobar [right or left] or bilobar); (b) number of abscesses (single or multiple); (c) maximal abscess diameter, with the largest abscess measured when there were multiple abscesses; (d) unilocular or multilocular (presence of ≥ 1 -mm-thick septations), multilocular abscess; (e) solid or cystic appearance ($>50\%$ of the abscess cavity appears hypodense or liquefied, with an attenuation value of ≤ 20 HU) in most of the sections showing the abscess cavity, cystic abscess; (f) gas within the abscess cavity; and (g) spontaneous rupture of the abscess (based on CT and clinical symptoms).

Results

The study group consisted of 32 patients with a median age of 53.3 ± 15.3 years (range, 24–85 years). There were more men than women in the study, with a male-to-female ratio of 1.9:1. Most of the patients were suffering from abdominal pain (96.9%), malaise and anorexia (96.9%), fever (78.1%), peritonitis (78.1%), and other positive clinical signs showed in Table 1.

Table 1: Clinical presentations of intraabdominal ruptured liver absceses.

Clinical Presentations	Frequency	Percentage (%)
Abdominal pain	31	96.9
Malaise and anorexia	31	96.9
Nausea / vomiting	20	62.5
Fever $> 38^{\circ}\text{C}$	25	78.1
Jaundice	5	15.6
Peritonitis	25	78.1
Hepatomegaly	4	12.5

The mean time interval between the onset of fever and CT was 8.3 days (range, 1–30 days). The mean size of the liver abscess was 8.4 cm (range, 4.0–14 cm). A single abscess was found in 11 patients, and multiple abscesses were seen in 21 patients. Unilobar involvement was seen in 29 patients, with

the right lobe affected more commonly (68.8%, 22 of 32). The abscesses were completely liquefied in 25 patients and gas in the abscess cavity in 7 patients. Free intraperitoneal fluid was seen in all patients. The liver CT characteristics of intraabdominal ruptured liver abscess was showed in Table 2, Figures 1 & 2.

Table 2: CT characteristics of intraabdominal ruptured liver abscess.

CT Findings	All Patients (n=32)
Abscess size (cm)	8.4 (4.0–14)
No. of abscesses	
1	11 (34.4)
>1	21 (65.6)
Lobar involvement	
Unilobar	29 (90.6)
Bilobar	3 (9.4)
Abscess appearance	
Completely liquefied	25 (78.1%)
Gas in the abscess cavity	7 (21.9)
Peritoneal effusion	32 (100%)




Figure 1: A 71-year-old woman with rupture of a pyogenic hepatic abscess into the peritoneal cavity resulting in peritonitis. The abscess is located in the right hepatic lobe with gas inside.



Figure 2: A 62-year-old man with rupture of a pyogenic hepatic abscess forming a perihepatic abscess. An enhanced helical computed tomography scan shows a heterogeneously hypodense abscess in the left hepatic lobe.

Discussion

Liver abscess is an important tropical gastrointestinal disorder [15,16]. Liver abscess can be classified into pyogenic and amebic, both having its serious implications, especially when presented late. In developing countries, it forms a major cause for mortality and morbidity [17,18]. With the advent of modern radiological modalities, diagnosis of hepatic abscess is possible in early stages resulting in nonsurgical management; however, fraction of patients either due to late presentation or refractory disease presents with ruptured liver abscess thereby increasing the mortality, presents with fatal disease course, and requires surgical intervention at the earliest [8,19].

Intraperitoneal rupture is one of the serious complication of liver abscesses [5-21]. The frequency of intraperitoneal rupture of an liver abscess varies from 2.5% to 17% [22]. Clinically, increasing hepatic tenderness should indicate an impending rupture [22]. Computed tomography may provide important information regarding the extent of intraperitoneal spread of the liver abscess. Dal Mo Yang et al. [23] study 81 patients who had a confirmed pyogenic hepatic abscess, the complication was encountered in 3 patients (3.7%), and 2 types could be recognized: a loculated perihepatic abscess ($n = 2$) and diffuse peritonitis ($n = 1$). In diffuse peritonitis secondary to rupture of a hepatic abscess, a large amount of ascites and diffuse thickening of the parietal peritoneum could be seen on CT. Although an amebic peritonitis secondary to rupture of a hepatic amebic abscess showed a poor survival rate despite surgical intervention, all the patients with intraperitoneal rupture of the pyogenic hepatic abscess could be treated with percutaneous drainage and antibiotic treatment.

The most common presentation was abdominal pain (96.9%), malaise and anorexia (96.9%), fever (78.1%), peritonitis (78.1%), nausea/vomiting (62.5%), jaundice and hepatomegaly (15.6% and 12.5%, respectively). This results were comparable to study by Hind S. Alsaif with the most common presentation was fever and/or chills, followed by gastrointestinal symptoms (eg, gastrointestinal upset, diarrhea, vomiting, nausea, discomfort, pain), respiratory symptoms (eg, cough, dyspnea, chest distress), and jaundice [24]. The CT appearance of liver abscess is variable and nonspecific. One or more round or oval low-density lesions 2-1 6 cm in diameter may be seen. The margin of the abscess may be smooth or nodular, and one or more internal septations may be present. An enhancing wall is common but not universal. However, the wall may not be apparent if unenhanced CT is not performed. Thus, the CT differential diagnosis of amebic liver abscess in the adult includes simple hepatic cyst, infected or hemorrhagic cyst, pyogenic liver abscess, echinococcal cyst, hematoma, biloma, cystic or necrotic hepatic metastasis, undifferentiated embryonal sarcoma, and biliary cystadenoma. In the few cases in which a rim of edema is seen peripheral to the lesion, diagnostic consideration may be limited to inflammatory conditions [25]. A feature of amebic liver abscess that may aid in

distinguishing it from other focal hepatic lesions is its tendency to extend beyond the surface of the liver.

Our results of liver CT characteristics were comparable to other studies. Alsaif HS et al showed that the median time interval between the onset of fever and CT was 3 days (range, 1-24 days). The median size of the liver abscess was 7 cm (range, 1.7-14 cm). A single abscess was found in 95 patients, and multiple abscesses were seen in 36 patients. Unilobar involvement was seen in 100 patients, with the right lobe affected more commonly (55.0%, 72 of 131). The abscesses were predominantly solid and masslike in 67 patients and cystic in 64 patients. Multilocular appearance of the abscesses was present in 115 patients (87.8%). Thrombophlebitis was present in 30 patients; pylephlebitis, in three; gas in the abscess cavity, in 17; and pneumobilia, in six. Three patients had both gas in the abscess cavity and pneumobilia [24]. Sutdy by Alexopoulou A et al, the majority (69.7%) of liver abscess involved the right lobe, they were 5-9 cm in size (63.6%) and solitary (75.7%). Multiple abscesses were observed in 24.2% of patients.

The presence of elevated hemidiaphragm (42.4%), pleural effusion (18.2%) and basilar infiltrate (6%) in chest radiography was also noted. Liver abscess was accompanied by pylephlebitis in 1 patient. Gas forming were observed in 4 patients (12.1%) and loculated subcollections in the abscess in 3 (9%) [3]. A single abscess was found in 54 (81.8%) patients, and multiple abscesses were seen in 12 (18.2%) patients in the study of Chang Z et al [26]. Among the patients with multiple abscesses, the average number of lesions per patient was 2.5 - 0.67. Unilobar involvement was seen in 54 (81.8%) patients, and multilocular abscess was present in 50 (75.8%) patients. The abscesses were predominantly solid in 38 (57.6%) patients and cystic in 28 (42.4%) patients. Thrombophlebitis was present in 9 (13.6%) patients, and gas in the abscess cavity was present in 11 patients (16.7%). There were 4 (6.1%) patients with liver abscess who experienced a spontaneous rupture of the abscess [26].

A definitive diagnosis of liver abscess requires imaging, with both sonography and CT being useful. Sonography is operator-dependent and the operator may have difficulty picking out a small, solitary abscess. Its sensitivity is around 79%, compared with 98% for CT [27], but emergency bedside sonography may be very useful in making a rapid diagnosis. Kim et al. suggested certain CT characteristics suggestive of a *Klebsiella* abscess, such as a hairball sign or air-fluid level [28]. However, while such imaging distinctions may be sought, culture and sensitivity results are the key to choosing the appropriate antibiotics. The advancement in radiologic techniques has been credited with the improvement in mortality rates. Ultrasonography and CT evaluation with contrast remain the radiologic modalities of choice in screening procedures and also can be used as techniques for guiding percutaneous aspiration and drainage. With advancement in multidetector CT scan technology, image quality has improved dramatically, allowing for improved

detection. Besides, gallium and technetium radionucleotides can be used in diagnosis.

Conclusion

A ruptured hepatic abscess is more life-threatening than an unruptured one. A rapid and accurate diagnosis and prompt surgical intervention are essential. Computed tomography is an ideal tool for diagnosing hepatic abscesses and its complications. The authors are grateful to physicians, administrative staff at Cho Ray Hospitals for allowing us to undertake this research.

References

1. Akhondi H, Sabih DE, Liver Abscess, in StatPearls. 2019: Treasure Island (FL).
2. Chia DWJ, Kuan WS, Ho WH, Sim TB & Chua MT (2019) Early predictors for the diagnosis of liver abscess in the emergency department. *Intern Emerg Med*.
3. Alexopoulou A, Dimopoulos H, Soultati A, Panetsos G, Dourakis SP (2010) Factors related to complications and mortality in pyogenic liver abscesses. *Annals of Gastroenterology* 23(4): 296-301.
4. Mukhopadhyay M, Saha AK, Sarkar A, Mukherjee S (2010) Amoebic liver abscess: presentation and complications. *Indian J Surg* 72(1): 37-41.
5. Satish KR, Sathyaranayana BA, Madhu SL, Nataraj NR, Amit GM (2015) A study of predictors for identification of risk of complications in patients with liver abscess. *Trop Gastroenterol* 36(2): 96-100.
6. Elechi EN, Etawo SU (1991) Management of intraperitoneal rupture of amoebic liver abscess: a 6-year experience with 11 cases in Port Harcourt, Nigeria. *Trop Doct* 21(1): 43-44.
7. Sarda AK, Bal S, Sharma AK, Kapur MM (1989) Intraperitoneal rupture of amoebic liver abscess. *Br J Surg* 76(2): 202-203.
8. Memon AS, Siddiqui FG, Memon HA, Ali SA (2010) Management of ruptured amoebic liver abscess: 22-years experience. *J Ayub Med Coll Abbottabad* 22(2): 96-99.
9. Halvorsen RA, Korobkin M, Foster WL, Silverman PM, Thompson WM (1984) The variable CT appearance of hepatic abscesses. *AJR Am J Roentgenol* 142(5): 941-946.
10. Munoz LE, Botello MA, Carrillo O, Martinez AM (1992) Early detection of complications in amebic liver abscess. *Arch Med Res* 23(2): 251-253.
11. Mondragon-Sanchez R, Cortes-Espinoza T, Sanchez-Cisneros R, Parra-Silva H, Hurtado-Andrade H (1994) Rupture of an amebic liver abscess into the pericardium. Presentation of a case and review of current management. *Hepatogastroenterology* 41(6): 585-588.
12. Mowji PJ, Cohen AJ, Potkin B & Viltuznik J (1987) Amebic liver abscess with hepatoduodenal fistula. *Am J Gastroenterol* 82(6): 558-559.
13. Tandon N, Karak PK, Mukhopadhyay S & Kumar V (1991) Amoebic liver abscess: rupture into retroperitoneum. *Gastrointest Radiol* 16(3): 240-242.
14. Yamada S, Maruo H, Mori K & Kosaka A (1996) [An unusual case of pyogenic hepatic abscess rupturing into the pericardial cavity]. *Nihon Kyobu Geka Gakkai Zasshi* 44(11): 2072-2075.
15. Cook GC (1991) Gastroenterological emergencies in the tropics. *Baillieres Clin Gastroenterol* 5(4): 861-886.
16. Reeder MM (1975) Tropical diseases of the liver and bile ducts. *Semin Roentgenol* 10(3): 229-243.
17. Blessmann J, Van Linh P, Nu PA, Thi HD, Muller-Myhsok B (2002) Epidemiology of amebiasis in a region of high incidence of amoebic liver abscess in central Vietnam. *Am J Trop Med Hyg* 66(5): 578-583.
18. Kannathasan S, Murugananthan A, Kumanan T, de Silva NR, Rajeshkannan N (2018) Epidemiology and factors associated with amoebic liver abscess in northern Sri Lanka. *BMC Public Health* 18(1): 118.
19. Short M & Desai AP (2008) Laparoscopy and transdiaphragmatic thoracoscopy in management of ruptured amoebic liver abscess. *J Laparoendosc Adv Surg Tech A* 18(3): 473-476.
20. Adams EB & MacLeod IN (1977) Invasive amebiasis. II. Amoebic liver abscess and its complications. *Medicine (Baltimore)* 56(4): 325-334.
21. Jolobe OM (2010) Intrahepatic expansion of amoebic liver abscess can also generate life-threatening complications. *South Med J* 103(11): 1189-1090.
22. Eggleston FC, Handa AK, Verghese M (1982) Amebic peritonitis secondary to amoebic liver abscess. *Surgery* 91(1): 46-48.
23. Yang DM, Kim HN, Kang JH, Seo TS, Park CH (2004) Complications of pyogenic hepatic abscess: computed tomography and clinical features. *J Comput Assist Tomogr* 28(3): 311-317.
24. Alsaif HS, Venkatesh SK, Chan DS, Archuleta S (2011) CT appearance of pyogenic liver abscesses caused by *Klebsiella pneumoniae*. *Radiology* 260(1): 129-138.
25. Gabata T, Kadoya M, Matsui O, Kobayashi T, Kawamori Y (2001) Dynamic CT of hepatic abscesses: significance of transient segmental enhancement. *AJR Am J Roentgenol* 176(3): 675-679.
26. Chang Z, Zheng J, Ma Y, Liu Z (2015) Analysis of clinical and CT characteristics of patients with *Klebsiella pneumoniae* liver abscesses: an insight into risk factors of metastatic infection. *Int J Infect Dis* 33: 50-54.
27. Pearl R, Pancu D, Legome E (2005) Hepatic abscess. *J Emerg Med* 28(3): 337-339.
28. Kim SB, Je BK, Lee KY, Lee SH, Chung HH (2007) Computed tomographic differences of pyogenic liver abscesses caused by *Klebsiella pneumoniae* and non-*Klebsiella pneumoniae*. *J Comput Assist Tomogr* 31(1): 59-65.



This work is licensed under Creative
Commons Attribution 4.0 License
DOI: [10.19080/CTCMI.2019.03.555610](https://doi.org/10.19080/CTCMI.2019.03.555610)

**Your next submission with Juniper Publishers
will reach you the below assets**

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission

<https://juniperpublishers.com/online-submission.php>