

5-Azadeoxycytidine and Trichostatin A in FMR1 Gene Reactivation



Alexander A Dolskiy, Irina S Kolesnikova and Dmitry V Yudkin*

Department of Chromosome Pathology, Institute of Molecular and Cellular Biology, Russia

Submission: February 20, 2017; **Published:** March 20, 2017

***Corresponding author:** Dmitry V Yudkin, Chromosome Pathology Group, Institute of Molecular and Cellular Biology SB RAS, Lavrentievave 8/2, Novosibirsk, 630090, Russia, Tel: +7-383-363-90-62; Email: dim@mcb.nsc.ru

Mini Review

Here we review the ways of chromatin modifications by 5-azadeoxycytidine and trichostatin A for gene reactivation. Therapeutic effects for different diseases treatment are discussed.

Many existing diseases are caused by suppression of specific gene expression. As an example, for oncological diseases these are tumor suppressor genes [1]. In case of Fragile-X syndrome, expression of FMR1 gene is suppressed. Nowadays numerous studies to search for any ways to reactivate genes using small molecules like 5-azadeoxycytidine, and trichostatin A which act on the whole genome are performed. Mechanisms, advantages and disadvantages of these methods are briefly described below.

First of all, we have to look closer the gene silencing system using FMR1 gene as an example to explain the gene reactivation mechanism. Methylation of CpG islands in promoter area is involved in regulation of gene expression and cell differentiation. Repeat expansion in FMR1 gene promoter area up to more than 200 triplets leads to recognition of CpG islands by DNA-methyltransferases (DNMT), it add methyl groups to cytosines in CGG-repeats, this serves as a signal for hetero chromatinization and for switching off the gene. Methylated cytosine becomes recognized by MeCP2 protein, who activates histone deacetylases (HDAC), which further remove acetyl groups from N-terminal domain of H3 and H4 histones at the methylated area, and this leads to chromatin condensation [2].

A number of studies showed a possibility of methylated FMR1 reactivation using 5-azadeoxycytidine-a cytidine analog. During DNA chain elongation 5-azadeoxycytidine becomes inserted in instead of cytidine and an irreversible covalently binding with DNA-methyltransferase 1 (DNMT1) occurs, because this enzyme tries to methylate a trivalent nitrogen in 5 position of pyrimidine ring. Thereafter during cell division a passive demethylation process takes place [3].

Simultaneous treatment of cell lines with 5-azadeoxycytidine and histone deacetylases inhibitors has led to greater gene

reactivation. So, simultaneous treatment of mammal gland cell lines with 5-azadeoxycytidine, butyrate and trichostatin A led to more efficient FMR1 reactivation than when only 5-azadeoxycytidine was used [4]. It should be noted that these substances can reactivate tumor suppressors too [5]. When treated simultaneously with 5-azadeoxycytidine and trichostatin A prostate cells showed increased estrogen receptor β expression and apoptosis. It was found that cell death is connected with estrogen receptor β expression elevation [6]. To investigate an influence of histone H3 lysine 9 acetylation and DNA methylation of P16, HMLH1 and MGMT genes in ovary tumor cells, treatment with trichostatin A and 5-azadeoxycytidine was performed. This experiment showed that DNA demethylation has a better effect on gene reactivation than deacetylation [7]. A problem which arises when using 5-azadeoxycytidine for clinical treatment is its toxicity. It is important to note that 5-azadeoxycytidine and deacetylase inhibitors action is not specific. It was determined using microarray screening of 10814 genes that 5-azacitidine acts on 51 gene, and trichostatin A-on 23 genes [8].

It was shown that 4 weeks after 5-azadeoxycytidine treatment DNA methylation level in most of experimental cells increases again. In cell lines which showed gene reactivation recurrent gene suppression can be induced due to de novo appearing methyltransferases. It was shown too; that many cells after 5-azadeoxycytidine treatment undergo apoptosis because of numerous DNMT1-DNA adducts [9]. 5-azacitidine and 5-azadeoxycytidine have side effects. Patients treated with these chemicals had sickness, vomit, diarrhea and neutropenia. This effect restricts maximal permissible doses of these chemicals and patients therapy duration.

Now a days, causes of such hereditary disorders as Fragile X syndrome are well studied. This causes lie in DNA methylation and histone deacetylation mechanisms. At the same time, generation of therapy methods for these diseases remains a substantial problem. Laboratory investigations presented data about chemicals, which are able to reactivate genes. However,

these agents cannot be put into clinical practice for a range of reasons: high cytotoxicity or low reactivation capacity. That's why search of new drugs which allow to reactivate FMR1 gene is an actual problem. This is especially important considering that Fragile X syndrome is the most often cause of mental retardation, and such patients have difficulties in socialization owing to memory and learning problems and hyperactivity.

Funding Support

The work is supported by fundamental scientific research (FASO Russia) on the project 0310-2016-0004.

References

1. Baylin SB (2005) DNA methylation and gene silencing in cancer . Nat Clin Pract Oncol (2): 4-11.
2. Saldarriaga W, Tassone F, Gonzalez-Teshima LY, Forero-Forero JV, Ayala-Zapata S, et al. (2014) Fragile X syndrome. Colombmedica 45(4): 190-198.
3. Chiurazzi P, Pomponi MG, Pietrobono R, Bakker CE, Neri G, et al. (1999) Synergistic effect of histone hyperacetylation and DNA demethylation in the reactivation of the FMR1 gene. Hum Mol Genet 8(12): 2317-2323.
4. Meng CF, Dai DQ, Guo KJ (2008) Effects of 5-Aza-2'-deoxycytidine and trichostatin A on DNA methylation and expression of hMLH1 in ovarian cancer cell line COC1/DDP. Ai Zheng 27(12): 1251-1255.
5. Walton TJ, Li G, Seth R, McArdle SE, Bishop MC, et al. (2008) DNA demethylation and histone deacetylation inhibition co-operate to re-express estrogen receptor beta and induce apoptosis in prostate cancer cell-lines. Prostate 68(2): 210-222.
6. Meng CF, Su B, Li W (2011) DNA demethylation is superior to histone acetylation for reactivating cancer-associated genes in ovarian cancer cells. Mol Med Rep 4(6): 1273-1278.
7. Suzuki H, Gabrielson E, Chen W, Anbazhagan R, van Engeland M, et al. (2002) A genomic screen for genes upregulated by demethylation and histone deacetylase inhibition in human colorectal cancer. Nat Genet 31(2): 141-149.
8. Pietrobono R, Pomponi MG, Tabolacci E, Oostra B, Chiurazzi P, Neri G, et al. (2002) Quantitative analysis of DNA demethylation and transcriptional reactivation of the FMR1 gene in fragile X cells treated with 5-azadeoxycytidine. Nucleic Acids Res 30(14): 3278-3285.
9. Christman JK (2002) 5-Azacytidine and 5-aza-2'-deoxycytidine as inhibitors of DNA methylation: mechanistic studies and their implications for cancer therapy. Oncogene 21(35): 5483-5495.



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/IJCSMB.2017.01.555573](https://doi.org/10.19080/IJCSMB.2017.01.555573)

Your next submission with Juniper Publishers will reach you the below assets

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission
<https://juniperpublishers.com/online-submission.php>