

A Review on Major Causes of Anemia and its Prevention Mechanism



Hirpasa Teresa*

Department of Biology, Debre Tabor University, Ethiopia

Submission: August 08, 2019; Published: August 28, 2019

*Corresponding author: Hirpasa Teresa, Faculty of Natural and Computational Sciences, Department of Biology, Debre Tabor University, P. O. Box 272, Ethiopia

Abstract

Anemia is one of the most public health problems that cause serious health difficulty like impaired mental and psychomotor development, reduced work performance, increased susceptibility to parasitic infections, and growth retardation in children. The factors for anemia causation are low socio-economic status, nutritional deficiencies, helminthic infections and other infectious diseases, illiteracy, and blood disorder. The major causes of anemia are iron deficiency, foliate deficiency, hookworm infection and malaria. Out of two billion people suffering from anemia, about 50% cases are related to iron deficiency. The cost-effective interventions against anemia include treatment of the underlying causes, restoration of the hemoglobin concentration to normal levels, and prevention and treatment of complications. Furthermore, the simplest and best way to prevent anemia is to eat the gluten free food made from Ethiopian red teff (*Eragrostis tef*) which is full in iron and B-complex vitamins.

Keywords: Anemia; Nutritional deficiency; Helminthic infections; Injera; Teff

Introduction

Anemia is one of the most widespread public health problems throughout the world. It is much more common in developing countries where people suffer from malnutrition and helminthic infections. On the other hand, in developed countries iron deficiency in the food is the major cause of anemia [1]. Anemia is the condition resulted when the body has not enough red blood cells to transport oxygen from the lung to each cell [2]. This occurs when hemoglobin (Hb) concentration level is lower than 11g/dl for pregnant women and children six month to five years of age, 12g/dl for children 6-14 age and 13g/dl for men [1]. Anemia causes serious health problems like impaired mental and psychomotor development, reduced work performance, increased susceptibility to parasitic infections (decreased immunity), growth retardation in children and in severe cases increased rate of child and maternal mortality [3]. According to WHO report two billion people are suffering from anemia in the world of which a major proportion is from South Asia. Globally, almost half (47.4%) of all preschool aged children and 41.8% pregnant women suffer from anemia [1]. Now a day, there is not enough compiled information about the major causes of anemia, and its simplest prevention mechanism. Therefore, the aim of this paper is to review the major causes of anemia and to suggest the simplest way of anemia prevention mechanism.

Major Causes of Anemia

Anemia is caused by different factors including low socio-economic status, nutritional deficiencies, helminthes infections and other infectious diseases, illiteracy, and blood disorder such as sickle cell anemia [4,5]. The major causes in the societies are iron deficiency, hookworm infection, foliate deficiency, and malaria [6]. In the current review the main causes of anemia: nutritional deficiency and parasitic infections are discussed.

Nutritional deficiency

Nutritional deficiency occurs when an individual's daily nutrient intake consistently falls below the recommended requirement. About 1,200 million adolescents suffer from poor nutrition that affects their development and growth [1]. Nutritional deficiencies can lead to anemia, which is the most prevalent health problem in both developed and developing countries. Anemia caused by nutritional deficiency includes iron deficiency anemia, vitamin B12 deficiency anemia and foliate deficiency anemia [7].

Iron deficiency: Iron deficiency is the most widespread condition that mainly causes anemia [1]. WHO estimated that about 50% of anemia is related to iron deficiency [8] It causes reduction in hemoglobin content of red blood cells, the iron containing pigment that carries oxygen from the lungs to the cells [9]. Iron deficiency anemia develops when the amount of iron absorbed

from the diet is lower than the mean daily intakes requirements [10]. The mean daily intake is 0.5mg per day for breast feed infants (0-6 months); 9mg/day for 7-12-month olds; 6-8mg/day for young children and 12-16mg/day for lactating women. Low iron absorption by the body can further be the cause for iron deficiency [11]. Iron deficiency is a major problem affecting health, cognitive development, school achievement and work performance of an individual [12]. Infants, children, adolescents who need additional iron requirements for growth and pregnant women that need additional iron to supply the mother's expanding blood volume and to support the needs of the growing fetus are highly at risk of iron deficiency [1]. Women of childbearing age, who have higher iron losses due to menstrual blood loss also develop iron deficiency [10].

Vitamin B deficiency: Vitamin B-complexes can cause anemia if there is deficiency in the food one eats. Vitamin B₆ deficiency has been associated with anemia because it is important in making hemoglobin that result in increase of oxygen transportation. A form of anemia which results from Vitamin B₆ deficiency is similar to iron deficiency anemia [13]. Riboflavin (Vitamin B₂) deficiency can also cause anemia. It alters iron metabolism by impairing iron absorption, increasing intestinal loss of iron, and/or impairing iron utilization for the synthesis of hemoglobin. In humans, correction of riboflavin deficiency has been known to increase circulating hemoglobin levels thereby improving iron deficiency anemia [1]. Deficiency of vitamin B₁₂ most commonly results in megaloblastic anemia which causes neuro-developmental effects and abnormal movements in children due to its role in foliate metabolism. Megaloblastic anemia is a type of anemia characterized by typical morphological changes in red blood cell precursors which are larger than the cells of same stage and maturation [7]. Most common cause of B₁₂ deficiency in young children has been particularly related to maternal deficiency which leads to decreased stores at birth [13]. Infants who are exclusively breastfed for prolonged period (3-5 years) tend to develop B₁₂ deficiency because of the low content of B₁₂ in mother's milk [14]. In developed countries this problem has been reported from infants born to vegetarian mothers as B₁₂ is not present in plants [15]. Anemia can also be resulted from impaired absorption of Vitamin B₁₂ [1].

Folate deficiency: Margo et al. [16] described foliate deficiency as a frequent complication of protein-energy malnutrition. Like other nutritional deficiencies, foliate deficiency may also lead to anemia and it occurs when absorbed foliate does not meet requirements level over time [17]. Since it is required for the normal production of red blood cells in bone marrow, foliate deficiency results in anemia by bringing red blood cells deformation and production reduction [18]. During the period of pregnancy and lactation, the problem can be encountered as a result of high demand for foliate. Folate stored in tissue may be depleted in up to one third of pregnant women worldwide [19].

Vitamin A deficiency: Vitamin A deficiency contributes to anemia by affecting hemoglobin levels as it is involved in iron metabolism and red blood cell production [7]. It brings risk of anemia

by influencing tissue storage and release of iron into circulation and having a direct regulatory effect on red blood cell production and development [1].

Parasitic infections

Intestinal parasitic infections, which are more prevalent among poor developing countries of the tropics affect over one quarter of the world's population at any one time [20]. Intestinal helminthes infections bring serious public health problems including causing iron deficiency anemia. Chronic blood loss due to hookworm, schistosomes, *Trichuris trichiura* and iron absorption blockage due to *Ascaris lumbricoides* are a significant cause of anemia [12].

Hookworms: Hookworm is a blood feeding intestinal nematode currently infecting over 740 million people throughout the tropics [21]. The adult hookworm attaches to the intestinal mucosa where it chews and digests tissue within its buccal capsule leading to loss of red blood cells. In other case, up on attachment to the host intestine, hookworms produce numerous proteases which have the ability to degrade hemoglobin for its feeding and survival [22]. The major clinical features of chronic hookworm infection are anemia and malnutrition which resulted from blood sucked by the parasite [22]. In chronic intestinal blood loss, iron deficiency anemia develops when blood loss exceeds the intake and reserves of iron and protein in the host. Depending on the status of the iron in the host, a hookworm burden of 40-160 worms is related with hemoglobin levels below 110g/l [23]. Iron deficiency anemia induced by hookworms depends on the species. The two hookworm species that commonly infect human are *Ancylostoma duodenale* and *Necator americanus*. Infection with *Ancylostoma duodenale* causes greater blood loss up to 0.2ml/day/individual whereas infection with *Necator americanus* brings a loss of 0.03ml/day/individual. On the other hand, the depletion of iron stores in the host has a direct correlation with the intensity of hook worm infection [24]. Women of the reproductive age are considerably affected by hookworm infection with adverse effects on the outcome of pregnancy [23]. Crompton (2000) reported that 44 million pregnant women are infected with hookworm worldwide with 7.5 million women in sub-Saharan African alone. According to WHO report severe iron deficiency anemia during pregnancy has resulted increased maternal mortality, impaired lactation, prematurity, and low birth weight [25].

Schistosomes: Schistosomes infection is one of man kind's oldest parasitic infections currently affecting more than 200 million people worldwide [26]. The disease is affecting many people in most African countries. It also presents in limited areas of South America, the Caribbean, the Middle East, and Asia [27]. The species which cause anemia are *Schistosoma mansoni* and *Schistosoma japonicum* (intestinal parasite), and *Schistosoma haematobium* (urinary schistosomiasis). Severe anemia is more firmly resulted from *Schistosoma haematobium* although all three forms can cause it [28]. As reported by Ezeamama et al. [29], blood loss in the stool as emerging eggs rapture intestinal lining, adult in-

gestion of host red blood cells, sequestration of red blood cells in an enlarged spleen, and destruction of red blood cells by autoimmune, results in anemia.

***Trichuris trichiura*:** *Trichuris trichiura* which is endemic in tropical and sub-tropical countries is affecting many people especially in developing countries. *T. trichiura* is most prevalent in warm and humid climate and rare or non-existent in arid, very hot or very cold regions. High intensity of *Trichuris* infection which affects iron status through blood loss in the stool has been reported to cause iron deficiency anemia [30]. The adult *T. trichiura* invade the mucosa and produce minor inflammatory changes. Inflammation at the site of attachment caused both from destruction of the normal colonic architecture as well as post inflammation comprised of macrophage and cytokines which leads to mucoid diarrhea and occasional bleeding [31].

***Plasmodium parasites*:** Malaria which is caused by *Plasmodium parasites* is a devastating disease that affects approximately half of the world population. WHO report indicated that 247 million cases of malaria of which most deaths occur among children living in Sub-Saharan Africa [32]. This devastating disease is thought to be the primary cause of severe anemia (Hb<7g/dl) [25]. Even though the mechanisms by which *plasmodium parasites* cause anemia are quite complex, in its simple terms, the parasites cause anemia both by rupturing red blood cells and by suppressing the production of new red blood cells. It also stimulates the activity of macrophages in the spleen which then destroy both parasitized and unparasitized red blood cells [33]. When red blood cells fall down, hemoglobin, the oxygen carrying protein in red blood cells, will also fall to the extent that cell cannot have the ability to carry oxygen. If this is severe enough, it can bring a significant risk of death by profound hypoxia due to the blockage of capillaries by infected red blood cells, and congestive heart failure, or rarely cerebral malaria. Young children are mostly at risk of malaria because they cannot develop protective immunity against the severe forms of the disease which results in anemia [32]. Recently, researchers discovered that a protein produced by immune cells during malaria infection can also trigger severe anemia [34]. The severity of anemia caused by *plasmodium parasites* depends upon many factors including age, nutritional status, host's and parasites' genotype and immunological history. Children under age five do not produce immunity to resist parasitic damaging when they have been repeatedly infected with *plasmodium parasites*. For this reason, young children account for vast majority of deaths due to anemia caused by plasmodium [35]. Factors such as nutrients malnutrition, pregnancy and HIV infection weaken the immune system to increase incidence of plasmodium infection [36]. Individuals with certain genotypes, for example, with sickle-cell hemoglobin allele, have been known to be protected against infection by *plasmodium parasites*. An outcome of the infection also depends upon the genotypes of *plasmodium parasites*. This is obviously illustrated by the existence of localized varieties of *plasmodium falciparum* that show an inherited resistance to anti-malarial drugs [35]. As a result, many children with severe ane-

mia die and others have a depressed reticulocyte response and a significant number of children had abnormalities of erythropoietic cells in the bone marrow [37]. Severe anemia also affects African pregnant woman in malaria endemic areas [38]. As Guyatt, et al. [39] estimated, approximately 400,000 pregnant women develop moderate or severe anemia annually in sub-Saharan Africa as a result of malaria infection.

Anemia Prevention and Control

Since anemia brings serious public health problems its cost-effective prevention and control is in need worldwide. Many cost-effective interventions against anemia that are well documented include treatment of the underlying causes, restoration of the hemoglobin concentration to normal levels, and prevention and treatment of complications [40]. A number of strategies are used to deliver additional iron to humans, but food fortification and dietary diversification with iron has the greatest potential to prevent iron deficiency anemia [41]. On the other hand, in the areas where vitamin A deficiency is found, WHO recommends routine vitamin A supplementation during pregnancy or at any time during lactation [42]. In case of malaria prevention WHO has recommended a combination of integrated vector management, indoor spraying of insecticide and larval control [8]. In areas endemic with parasitic infections, the International Nutritional Anemia Consultative Group (INACG), WHO, and United Nations Children's Fund (UNICEF) recommended certain control measures such as controlling hookworm and other helminthic infections including sanitary disposal of feces and educational campaigns on proper use of latrines All forms of schistosomiasis including intestinal and urinary types can be prevented by eliminating water dwelling snails, which serve as natural reservoirs of schistosomes [42,43]. Since anemia is mainly a condition with iron and vitamin B deficiencies, the other simplest and best strategy to prevent it is to eat the gluten-free food called Injera made from Ethiopian teff (*Eragrostis tef*). Injera is a major Ethiopian food staple and provides approximately two-thirds of the diet in Ethiopia. Teff is also eaten as porridge and a good thickener for soups, stews, gravies, and puddings and can also be used in stir-fry dishes, and casserole dishes. Furthermore, teff flour is used for making traditional alcoholic drinks like tella (local beer), katikalla (local spirit), kitta (sweet dry unleavened bread), and muk (gruel) [44] Figure 1. Teff is full in iron and B-complex vitamins although many gluten free foods are not enriched with these nutrients. Researchers identified that the iron content of red teff can be up to 150mg/100g and one cup of raw grain teff can provide as high as 82% of the recommended daily intake of iron. As a result, in people living in areas of Ethiopia where consumption of red teff is most prevalent, hemoglobin levels were found to be higher with a decreased risk of anemia [45-47]. In addition to iron, teff contains B-complex vitamins that are used in the formation of red blood cells and facilitate nutrients' metabolism. One cup of uncooked teff contains 47% of the recommended daily intake for vitamin B₆, 50% for vitamin B₁, 31% for vitamin B₂, 32% for vitamin B₃ and 18% for vitamin B₅. Furthermore, teff is full in vitamin C that is used in iron absorp-

tion but contains small amount of vitamin A [48]. In another study, Gamboa [49] identified different vitamins found in teff as stated on Table 1. Currently, there has been a growing global interest in the use of teff as a food. This interest is mainly attributed to teff

being gluten-free and having relatively high nutrient content and thus a candidate ingredient for food products destined for people with celiac disease [50,51].



Figure 1: Injera with Mesob (handling material), captured by the author immediately after preparation at home.

Table 1: Amount of vitamins in the flour of teff grain.

Vitamin	mg/100g	Advised daily amount for a 75kg human (mg)
Thiamin (B ₁)	0,51	1,0
Riboflavin (B ₂)	< 0,1	1,5
Niacin (B ₃)	0,80	16
Pyridoxin (B ₆)	<0,1	3
Ascorbic acid (C)	0,25	70
Folic acid (M)	<0,02	0,4

Conclusion

Anemia, the most widespread public health problems, causes impaired mental and psychomotor development, reduced work performance, and growth retardation in children. It is mainly caused by nutritional deficiencies and parasitic infections. Nutritional deficiency including iron deficiency anemia, vitamin B₁₂ deficiency anemia and foliate deficiency anemia are the major causes of anemia. Blood loss due to hookworm, schistosomes, *Trichuris trichiura* and iron absorption blockage due to *Ascaris lumbricoides* are a significant cause of anemia. Ethiopian teff (*Eragrostis tef*), which has recently got international acceptance, is recommended as the best remedy to prevent anemia.

Acknowledgement

I would like to acknowledge all the previous Investigators for doing different studies that I have used for this review paper.

References

- Badham J, Zimmermann MB, Kraemer K (2007) The Guidebook Nutritional Anemia. Sight and life press, Basel, Switzerland. pp. 11-42.
- LeHesran JY, Akiana J, Ndiaye HM, Dia M, Senghor P (2004) Severe malaria attack is associated with high prevalence of *Ascaris lumbricoides* infection among children in rural Senegal. *Trans R Soc Trop Med Hyg* 98(7): 397-399.
- Georgieff MK (2011) Long-term brain and behavioural consequences of early iron deficiency. *Nutrition Reviews* 69 (S1): S43-S48.
- World Bank (1980) Poverty and development of human resources: regional perspectives. Washington, DC: World Bank staff working paper no. 406.
- Menendez C, Fleming AF, Alonso PL (2000) Malaria-related anemia. *Parasitol Today* 16(11): 469-476.
- Fleming AF, Weatherall DJ, Ledingham JGG, Warrell DA (1987) Anaemia as a world health problem. *Oxford textbook of medicine* 2nd (edn). Oxford: Oxford University Press. pp. 72.
- Hahn JS, Nam Dk, Lee EJ, Lee SJ, Koo YW, et al. (1988) A Case of Vitamin B₁₂ Deficiency *Megaloblastic Anemia* Following Total Gastrectomy. *Yonsei Medical Journal* 29(3): 270-277.
- World Health Organisation (2011) WHO Vitamin and Mineral Nutrition/Anaemia.
- Ramzan M, Ali I, Salam A (2009) Iron Deficiency Anemia in School Children of Dera Ismail Khan, Pakistan. *Pakistan Journal of Nutrition* 8 (3): 259-263.
- Cook JD, Finch CA (1979) Assessing iron status of a population. *Am J Clin Nutr* 32(10): 2115-2119.
- Bothwell TH, Baynes RD, Mac Farlane BJ, Mac Phail AP (1989) Nutritional iron requirements and food iron absorption. *J Inter Med* 226(5): 357-365.
- Osório MM, Lira PI, Batista Filho M, Ashworth A (2001) Prevalence of anemia in children 6-59 months old in the state of Pernambuco, Brazil. *Pan Am J Public Health* 10(2): 101-106.
- Jain V (1999) Clinico hematological study of *megaloblastic anemia*. MD Thesis, Delhi University. pp. 50-52.

14. Casterline JE, Allen LH, Ruel MT (1997) Vitamin B₁₂ deficiency is very prevalent in Lactating Guatemalan women and their infants at three months postpartum. *J Nutr* 127(10):1966-1972.
15. Smolka V, Bekarek V, Hlilkova (2001) Metabolic complications and neurologic manifestations of vitamin B₁₂ deficiency in children of vegetarian mothers. *Cas Lek Cesk*, 140(23): 732-735.
16. Margo G, Baroni Y, Wells G, Green R, Metz J (1978) Protein energy malnutrition and nutritional anemia in preschool children in rural Zwazulu. *S Afr Med J* 53(1): 21-26.
17. Fleming AF, Werblinska B (1982) Anemia in childhood in the Guinea Savanna of Nigeria. *Ann Trop Paediatr* 2(4): 161-173.
18. Fleming AF (1989) The etiology of severe anemia in pregnancy in Zambia. *Ann Trop Med Parasitol* 83(1): 37-49.
19. FAO/WHO (1970) Requirements of ascorbic acid, vitamin D, vitamin B12, foliate and iron. World Health Organization, Geneva, Switzerland. WHO Technical Report Series 452.
20. Chan MS (1997) The global burden of intestinal nematode infections. Fifty years on *Parasitol Today* 13(11): 438-443.
21. De Silva NR, Brooker S, Hotez PJ (2003) Soil-transmitted helminth infections: updating the global picture. *Trends Parasitol* 19(12): 547-551.
22. Cappello M, Harrison LM, Bungiro RD (2003) Molecular pathogenesis of hookworm anemia: prospects for a disease-based vaccine. *J Parasitol* 89: 158-164.
23. Bundy DA, Chan MS, Savioli L (1995) Hookworm infection in pregnancy. *Trans R Soc Trop Med Hyg* 89(5): 521-522.
24. Oisen A, Magnussen P, Ouma JH, Andreassen J, Friis H (1998) The contribution of hookworm and other parasitic infections to hemoglobin and iron status among children and adults in western Kenya. *Trans R Soc Trop Med Hyg* 92(6): 643-649.
25. World Health Organization (2002) Prevention and control of schistosomiasis and soil-transmitted helminthiasis. WHO technical report series 912. Geneva, Switzerland.
26. Tan SY, Ahana A (2007) *Medicine in Stamps*. Singapore Med J 48: 184.
27. Charles D, Ericsson, Steffen R (2002) Schistosomiasis and International Travel. *Travel Medicine*. CID: 35.
28. World Health Organization (1991) Basic laboratory methods in medical parasitology. Geneva, Switzerland.
29. Ezeamama AE, Mc Garvey ST, Acosta LP, Ziorler S, Manalo DL, et al. (2008) The synergistic effect of concomitant Schistosomiasis, Hookworm, and *Trichuris infections* on Children's Anemia Burden. *Plos Negl Trop Dis* 2(6): 1-9.
30. Ramdath DD, Simeon DT, Wong MS, Grantham Mc Gregor SM (1995) Iron status of schoolchildren with varying intensities of *Trichuris trichiura* infection. *Parasitology* 110(pt 3): 347-351.
31. Mac Donald TT, Spencer J, Murch SH (1994) Mucosal macrophages and cytokine production in the colon of children with *Trichuris trichiura* dysentery. *Trans R Soc Trop Med Hyg* 88(3): 265-268.
32. World Health Organization (2010) General Assembly.
33. Abdalla S, Weatherall DJ, Wickramasinghe SN, Hughes M (1980) The anemia of *Plasmodium falciparum* malaria. *Br J Haematol* 46(2): 171-183.
34. Kai OK, Roberts DJ (2008) The pathophysiology of malarial anemia: where have all the red cells gone? *BMC Medicine* 6: 24.
35. Mendis KN, Carter R (1995) Clinical Disease and pathogenesis in Malaria. *Parasitology today* 11(5): 2-16.
36. Shulman CE, Dorman EK (2003) Reducing childhood mortality in poor countries: importance and prevention of malaria in pregnancy. *Transitions of the Royal Society of Tropical Medicine and Hygiene* 97: 30-35.
37. Allen LH, Rosado JL, Casterline JE (1995) Vitamin B₁₂ deficiency and malabsorption are highly prevalent in Mexican communities. *Amer J Clin Nutr* 65(2): 1013-1019.
38. Marsh K (1992) Malaria, a neglected disease. *Parasitology Suppl* 104: S53-S69.
39. Guyatt, HL, Snow RW (2001) The epidemiology and burden of *Plasmodium falciparum*-related anemia among pregnant women in sub-Saharan Africa. *Am J Trop Med Hyg* 64(1-2): 36-44.
40. Osungbade KO, Oladunjoye AO (2012) Anaemia in Developing Countries: Burden and Prospects of Prevention and Control.
41. World Health Organization (2005) WHO Global Database on Iron Deficiency and Anaemia, Micronutrient Deficiency Information System. Geneva, Switzerland.
42. ACC/SCN (2001) What Works? A Review of the Efficacy and Effectiveness of Nutrition Interventions. Geneva in collaboration with the Asian Development Bank, Manila.
43. World Health Organization (1996) Report of the WHO informal consultation on hookworm infection and anaemia in girls and women. Geneva, Switzerland.
44. Bultosa G (2002) Physico-chemical Characterization of Grain Tef [*Eragrostis tef* (Zucc.) Trotter] *Starch*. 54: 461-468.
45. <http://ethnomed.org/cultures/ethiop/teff.html>
46. Demeke M, Marcantonio F (2013) Analysis of incentives and disincentives for teff in Ethiopia. Technical notes series, MAFAP, FAO, Rome.
47. Kaleab Baye (2014) Teff: nutrient composition and health benefit. Ethiopian Development Research Institute (EDRI). Addis ababa.
48. <http://www.livestrong.com/article/543343-nutritional-value-of-teff-grain>
49. Gamboa PA (2008) TEFF Survey on the nutritional and health aspects of teff (*Eragrostis Tef*). Instituto Tecnológico de Costa Rica, Sede Central, Apdo. 159-7050 Cartago, Costa Rica.
50. Hager, Anna Sophie, Elke K Arendt (2013) Influence of Hydroxypropyl methylcellulose (Hpmc), Xanthan Gum and Their Combination on Loaf Specific Volume, Crumb Hardness and Crumb Grain Characteristics of Gluten-Free Breads Based on Rice, Maize, Teff and Buckwheat. *Food Hydrocolloids*. 32 (1): 195-203.
51. García Manzanares Á, Lucendo AJ (2011) Nutritional and Dietary Aspects of Celiac Disease. *Nutrition in Clinical Practice* 26 (2): 163-173.



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/IJCSMB.2019.06.555686](https://doi.org/10.19080/IJCSMB.2019.06.555686)

**Your next submission with Juniper Publishers
will reach you the below assets**

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission
<https://juniperpublishers.com/online-submission.php>