

Mini Review

Volume - 5 Issue - 5 October 2017  
DOI: 10.19080/JFSCI.2017.05.555678

J Forensic Sci & Criminal Inves

Copyright © All rights are reserved by Manoj Balachandran

# Medicinal Poisons



ManojBalachandran<sup>1</sup>, Vin Ravi Vaswani<sup>2</sup> and Hashim A<sup>3</sup>

<sup>1</sup>Post Graduate, Department of Forensic Medicine & Toxicology, Yenepoya Medical College, India

<sup>2</sup>Prof and Head, Department of Forensic Medicine & Toxicology, Yenepoya Medical College, India

<sup>3</sup>Lecturer/Toxicologist, Department of Forensic Medicine & Toxicology, Yenepoya Medical College, India

Submission: October 03, 2017; Published: October 16, 2017

\*Corresponding author: Manoj Balachandran, Post Graduate, Department of Forensic Medicine & Toxicology, Yenepoya Medical College, Mangalore, India, Email: bmanoj7@yahoo.co.in

### Abstract

Almost all pharmaceutical substance is potentially poisonous if taken in sufficient dosages or for a long period. However, toxic and fatal consequences are most commonly accidental or suicidal, although homicide should always be borne in mind, even though it is rare [1].

### Introduction

There is great variation in the drugs prescribed, used and most importantly available to the public in different countries. In many countries there are strict controls on the supply of drugs but many potentially lethal compounds are still available to the public without prescription in the shops. Where drugs are easily available they are by far the most common method of suicide and suicidal gestures. The subject has expanded so greatly that it properly belongs in the field of clinical toxicology rather than legal medicine [1].

### From Past to Present

The history of poison stretches from before 4500 B.C. to the present day. Poisons have been used for many purposes across the span of human existence, most commonly as weapons, anti-venoms and medicines. Poison has allowed much progress in branches, toxicology, and technology, among other sciences



[1]. Some medicinal plants like Corymbia citriodora were used against malaria due to their antiplasmodial activity [2]. Datura Stramonium is a widely used poisonous plant with great medicinal and economic value [3].

### Common Medicinal Poisons

- a) Analgesics,
- b) Antidepressant and sedative drugs,
- c) Barbiturates,
- d) Seda-tive hypnotic non-barbiturate,
- e) Analgesic- antipyretic,
- f) Antimanic agents
- g) Insulin [1] (Table 1).

Table 1: Tests for Detecting Medicinal Poisons.

Serial No.	Name of drug	Test performed	Results
1	Paracetamol 	Qualitative test A trong, royal blue colour Applicable to urine, stom- developing immediately ach contents and scene indicates the presence of residues. <b>Method</b> 1. Add 0.5 ml of hydro-chloric acid to 0.5 ml of sample, boil for 10 minutes and cool. 2. Add 1 ml of o-cresol solution to 0.2 ml of the hydrolysate. 3. Add 2 ml of ammonium hydroxide solution and mix for 5 seconds.	A strong royal blue colour developed immediately indicates the presence of Paracetamol

2	<p>Amitriptyline</p> 	<p>There is no simple qualitative test for amitriptyline, but this compound and other tricyclic antidepressants can be easily detected and identified by thin-layer chromatography of a basic solvent extract of urine, stomach contents or scene residues</p>	<p>Mandelin's reagent (basic extract) gives colours ranging from blue and green to orange and red with a variety of basic compounds</p>
3	<p>Barbiturates</p> 	<p>A qualitative analysis is best performed by thin-layer chromatography of a solvent extract of urine, stomach contents or scene residues.</p>	<p>Mercurous nitrate reagent (acidic extract) gives white spots with a grey centre on a darker background</p>
4	<p>Chloral hydrate</p> 	<p>Applicable to urine. Fujiwara test</p>	<p>An intense red/purple colour in the upper, pyridine layer indicates the presence of chloral compounds.</p>
5	<p>Phenacetin</p> 	<p>Applicable to urine, o- Cresol/ammonia test</p>	<p>A strong, royal blue colour developing immediately indicates the presence of paracetamol.</p>
6	<p>Lithium</p> 	<p>Applicable to scene residues. Method 1. Dip the end of the platinum wire in the concentrated acid. 2. Dip the moistened end of the wire into the test material. 3. Place the material in the hot part of the flame of a microburner.</p>	<p>A crimson red flame denotes the presence of lithium salts</p>

**Doctrine of Double Effect**

The doctrine of double effect says that if doing something morally good has a morally bad side-effect, it is ethically correct to do it provided the bad side-effect wasn't intended. This is true even if the bad effect is unavoidable. This principle has been seriously argued in ethics. According to the traditional Catholic conviction if there are exception less norms prohibiting inflicting some kinds of harms on people is correct, then double effect is

justified and necessary [4]. The principle is used to justify the case where a doctor gives drugs to a patient to relieve distressing symptoms even though he knows doing this may shorten the patient's life. This is because the doctor is not aiming directly at killing the patient - the bad result of the patient's death is a side-effect of the good result of reducing the patient's pain. Many doctors use this doctrine to justify the use of high doses of drugs such as morphine for the purpose of relieving suffering

in terminally-ill patients even though they know the drugs are likely to cause the patient to die sooner [1]. The value of doctrine of double effect is of special importance in euthanasia and abortion [5].

### Conclusion

All pharmaceutical substances are to be taken in the correct dosage for the correct period of time. In case of over-dosage, it is important to identify the drug ingested so that resuscitative measures can be initiated rapidly. Doctrine of double effect where the doctor gives drug to a patient to relieve distressing symptoms even though he knows doing this may shorten the patient's life should be used with caution.

### Acknowledgement

I would like to thank the teaching, non teaching staffs and PGs of Department of Forensic Medicine, Yenepoya Medical

College, Mangalore for their help and guidance in making this poster.

### References

1. Shepherd R (2003) Simpson's Forensic Medicine Twelfth Edition. Arnold Publications 25:170-174.
2. Singh N, Kaushik NK, Mohanakrishnan D, Tiwari SK, Sahal D (2015) Antiplasmodial activity of medicinal plants from Chhotanagpur plateau, Jharkhand, India- J Ethnopharmacol 165:152-62.
3. Yang Y, Dang Y, Li Q, Lu J, Wang Y (2014) Complete chloroplast genome sequence of poisonous and medicinal plant *Datura Stramonium* organisations and implications for genetic engineering. PLoS One 9(11): e110656.
4. Shaw AB (2002) Two challenges to the double effect doctrine: euthanasia and abortion. J Med Ethics 28: 102-104.
5. Boyle Joseph (1991) Who is entitled to double effect. The Journal of Medicine and Philosophy 16: 475-494.



This work is licensed under Creative Commons Attribution 4.0 License  
DOI: 10.19080/JFSCI.2017.04.555678

#### Your next submission with Juniper Publishers

will reach you the below assets

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats  
( Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission

<https://juniperpublishers.com/online-submission.php>