



Case Report

Volume 7 Issue 3 - October 2017  
DOI: 10.19080/JGWH.2017.07.555711

J Gynecol Women's Health

Copyright © All rights are reserved by Angel Danchev Yordanov

# Very Large, Rapidly Growing Myoma during Second Trimester of Pregnancy - Outcome



Angel Danchev Yordanov<sup>1\*</sup>, Krasimira Zhivkova Zhelyaskova<sup>2</sup>, Ilko Ivanov Iliev<sup>1</sup>, Polina Petkova Vasileva<sup>3</sup>, Nikola Kalinov Popovski<sup>3</sup> and Momchil Denev Ivanov<sup>4</sup>

<sup>1</sup>Clinic of Gynecologic Oncology, University Hospital "Dr. Georgi Stranski"-Pleven, Bulgaria

<sup>2</sup>Medical Center "Petinka Tzvetkova"-Varna, Bulgaria

<sup>3</sup>Department of Obstetrics and Gynecology, Medical University, Bulgaria

<sup>4</sup>MHAT "Saint Paraskeva"-Pleven, Bulgaria

Submission: October 18, 2017; Published: October 26, 2017

\*Corresponding author: Angel Danchev Yordanov, Clinic of Gynecologic Oncology, University Hospital "Dr. G. St. ranski"-Pleven, Bulgaria, Email: [angel.jordanov@gmail.com](mailto:angel.jordanov@gmail.com)

## Abstract

**Introduction:** The leiomyoma is the most common gynaecological tumor. Uterine myomas during pregnancy occur in 2 to 10% of all cases. One tenth of those may lead to complications of the pregnancy. According to most ultrasounds studies myomas remain the same size or become smaller during pregnancy.

**Case:** This case report is about 34 year old woman, nullipara, in 14th week of pregnancy. Through ultrasound observation was found subserous pediculated myoma node with diameter of 20cm. When the pregnancy was found, the myoma was not. A median laparotomy with myomectomy was performed without any complications. On the 38th week of pregnancy a healthy, 3400g, full-term baby was delivered by cesarean section.

**Conclusion:** Though rarely, rapidly growing and very large myomas can be an indication for miomectomy during the second trimester of pregnancy. Laparotomic myomectomy should be preferred over laparoscopic myomectomy in cases of very large myomas, due to prolonged operative time, inapplicability of in-bag morcellation and related risks of dissemination of a leiomyoma or leiomyosarcoma.

**Keywords:** Very large myoma; Pregnancy; Myomectomy; Rapidly growing myoma

## Introduction

The leiomyoma is the most common benign gynaecological tumour [1]. Uterine myomas during pregnancy occur in 2 to 10% of all cases. One tenth of those may lead to complications of the pregnancy. According to most ultrasounds studies myomas remain the same size or become smaller during pregnancy [2-4]. Estrogen and progesterone were thought to play a major role in the development and growth of myomas, but recently this theory has been questioned.

Usually myomectomy is avoided during pregnancy, but in some situations it is indicated, despite the risk of hemorrhagic complications that may require hysterectomy due to the increased vascularity [5-7]. Laparotomic myomectomy during pregnancy has been reported as a safe approach more than a hundred years ago [8,9]. The most common indications for myomectomy during pregnancy are acute pelvic pain that is not responsive to medical therapy, signs of red infarction or torsion

of pedunculated myomas, and abdominal discomfort due to a large or rapidly growing myomas [7,10].

## Clinical Case

We present a 34 year old nulliparous, Caucasian woman at 14th week of gestation. The patient was admitted in our clinic with a 20cm myoma. A few months ago, before the conception, through ultrasound observation a normal gynecological status was found. At 5<sup>th</sup> gestational week a normal pregnancy was detected, without any signs of myoma. At 7<sup>th</sup> week of pregnancy a 3.5cm. subserous myoma was found. At 12<sup>th</sup> week of pregnancy the ultrasound observation showed a subserous pediculated myoma on the left edge of the uterine fundus with dimensions 15/10.9cm. The analysis of the ultrasound observations showed unexpected progressive increasing of the myoma size. Because of the rapid growth and the big size a median laparotomy with myomectomy was performed. A 23cm pedunculated subserosal

myoma with an implantation base of 4cm was found (Figure 1 & 2). The operation and the postoperative period take place without any complications. On the 38th week of pregnancy a healthy, 3400g, full-term, baby boy was delivered by cesarean section. Postpartum and the puerperium period had a regular course.



Figure 1: Intraoperative findings.



Figure 2: Myoma.

## Discussion

Different authors classify myomas by size in different ways. Most agree that myoma is large when it is over 5cm [11-14]. The Society of Obstetricians and Gynaecologists of Canada, classify myomas over 10cm. like very large [15].

Most ultrasound studies have shown that most of the fibroids during pregnancy (60%-78%) do not show significant changes in their volume. Out of the fibroids that did increase their volume (22% to 32%), the growth was limited in the first trimester [16,17]. There is a prospective study, according to which, if a myoma increases its size during pregnancy, that increase is 12±6% of the initial size, up to a maximum of 25% [16]. A variety

of factors, such as genetic and epigenetic mechanism, growth factors, cytokines and disorganization of extracellular matrix components have shown interaction with their pathogenesis [18].

How leiomyomas affect pregnancy is based on their number, size and localization. A significant part of myomas remain asymptomatic and do not require treatment, but 10% lead to complications such as: fetal growth restriction, pelvic pain, placental abruption, placenta previa / accrete, fetal deformities, dystocia, postpartum hemorrhagia [19]. Acceptable indications for myomectomy during pregnancy include intractable pain from a degenerating myoma especially if it is subserosal or pedunculated, a large or rapidly growing myoma, or any large myoma (> 5cm) located in the lower uterine segment [12]. The outcome, both obstetric and neonatal, of women, who underwent myomectomy in pregnancy is comparable to the outcome of those, who were managed conservatively [20,21]. According to few studies, antepartum myomectomy can be safely performed in the first and second trimester of pregnancy, if indicated [20-27].

There is enough scientific data about the safety of laparotomic myomectomy during pregnancy. Recently the number of case reports and studies about the possibility of performing a safe laparoscopic myomectomy during pregnancy is gradually increasing. It begins to slowly replace open surgery because of certain advantages less blood loss, diminished postoperative pain, fewer overall complications, faster recovery and significant cosmetic. However, there are also disadvantages, like the prolonged operative time because of the need of morcellation, especially time-consuming myomas larger than 10cm [16,28].

Morcellation by itself has additional risks of vascular or visceral trauma, and can also lead to leiomyoma dissemination (leiomyomatosis). The possibility of a leiomyosarcoma can't be ruled out, which would lead to a sudden decrease in survivability and upstaging of the disease [16]. Some of those risks can be brought down to a minimum through the use of in-bag morcellation, but when a large myoma is present and given the limited area, in cases like ours it is inapplicable. Another disadvantage of the laparoscopic myomectomy during pregnancy is the need of additional training, an experienced surgeon and specialized equipment [16].

## Conclusion

Though rarely, rapidly growing and very large myomas can be an indication for myomectomy during the second trimester of pregnancy, because of the limited space for fetus development.

Laparotomic myomectomy should be preferred over laparoscopic myomectomy in cases of very large myomas, due to prolonged operative time, inapplicability of in-bag morcellation and related risks of dissemination of a leiomyoma or leiomyosarcoma.

## Acknowledgement

The publication was funded by project BG05M2OP001-2.009-0031-C01.

## References

- Ryan GL, Syrop CH, Van Voorhis BJ (2005) Role, epidemiology, and natural history of benign uterine mass lesions. *Clin Obstet Gynecol* 48(2): 312-324.
- Hammoud AO, Asaad R, Berman J, Treadwell MC, Blackwell S, et al. (2006) Volume change of uterine myomas during pregnancy: do myomas really grow? *J Minim Invasive Gynecol* 13(5): 386-390.
- Muram D, Gillieson M, Walters JH (1980) Myomas of the uterus in pregnancy: ultrasonographic follow-up. *Am J Obstet Gynecol* 138(1): 16-19.
- Neiger R, Sonek JD, Croom CS, Ventolini G (2006) Pregnancy-related changes in the size of uterine leiomyomas. *J Reprod Med* 51(9): 671-674.
- Yumi H (2008) Guidelines for diagnosis, treatment, and use of laparoscopy for surgical problems during pregnancy. *Surg Endosc* 22: 849-861.
- Lolis DE, Kalantaridou SN, Makrydimas G, Sotiriadis A, Navrozoglou I, et al. (2003) Successful myomectomy during pregnancy. *Hum Reprod* 18(8): 1699-1702.
- Fanfani F, Rossitto C, Fagotti A, Rosati P, Gallotta V, et al. (2010) Laparoscopic myomectomy at 25 weeks of pregnancy: case report. *J Minim Invasive Gynecol* 17(1): 91-93.
- Evans HM (1899) A case of myomectomy for subperitoneal myoma complicating pregnancy. *Br Med J* 2: 1673.
- Doran A (1906) Myomectomy during pregnancy and labour at term in an elderly primipara: with notes on similar cases. *Br Med J* 2(2395): 1446-1447.
- Ardivino M, Ardivino I, Castaldi MA, Monteverde A, Colacurci N, et al. (2011) Laparoscopic myomectomy of a subserous pedunculated fibroid at 14 weeks of pregnancy: a case report. *J Med Case Rep* 5: 545.
- American Association of Gynecologic Laparoscopists (AAGL): Advancing Minimally Invasive Gynecology Worldwide (2012) AAGL Practice Report: Practice Guidelines for the Diagnosis and Management of Submucous Leiomyomas. *J Minim Invasive Gynecol* 19(2): 152-171.
- Joong Lee H, Norwitz ER, Shaw J (2010) Contemporary Management of Fibroids in Pregnancy. *Rev Obstet Gynecol* 3(1): 20-27.
- Marvelos D, Ben-Nagi J, Holand T, Hoo W, Naftalin J, et al. (2010) The natural history of fibroids. *Ultrasound Obstet Gynecol* 35(2): 238-242.
- Kanaoka Y, Hirai K, Ishiko O (2005) Microwave endometrial ablation for menorrhagia caused by large submucous myomas. *J Obstet Gynaecol Res* 31(6): 565-570.
- Vilos G, Allaire C, Laberge Ph, Leyland N (2015) The Management of Uterine Leiomyomas. *J Obstet Gynaecol Can* 37(2): 157-178.
- Aharoni A, Reiter A, Golan D, Paltiely Y, Sharf M (1988) Patterns of growth of uterine leiomyomas during pregnancy. A prospective longitudinal study. *Br J Obstet Gynaecol* 95(5): 510-513.
- Rosati P, Exacoustòs C, Mancuso S (1992) Longitudinal evaluation of uterine myoma growth during pregnancy. A sonographic study. *J Ultrasound Med* 11(10): 511-515.
- Ciavattini A, Di Giuseppe J, Stortoni P, Montik N, Giannubilo SR, et al. (2013) Uterine fibroids: pathogenesis and interactions with endometrium and endomyometrial junction. *Obstet Gynecol Int* 2013: 173184.
- Klatsky PC, Tran ND, Caughey AB, Fujimoto VY (2008) Fibroids and reproductive outcomes: a systematic literature review from conception to delivery. *Am J Obstet Gynecol* 198(4): 357-366.
- Exacoustòs C, Rosati P (1993) Ultrasound diagnosis of uterine myomas and complications in pregnancy. *Obstet Gynecol* 82(1): 97-101.
- Mollica G, Pittini L, Minganti E, Perri G, Pansini F (1996) Elective uterine myomectomy in pregnant women. *Clin Exp Obstet Gynecol* 23(3): 168-172.
- De Carolis S, Fatigante G, Ferrazzani S, Trivellini C, De Santis L, et al. (2001) Uterine myomectomy in pregnant women. *Fetal Diagn Ther* 16(2): 116-119.
- Wittich AC, Salminen ER, Yancey MK, Markenson GR (2000) Myomectomy during early pregnancy. *Mil Med* 165(2): 162-164.
- Glavind K, Palvio DH, Lauritsen JG (1990) Uterine myoma in pregnancy. *Acta Obstet Gynecol Scand* 69(7-8): 617-619.
- Michalas SP, Oreopoulou FV, Papageorgiou JS (1995) Myomectomy during pregnancy and caesarean section. *Hum Reprod* 10: 1869-1870.
- Febo G, Tessarolo M, Leo L, Arduino S, Wierdis T, et al. (1997) Surgical management of leiomyomata in pregnancy. *Clin Exp Obstet Gynecol* 24(2): 76-78.
- Celik C, Acar A, Çiçek N, Gezginc K, Akyürek C (2002) Can myomectomy be performed during pregnancy? *Gynecol Obstet Invest* 53(2): 79-83.
- Saccardi C, Visentin S, Noventa M, Cosmi E, Litta P, et al. (2014) Uncertainties about laparoscopic myomectomy during pregnancy: A lack of evidence or an inherited misconception? A critical literature review starting from a peculiar case. *Minim Invasive Ther Allied Technol* 24(4): 189-194.



This work is licensed under Creative Commons Attribution 4.0 License  
DOI: [10.19080/JGWH.2017.07.555711](https://doi.org/10.19080/JGWH.2017.07.555711)

### Your next submission with Juniper Publishers will reach you the below assets

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats  
( Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission  
<https://juniperpublishers.com/online-submission.php>