

OpinionVolume 1 Issue 1 - May 2019
DOI: 10.19080/JOJWB.2019.01.555555

JOJ Wildl Biodivers

Copyright © All rights are reserved by Pedro Manuel Villa

Inselbergs of Venezuelan Guayana Region: A Natural Laboratory for Plant Ecology Research

**Pedro Manuel Villa***

Program in Botany, Federal University of Viçosa, Brazil

Submission: January 17, 2019; **Published:** May 28, 2019***Corresponding author:** Pedro Villa, Researcher, Program in Botany, Federal University of Viçosa, CEP 36570000 Viçosa, Minas Gerais, Brazil

Opinion

The Guayana shield is one of the most important biogeographic regions of the world, with a complex matrix of ecosystems and different types of vegetation [1-4]. One of the most outstanding ecological units of the region are the granitic outcrops, also known as *inselbergs* (from the German Insel: island, and Berg: mountain), and locally as *lajas*. These rock formations have high plant species diversity, a high number of endemic species with restricted geographical distribution [2-3]. From a biogeographical approach, these geological formations are also considered as continental islands, mainly because of the ecological processes that depend to a large extent on geographic isolation [5]. This geographical isolation confers unique ecological conditions for the formation of different types of plant communities on a small spatial scale. However, since pioneering research on Guayana *inselbergs* vegetation types, e.g., [3-4], knowledge about plant community ecology is still limited as a basis for establishing conservation measures [6].

The Guayana *inselbergs* present different types of plant communities, which are shaped by a complex system of environmental gradients, mainly by the availability of water, topography and the depth of the substrates [5,7-8]. Thus, the mosaic of habitats present in the *inselbergs* of Guayana region, can occur discontinuously separated with different types of vegetation, such as epilithic (on bare rock), casmophytic (between cracks), forest in the depressions, continuous forests in flat sites of the peaks, and in accumulations of large blocks of granite. In addition, one of the most dominant types of vegetation are the patches of vegetation found in the depressions [3-5]. However, it is not yet known about the assembly rules of plant communities, especially the relative contribution of abiotic (i.e., microclimate, topography, water deficit) and biotic (i.e., facilitation) drivers, to test hypotheses in plant ecology.

The geomorphological and microenvironmental conditions of the *inselbergs* confer special ecological conditions where only

some plant species with different types of morphological and ecophysiological adaptations can be developed [5,9-11]. This leads to a greater specialization of the species on a small spatial scale in extreme conditions compared to other surrounding ecosystems, such as savannas and forests [3-4]. For example, many of these species grow on substrates accumulated in depressions of different sizes, and on which patches of vegetation are developed [6], also known as islands of vegetation, on islands of rocky outcrops [7].

Recently some researchers have presented results on the ecology of plant communities in inselberg of Venezuelan Guayana; this has been analyzing the effect of patch size on diversity patterns [6,12]. Thus, these studies carried out in the *Piedra la Tortuga* Natural Monument reveal that there is a considerable number of endemic species in fine-scale vegetation patches, with up to 50% of endemic species per patch of vegetation [6]. This pattern of diversity is exclusive from a biogeographical perspective, because it is not common in other plant communities present around the *inselbergs* of Guayana, as for example in the savannas and forests that have a wide distribution throughout the region. This consequence has been well described, that in the *inselbergs* there is an important representation of endemic species (approximately 144) in relation to the entire Guayana region, and that is why it is considered as an important center of endemism in Venezuela [2-4]. On the other hand, several studies have also reported the high level of endemism existing in rocky outcrops in Australia, Africa and tropical America [5]. Although, there are still limited studies on the species-area relationship and species abundances distribution as basic information to establish new conservation areas [6].

The *inselbergs* of western Guayana are dominated mainly by herbaceous and shrubby growth forms (Figure 1), such as *Tabebuia orinocensis*, *Acanthella sprucei*, *Commiphora leptophloes*, and *Pseudobomax croizatii* the most representative

shrub species in areas visited and studied on an ecological approach [6,12]. On the other hand, herbaceous species such as *Irlbachia alata*, *Utricularia subulata*, and *Sauvagesia ramosissima*, have been reported as ephemeral plants; which appear during the wetter periods when water stress is possibly less marked in these ecological formations [3-4,9-10]. In the case of shrubs, despite not being abundant in different types of plant communities in these *inselbergs*, they could be key species because probably generate appropriate microclimatic conditions for the establishment and survival of other species, for example in patch of vegetation found in the depression (Figure 2). Succulent stems (cactus) are other life forms characteristic of the study area, in relation to their strategies for storing water (eg *Melocactus* spp), also succulent leaves (*Pitcairnia pruinosa*, *Pitcairnia armata*), and geophytic as *Portulaca pygmaea* [4,9,13].



Figure 1: *Inselbergs* of western Guayana, Cuao river, Amazon state, Venezuela.



Figure 2: Patch of vegetation found in the depression, *Piedra la Tortuga* Natural Monument, Amazon state, Venezuela.

Final Remarks

Due to the importance value of the plant diversity present in the *inselbergs* of Guayana, it would be essential to continue research on the ecology of plant communities; especially due to the urgent need to create new conservation areas. For example, one of the few conservation units known in the region is the *Piedra la Tortuga* Natural Monument. Likewise, to establish new

conservation areas with *inselbergs* in the Guayana region, it is necessary to consider that species with restricted distribution tend to be more vulnerable to extinction. Therefore, it is essential to know the minimum viable population of all those threatened species of *inselbergs*. This would allow estimating the minimum number of individuals needed to ensure a high probability of survival in the immediate future.

Finally, despite the lack of moisture as a general microclimate pattern in *inselbergs*, there are locally restricted habitats with vegetation that depend on seasonally humid conditions. For this reason, it is considered important to characterize the habitats within *inselbergs* to have enough conservation bases within these systems. For example, identify the environmental gradients where a vegetation type is developed, as well as the key species within communities. There are no doubts, that these *inselbergs* are an ideal scenario for the development of research in plant ecology.

References

- Huber O (1995) Geographical and physical features. In: Flora of the Venezuelan Guayana. Steyermark JA, Berry PE, Holst BK (Eds.), Missouri Botanical Garden, St Louis, Timber Press, Portland, USA, pp 1-63.
- Huber O (1995) Conservation of the Venezuelan Guayana. In: Flora of the Venezuelan Guayana. Steyermark JA, Berry PE, Holst BK (Eds.), Missouri Botanical Garden, St Louis, Timber Press, Portland, USA, pp 193-218.
- Groger A (2000) Flora and vegetacion of *inselbergs* of Venezuelan Guayana In: *Inselbergs – Biotic diversity of isolated rock outcrops in tropical and temperate regions*. Porembski S., Barthlott W (Eds.), Springer-Verlag, Berlin, Heidelberg, Germany 146: 291-313.
- Groger A, Huber O (2007) Rock outcrop habitats in the Venezuelan Guayana lowlands; their main vegetation types and floristic components. *Rev Brasil Bot* 30(4): 599 – 609.
- Porembski S (2007) Tropical *inselbergs*: habitat types, adaptive strategies and diversity patterns. *Revista Basil Bot* 30(4): 579 – 586.
- Villa PM, de Siqueira Cardinelli L, Magnago LF, Heringer G, Martins SV, et al. (2018) Relacion especie-area y distribucion de la abundancia de especies en una comunidad vegetal de un *inselberg* tropical: efecto del tamaño de los parches. *Rev Biol Trop* 66(2): 937-951.
- Porembski S (2000) Biodiversity of Terrestrial Habitat Islands – the *Inselbergs* Evidence In: *Inselbergs – Biotic diversity of isolated rock outcrops in tropical and temperate regions*. Porembski S, Barthlott W (Eds.), Springer-Verlag, Berlin, Heidelberg, Germany 146: 507-513.
- Kluge M, Budel B (2002) *Inselbergs: Vegetation, Diversity and Ecology*. *Trop Biol Conserva Manag* 4.
- Luttge U (1997) *Physiological ecology of tropical plants*. Springer-Verlag, Berlin, Heidelberg, Germany 320-346.
- Biedinger N, Porembski S, Barthlott W (2000) Vascular Plants on *Inselbergs: Vegetative and Reproductive Strategies*. In: *Inselbergs – Biotic diversity of isolated rock outcrops in tropical and temperate regions*. Porembski S, Barthlott W (Eds.), Springer-Verlag, Berlin, Heidelberg, Germany 146: 117-140.
- Bremer H, Sander H *Inselbergs: Geomorphology and Geoecology*. In: *Inselbergs – Biotic diversity of isolated rock outcrops in tropical and temperate regions*. Porembski S, Barthlott W (Eds.), Springer-Verlag, Berlin, Heidelberg, Germany 146: 7-34.

12. Villa PM, Gastauer M, Martins SV, Carrion JF, Campos PV, et al. (2018) Phylogenetic structure is determined by patch size in rock outcrop vegetation on an inselberg in the northern Amazon region. *Acta Amazonica* 48(3): 248-256.
13. Groger A (1994) Analisis preliminar de la florula y vegetacion del Monumento Natural *Piedra la Tortuga*, Estado Amazonas, Rheinische Friedrich - wilhelms - Universitat Bonn, Meckenheimer Allee 170 53115 Bonn, Germany.



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/JOJWB.2019.01.555555](https://doi.org/10.19080/JOJWB.2019.01.555555)

**Your next submission with Juniper Publishers
will reach you the below assets**

- Quality Editorial service
 - Swift Peer Review
 - Reprints availability
 - E-prints Service
 - Manuscript Podcast for convenient understanding
 - Global attainment for your research
 - Manuscript accessibility in different formats
- (Pdf, E-pub, Full Text, Audio)**
- Unceasing customer service

Track the below URL for one-step submission
<https://juniperpublishers.com/online-submission.php>