



Research Article

Volume 11 Issue 2 - August 2025
DOI: 10.19080/JPFMTS.2025.11.555810

J Phy Fit Treatment & Sports

Copyright © All rights are reserved by Simone Katrin Manai

Use of a simple stair-climbing test to assess cardiopulmonary fitness in clinical practice. An overview of the published literature aiming for a future goal



Johanna Wander¹, Maurice Pablo Mall¹, Anne Lentz¹, Nikolaus Alexander Haas^{1*}, Meike Schrader¹ and Simone Katrin Manai^{1*}

¹Department of Pediatric Cardiology and Pediatric Intensive Care Medicine, Ludwig-Maximilians University Hospital Munich-Großhadern, 81377 Munich, Germany

Submission: August 16, 2025; **Published:** August 28, 2025

***Corresponding author:** Simone Katrin Manai and Nikolaus Alexander Haas, Department of Pediatric Cardiology and Pediatric Intensive Care Medicine, Ludwig-Maximilians University Hospital Munich-Großhadern, 81377 Munich, Germany

Abstract

Background

Cardiorespiratory fitness (CRF) is a prognostic factor regarding long time morbidity and mortality. To assess CRF the well-known gold standard is cardiopulmonary exercise testing (CPET), which is a time consuming and expensive test. Therefore, alternative methods for routine evaluation of CRF are needed for the general population as well as for specific patient groups such as children and young adults with congenital heart disease. The Stair Climbing Test (SCT) is a simple and resource saving test reported in the current literature. This test is used in various settings to assess different health associated risk factors including cardiorespiratory fitness, muscular resilience, therapy effectiveness or pre- and postoperative morbidity and mortality.

Objectives

To summarize the current knowledge on the use and the different protocols of the SCT in the medical context.

Method: An internet-based literature search for research articles on both clinical trials and controlled randomized trials containing the SCT was undertaken.

Results

We used Pub Med and Google Scholar for literature research. A total of 200 articles were included. The SCT is used for multiple purposes on various patient groups. Significant correlations to the results from established clinical tests such as spiroergometry and 6-minute walking test were shown repetitively. However, the SCTs were conducted and evaluated in various, not standardized ways.

Conclusion

The SCT is a simple, cost saving test with promising reliability for the assessment of physical and cardiorespiratory fitness. Due to its easy approach, it can be used for various objectives, such as the general fitness and for specific patients-groups (i.e. children and young adults with congenital heart disease). Unfortunately, the SCT is not universally accepted yet due to the missing standardization. This standardization of the SCT protocol is however required to establish the SCT as a comprehensive test in the medical field.

Keywords: Stair Climbing Test; Cardiorespiratory Fitness; Exercise Test, Children; Congenital Heart Disease; spiroergometry; adolescents; physiotherapy

Abbreviations: CHD congenital heart disease; CPET cardiopulmonary exercise testing; CT clinical trial; CCT controlled clinical trial; HR heart rate; 6MWT 6-minute walking test; RCT randomized controlled trial; SCT stair climbing test; VO2max max. oxygen consumption/uptake

Introduction

Health-associated significance of cardiorespiratory- and physical fitness

During the last decades medicine and health care has significantly changed. Prevention is gaining more and more significance. Today it is generally accepted that cardiorespiratory fitness (CRF) is another important risk factor regarding cardiovascular disease, morbidity and mortality. The higher the level of CRF, the lower the mortality rate. [1] CRF not only reduces the risk of the occurrence of obvious diseases such as cardiovascular disease or comorbidities like high blood pressure and atrial fibrillation but is also playing an increasingly important role in other areas such as cancer prevention. [2,3]

Therefore, current WHO recommendations suggest a physical activity level for different age groups, e.g. for adults, including at least 150-300 minutes moderate-intensity aerobic physical activity or alternatively at least 75-150 minutes of vigorous-intensity aerobic physical activity per week. Physical activity is defined as any movement, that increases the resting metabolic rate. Nevertheless, a lot of people do not meet these recommendations. Especially in adolescents the recommendations are not fulfilled. [4]

This is of raising concern since the cornerstones of a healthy, active lifestyle are placed in early childhood. Furthermore, this is even more important in people with congenital diseases, for instance children and adolescents with congenital heart disease. [5] As the consequences of a sedentary lifestyle such as metabolic syndrome, diabetes mellitus, cardiovascular disease and others might reduce quality of life severely in these patient groups and increase overall health care costs, it is important to assess and evaluate cardiorespiratory fitness levels more often, preferentially as a routine check-up.

Cardiorespiratory fitness defines the ability of the lungs and the cardiovascular system to supply the skeletal musculature with oxygen during physical activity. If measured and interpreted accurately, the level of an individual's CRF can play an important role in the serious decision-making process for prevention and treatment plans. It could even be the decisive factor to evaluate whether the patient receives curative or palliative therapy. [6,7] Physical fitness defines the performance of the lungs, heart and skeletal muscles, therefore physical fitness testing actually includes the whole body and evaluates not only cardiorespiratory fitness but also, for example psychoneurological- or skeletomuscular function and mobility. [8]

Established tests for the evaluation of cardiorespiratory- and functional exercise capacity in the medical context

Currently, the clinical gold standard for evaluating cardiorespiratory fitness is cardiopulmonary exercise testing (CPET). [6,7] CPET evaluates the body's overall response to physical exertion and therefore incorporates several organ systems. [9] In the standard usage, CPET is used as a maximum exercise test, it involves

progressively increasing exercise intensity until reaching exhaustion or encountering symptoms or signs that impose limits. [10] Bicycle ergometer and treadmill are usually used for the physical exercise. [11]

While previously used clinical fitness tests such as bicycle ergometry simply recorded vital parameters including pulse, blood pressure and oxygen saturation, CPET adds a ventilatory expired gas analysis to these values. [7] By measuring the peak oxygen uptake (VO₂), CPET accurately identifies the highest exercise capacity of the person examined. [21] Nevertheless, CPET currently is not used in routine check-ups, as it is a costly and time-consuming examination, which requires special equipment and trained health care professionals. It is not universally available and often limited to special institutions.

The other commonly utilized test for evaluation of the functional exercise capacity is the 6-min walk test (6MWT). [13] For the 6MWT the patient is tasked with walking (along a 30-meter route) for 6 minutes, aiming to cover the maximum distance achievable. The key metric assessed is the 6-minute walk distance (6MWD), measured in meters. [13] Three major advantages of the 6-minute walk test are that it is easy to perform, that patients usually tolerate it well and that it is inexpensive. [14] Particularly the ease with which the 6MWT can be performed and the good tolerance of patients makes it suitable for patient groups with limited fitness such as patients with heart failure. [14] Nevertheless, it must be considered, that the 6MWT is a submaximal exercise test for most patients. Only in patients with a significantly reduced CRF it can be used as a maximum exercise test.

Currently there is no cheap, universally available maximum exercise test to assess cardiorespiratory fitness. With the gaining importance of the cardiorespiratory fitness level regarding preventive and therapeutic approaches, a lot of working groups investigate alternative approaches such as the Stair-Climbing-Test (SCT).

The SCT is used in various approaches and can give a lot of information about an individual's general fitness, as well as its CRF and its motor function. Depending on its approach it also can differentiate between a loss in strength and a lack of coordination as well as whether there is a respiratory or cardiological restriction.

The aim of this review is to give an overview and compare the different SCT approaches used so far.

Materials and Methods

Search strategy

We conducted an internet-based literature search. For this purpose, we primarily used Pub Med as well as Google Scholar and the Cochrane Library to cross check. The search terms "stair climbing test OR stair-climbing test" was used on Pub Med to find original research articles on the topic. (The search was last con-

ducted on January 03rd 2024) A total of 973 articles matched our initial searching criteria.

Inclusion Criteria

Randomized controlled trials, clinical trials and pilot studies, available as full text in English language that used the SCT in any form and described its use and implementation were included. Our literature search was not restricted to an age group.

Exclusion Criteria

We excluded articles that were not relevant to our research question, review articles, meta-analysis and articles that were only available as abstracts. Studies that used the SCT but did not describe the exact application and design were also eliminated.

Data Extraction

Of the 973 articles that matched our initial inclusion criteria on Pub Med, 319 mentioned the SCT and were accessible as full text. Ultimately, 200 of them matched our final inclusion criteria

and were therefore included in our review. The others were discarded because they either mentioned the SCT but did not describe its implementation in more detail, or they were not original texts, or because they were not thematically relevant to our research question.

Results

In the Pub Med search a total of 200 articles matched the final searching criteria, described the usage of the stair climbing test and were therefore included in this review. In these 200 trials the SCT was used with different intentions and for the evaluation of various body functions. They were performed between 1986 and 2023. The most frequently investigated parameter during SCT was physical and neuromuscular function. (table 1) Cardiorespiratory fitness and exercise tolerance was evaluated in a smaller number of trials (table 2). Some studies evaluated additional effects such as blood sugar, the ability of complex thinking and others (table 3).

Table 1. 155 Studies evaluating neuromuscular/physical function

Author, Study Design, Number of Participants	Study Background	SCT-outcome	SCT-technical background	SCT-implementation	SCT- parameters surveyed
0	Effects of Resistance Training on Strength, Power, and Selected Functional Abilities of Women Aged 75 and Older	functional ability	6 flights 12 steps Flight height: 1,885 m	climb up a staircase as far as possible without stopping at a comfortable pace without using the handrail.	-flights per second -heart rate (mean of the final 15 seconds)
Schwid SR et al. 1997 CT, n= 10 [91]	Quantitative assessment of sustained-release 4-aminopyridine for symptomatic treatment of multiple sclerosis	motor function	4 steps step height: 15,24 cm	climb four steps of stairs as quickly as possible.	-time to ascend
Sharp SA et al. 1997 CT, n= 15 [99]	Isokinetic Strength Training of the Hemiparetic Knee: Effects on Function and Spasticity	physical function	1 set, 4 stairs, step height: 17,7 cm	climbed up at a comfortable speed, using their normal pattern of foot placement and hand support.	-time to ascend (average of 3 trials) -calculated cadence (stairs/minute)
Teixeira-Salmela LF et al. 1999 CT, n= 13 [100]	Muscle Strengthening and Physical Conditioning to Reduce Impairment and Disability in Chronic Stroke Survivors	functional performance	5 steps, Step height: 17,7 cm	climb up 5 steps of stairs at a comfortable speed, using their usual patterns of foot placement and hand support. Two trials were completed.	-average time and cadence (stairs per minute)
Kovacs I et al. 2001 CT, n=58 [53]	The therapeutic effects of Cserkeszo "lo" thermal water in osteoarthritis of the knee: a double blind, controlled, follow-up study	physical function	20 steps	ascend and then to descend 20 steps	-time

Sartorio A et al. 2001 CT, n=230 [106]	Changes in motor control and muscle performance after a short-term body mass reduction program in obese subjects	motor control maximal lower limb muscle power	13 steps Step height: 15.3 cm total height: 1.99 m	climb up ordinary stairs at the highest possible speed. Instructor classified the performance	-time -calculated power output
Rutkove SB et al. 2002 RCT, n= 16 [211]	A pilot randomized trial of oxandrolone in inclusion body myositis	physical function	steps unlimited	had to climb as many stairs as possible in 15 seconds.	-number of steps climbed in 15 seconds (best out of two)
Hiroyuki S et al. 2003 CT, n= 34 [125]	Specific effects of balance and gait exercises on physical function among the frail elderly	walking assessment	5 steps step height: 15 cm	climb and descend 5 steps of stairs as fast as possible.	-total time
Katz-Leurer M et al. 2003 RCT, n= 92 [93]	The Influence of Early Aerobic Training on the Functional Capacity in Patients With Cerebrovascular Accident at the Subacute Stage	functional walking	steps unlimited	climb as many stairs as possible at a comfortable speed. Any assisted device was allowed. The test ended when the patient felt fatigue.	-number of stairs climbed
Ljungquist T et al. 2003 CT, n= 186 [83]	Physical performance tests for people with spinal pain—sensitivity to change	physical performance	1 flight, different heights	ascend and descend 1 flight of stairs at a self-selected speed.	-total time
Bonan IV et al. 2004 RCT, n= 20 [101]	Reliance on Visual Information After Stroke. Part II: Effectiveness of a Balance Rehabilitation Program With Visual Cue Deprivation After Stroke	gait	1 set 10 steps	ascend and descend a set of 10 stairs.	-total time
Grant S et al. 2004 CT n= 26 [107]	The effects of a 12-week group exercise programme on physiological and psychological variables and function in overweight women	physical performance	12 steps, step height: 16 cm	climb up 12 stairs, turn on the landing and descend as fast as possible without using the handrail.	-time to ascend -time to descend -total time
Kraemer WJ et al. 2004 CT, n=40 [54]	Effect of a Cetylated Fatty Acid Topical Cream on Functional Mobility and Quality of Life of Patients with Osteoarthritis	functional Mobility	1 flight 11 steps Step height: 13,5 cm	ascend and descend a flight of eleven steps as quickly as possible. 3 to 5 trials were performed, the best times were recorded. Use of the handrails was allowed; 2 members of the research staff accompanied the patient to assure maximal safety.	-total time -time to ascend -time to descend
Mengshoel AM et al. 2004 CT, n=31+26 [86]	Associations between walking time, quadriceps muscle strength and cardiovascular capacity in patients with rheumatoid arthritis and ankylosing spondylitis	physical function	22 steps step height 20 cm	walk as rapidly as possible without running up and down a staircase without a railing.	-time

Molsted S et al. 2004 CT, n= 33 [157]	Five Months of Physical Exercise in Hemodialysis Patients: Effects on Aerobic Capacity, Physical Function and Self-Rated Health	physical function	2 flights	ascend and descend two flights of stairs as quickly as possible for 2 minutes.	-number of steps (ascending and descending)
Seynnes O et al. 2004 CT, n= 22 [126]	Physiological and Functional Responses to Low-Moderate Versus High-Intensity Progressive Resistance Training in Frail Elders	functional limitation	4 steps, Step height: 0,15 m Step length: 0.30 m	climb 4 risers of stairs as fast as possible without using a hand-rail. They did 3 repetitions with 2 minutes break in between.	- Stair-climbing power (force 3distance/ time), defined as body weight 3 vertical height climbed/ time to ascend steps, expressed in watts
Zaino et al. 2004 CT, n= 47 [104]	Timed Up and Down Stairs Test: Preliminary Reliability and Validity of a New Measure of Functional Mobility	functional mobility musculoskeletal and neuromuscular systems	14-steps step height 19.5-cm	stand 30 cm from the bottom of a flight of stairs "Quickly, but safely go up the stairs, turn around on the top step (landing) and come all the way down until both feet land on the bottom step (landing)." The subjects were allowed to choose any method of traversing the stairs.	-time
Capodaglio D et al. 2005 CT, n=60 [129]	Muscle function and functional ability improves more in community-dwelling older women with a mixed-strength training programme	functional ability	2 flights 12 steps	climb up a staircase as quickly as possible without stopping and without using the handrail as support, to turn around on the top platform and then walk down.	-time
Galvao DA et al. 2005 RCT, n= 28 [148]	Resistance Exercise Dosage in Older Adults: Single- Versus Multiset Effects on Physical Performance and Body Composition	physical Performance	1 flight, 11 stairs, step height: 16 cm	climb a flight of stairs as fast as possible while staying safe and without use of the handrails.	-time to ascend
Laufer Y et al. 2005 CT, n= 103 [55]	Effect of pulsed short-wave diathermy on pain and function of subjects with osteoarthritis of the knee	functional mobility	1 flight 15 stairs step height: 15 cm	First trial: climb 15 steps, using the handrail or a cane was allowed Second trial: descend the same 15 steps of stairs.	-time to ascend -time to descend
Mizner RL et al. 2005 CT, n=40 [18]	Preoperative Quadriceps Strength Predicts Functional Ability One Year After Total Knee Arthroplasty	functional performance	12 steps step height: 18 cm step depth: 28 cm	climb as quickly as they felt safe and comfortable. use of one handrail was allowed if necessary, but encouraged to minimize their use of the handrail. 1practice test 2 tests Assistive devices were allowed only if the subject was unsafe or could not complete the test without the assistance of a cane or walker	- time (average)

Mizner RL et al. 2005 CT, n=40 [17]	Quadriceps Strength and the Time Course of Functional Recovery After Total Knee Arthroplasty	functional performance	1 flight 12 steps step height: 18cm depth 28 cm	ascend and descend a flight as quickly as it feels safe and comfortable. handrail was allowed if required	-time
Storer TW et al. 2005 CT, n= 12 [158]	Endurance exercise training during haemodialysis improves strength, power, fatigability and physical performance in maintenance haemodialysis patients	physical performance	1 staircase, 4-steps Total height: 0,625 m	ascend a 4-step staircase as fast as possible. One test round, three trials, best time out of the three was taken as stair-climb score.	-time to ascend (stair-climb score) - Power (calculated from subject's body weight, vertical ascent and ascent time)
Eyyigor et al. 2006 CT, n= 20 [150]	Effects of a group-based exercise program on the physical performance, muscle strength and quality of life in older women	Physical performance	10 steps step height 20cm	ascend a staircase and turn on the landing and descend the stairs without stopping and without using the handrail for support.	-time to ascend -time to descend
Galvao DA et al. 2006 CT, n= 10 [122]	Resistance Training and Reduction of Treatment Side Effects in Prostate Cancer Patients	physical performance	1 flight, 13 stairs, step height: 17 cm	climb the flight of stairs as fast as possible while staying safe.	-total time
Henwood TR et al. 2006 RCT, n= 67 [127]	Short-term resistance training and the older adult: the effect of varied programmes for the enhancement of muscle strength and functional performance	functional performance	1 flight 11 steps step height: 16 cm	ascend 11 stairs; without the use of a handrail.	-time to ascend -stairclimbing power (using time to ascend, body weight, gravity, step height, number of steps)
Bar-Haim et al. 2007 CT, n=36 [105]	Prediction of mechanical efficiency from heart rate during stair-climbing in children with cerebral palsy	mechanical efficiency	4 steps step height adjustable 1-17 cm,	walk up and down 4 steps for 4 minutes at a pace of their choice. stopping time and number of ascents recorded.	-number of ascents -stopping time -work - breath-by-breath V _{O2} and HR during exercise
Capodaglio P et al. 2007 CCT, n= 58 [128]	Long-term strength training for community-dwelling people over 75: impact on muscle function, functional ability, and lifestyle	functional abilities	2 flights 12 steps each	climb up as fast as possible without stopping and without using the handrail as support, to turn on the top and walk down again.	-total time
Nyland J et al. 2007 Retro. study n=31 [19]	Self-reported chair-rise ability relates to stair-climbing readiness of total knee arthroplasty patients: A pilot study	functional performance	1 flight 10 steps step height 7-inch (17.8 cm) Step depth 11-inch (27.9 cm)	ascend a flight as quickly as possible without compromising safety. Following a 30-second rest period at the top of the steps, subjects were instructed to descend the steps as quickly as possible without compromising safety. Using a handrail was allowed	-time

Smeets RJEM et al. 2007 RCT, n= 221 [84]	Physical capacity tasks in chronic low back pain	physical capacity	five steps circuit, shaped like an eight	walk a stair up and down for 1 min.	-number of steps climbed
Westlake KP et al. 2007 CT, n=46+24 [130]	Velocity discrimination: Reliability and construct validity in older adults	Physical performance	13 standard steps	ascend and descend steps using one rail at a "quick, but safe speed". 1-min rest period was allowed at the top and bottom landing. 2 tests	-ascend time -descend time
Bruun-Olsen V et al. 2008 RCT, n= 57 [20]	The immediate and long-term effects of a walking-skill program compared to usual physiotherapy care in patients who have undergone total knee arthroplasty (TKA)	physical function	16 stairs, step height: 16 cm	ascend and descend 16 steps using alternate legs, it was allowed to support themselves by holding onto the rail.	-total time
Dreher M et al. 2008 RCT n= 16 [160]	Exercise in severe COPD: Is walking different from stair-climbing	physical function	44 steps step height: 0.16 m	climb up 44 steps. One group with supplemental oxygen during exercise, one without.	-total time -vitals -blood gas -blood lactate
Galea MP et al. 2008 RCT, n= 23 [43] [116]	A Targeted Home- and Center-Based Exercise Program for People After Total Hip Replacement Effects of a supra-physiological dose of testosterone on physical function, muscle performance, mood, and fatigue in men with HIV-associated weight loss	physical function physical performance	1 set, 4 stairs, step height: 17,5 cm step depth: 29,8 cm adjustable rail 12 steps, 4 middle steps recorded (0,66m height)	climb 4 steps as fast as possible. Using the handrail was allowed, without pulling themselves up. Patients did one test trial and completed 2 trials. ascend 12 steps as rapidly as possible. ascent time recorded over the middle four steps (0.66 m). Handrail holding was allowed for balance.	-time to ascend -lower-limb power (using weight, height of stairs and time to calculate) -ascent time over the middle four steps -stairclimbing power (the product of body mass, total step rise, and the acceleration of gravity all divided by time to traverse the middle four steps)
Kortebein P et al. 2008 CT, n=11 [131]	Functional Impact of 10 Days of Bed Rest in Healthy Older Adults	lower extremity strength and power	stairs	climb stairs as fast as comfortably possible with one hand near, but not on, the handrail.	-stair ascent power participant's weight (N) and the time in seconds to ascend 10 steps (Power 1/4 (Distance/Time) 3 Weight)
Lee MJ et al. 2008 RCT, n= 52 [94]	Comparison of Effect of Aerobic Cycle Training and Progressive Resistance Training on Walking Ability After Stroke	walking ability	10 steps	ascend a standardized flight of 10 stairs as quickly and safely as possible, using a hand-rail for support was allowed.	-time to ascend - Stair climb power (calculated from the time taken to ascend the stairs, the known vertical height, and body mass)
Schmitt LC et al. 2008 CT n=52 [56]	Instability, Laxity, and Physical Function in Patients With Medial Knee Osteoarthritis	physical function	12 stairs step height: 18 cm	ascend and descend a set of 12 stairs as quickly as they felt safe and comfortable. They were encouraged not to use the handrail, but were not prohibited from doing so for safety.	-time

Storer TW et al. 2008 RCT n=44 [152]	Changes in Muscle Mass, Muscle Strength, and Power; but not Physical Function are Related to Testosterone Dose in Healthy Older Men	physical function	SCT 1: 4 steps, total height: 0,66 m SCT 2: 12 steps, total recorded height: 0,69 m	SCT 1: climb up 4 steps of stairs as fast as possible without using the handrail. SCT 2: climb up 12 steps of stairs, but only from step 4-8 the time was recorded.	-time to ascend 4 steps of stairs -stair climbing power (fastest time achieved, subjects' body weight, total rise height, and the acceleration of gravity)
Vogt L et al. 2008 CT, n= 179 [147]	Cognitive status and ambulatory rehabilitation outcome in geriatric patients	ambulatory status	1 flight step height: 19 cm step depth: 28 cm	ascend and descend a flight of steps step by step as fast and comfortably as possible.	-performance (rated by using a 4-point ordinal scale scoring system based on the subject's difficulty in performing the task and the use of the handrail for support and balance)
Bruun-Olsen V et al. 2009 RCT, n= 63 [212]	Continuous passive motion as an adjunct to active exercises in early rehabilitation following total knee arthroplasty	physical function and mobility	8 steps, step height: 16 cm	walk up and down eight steps, using alternate legs no handrail or use of a walking aid.	-total time
Eyigor S et al. 2009 RCT, n= 40 [149]	A trial of Turkish folklore dance on the physical performance, balance, depression, and quality of life in older women	physical performance	10 steps, step height: 20 cm	climb up 10 steps of stairs, turn on the landing and descend, as fast as possible. Using the handrail for support was not allowed.	-total time
Farquhar S et al. 2009 CT, n=183 [22]	The Chitranjan Ranawat Award The Nonoperated Knee Predicts Function 3 Years after Unilateral Total Knee Arthroplasty	physical function	1 flight 12 steps step height 18 cm	ascend and descend a flight as safely and as quickly as possible; use of one handrail is allowed if required. One practice test, two tests used for	-time
Harmer AR et al. 2009 RCT, n=102 [21]	Land-Based Versus Water-Based Rehabilitation Following Total Knee Replacement	functional mobility	18 steps 1 landing in between	ascend 18 stairs as rapidly as possible. Using handrails and walking aids as required was allowed.	-time to ascend -stair climbing power (=calculated by using body mass, total stair height and ascent time)
LeBrasseur KL et al. 2009 RCT, n=252 [132]	Effects of testosterone therapy on muscle performance and physical function in older men with mobility limitations (The TOM Trial): design and methods	lower extremity function mobility	12 steps	ascend a flight of stairs as fast as possible and allowed to use the handrail only if needed. using a switch mat timing system (Lafayette Instrument Company, Lafayette, IN).	-time
Petterson SC et al. 2009 RCT, n= 200 [23]	Improved Function from Progressive Strengthening Interventions After Total Knee Arthroplasty	functional performance	12 steps step height: 7,9 cm	ascend and descend 12 steps. Two trials were completed.	-average total time of two trials

Pua YH et al. 2009 CT, n=92 [67]	Self-Report and Physical Performance Measures of Physical Function in Hip Osteoarthritis: Relationship to Isometric Quadriceps Torque Development	Physical performance	6 steps step height: 18 cm step depth: 30 cm	climb up and down 6 stairs in their usual manner. Handrails were on the right side of the stairs, and participants held them loosely for safety if necessary.	-time
Yoshida Y et al. 2009 CT, n=12 [24]	Examining outcomes from total knee arthroplasty and the relationship between quadriceps strength and knee function over time	Physical function	1 flight	go up and down a flight of stairs as quickly as an individual feels safe and comfortably referred to literature	-time
Andersson et al. 2010 CT, n= 198 [85]	Performance Tests in People with Chronic Low Back Pain	physical performance	5 stairs unlimited	climb up and down 5 stairs for 1 minute. Using the handrail was allowed.	-number of steps climbed in 1min
Heiberg KE et al. 2010 CT, n=63 [41]	Pain and recovery of physical functioning nine months after total knee arthroplasty	physical function	8 steps step height 16 cm	walking up and down a flight of stairs, using alternate legs, with no support from a rail or walking aid.	-time
Hirota et al. 2010 CT, n = 493 [133]	Association between the Trail Making Test and physical performance in elderly Japanese	movement parameters	4 steps, step height: 10 cm	ascend and descend 4 steps as fast as possible, using the handrail if needed.	-total time
Stevens-Lapsley JE et al. 2010 CT, n=140 [25]	Impact of Body Mass Index on Functional Performance After Total Knee Arthroplasty	functional performance	12 steps step height 20.1 cm	ascend and descend 12 steps	-time
Wetzel JL et al. 2010 CT, n=64 [87]	Six-Minute Walk Test for Persons with Mild or Moderate Disability from Multiple Sclerosis: Performance and Explanatory Factors	functional lower-extremity strength power	4 steps	ascend, turn, and descend, using a handrail as necessary.	-power: calculated using the following equation: (number of steps * step height [m] * body weight) -time
Zeni JA Jr et al. 2010 CT, n=40 [27]	Clinical predictors of elective total joint replacement in persons with end-stage knee osteoarthritis	functional mobility	12 stairs	ascend the steps on the investigators command, turn around and descend the stairs. Light handrail use was permitted for balance.	-time
Zeni Jr JA et al. 2010 CT, n=155 [26]	Early Postoperative Measures Predict 1- and 2-Year Outcomes After Unilateral Total Knee Arthroplasty: Importance of Contralateral Limb Strength MK-0677 (ibutamoren mesylate) for the treatment of patients recovering from hip fracture	functional performance lower extremity power	1 flight 12 steps 4 steps	ascend and descend a flight of 12 steps as quickly as possible in a safe manner. Use of the handrail allowed if needed for balance. 1 test and then 2 timed trials, (average time was recorded). ascend 4 steps of stairs as fast as possible.	-time -time to ascend -stairclimbing power (=product of patient's weight, gravitational force, and vertical velocity (staircase height/time))

Christiansen CL et al. 2011 CT, n=36+17 [28]	Weight-Bearing Asymmetry During Sit-Stand Transitions Related to Impairment and Functional Mobility After Total Knee Arthroplasty	functional performance	12-steps	ascend, turn around, and descend a flight of stairs. 2 trials. There was a handrail on the staircase, which participants were encouraged to not use during the test.	-time
Sattler F et al. 2011 RCT, n= 112 [120]	Testosterone Threshold Levels and Lean Tissue Mass Targets Needed to Enhance Skeletal Muscle Strength and Function	muscle performance and physical function	12 steps	ascend 12 steps of stairs, time to ascend the middle four steps in the 12-step staircase was measured. photocells used to measure time for stair climbing power	-time to ascend the middle 4 steps in 12-step staircase -stair climbing power
Travison TG et al. 2011 RCT, n=165 [121]	Clinical Meaningfulness of the Changes in Muscle Performance and Physical Function Associated with Testosterone Administration in Older Men With Mobility Limitation	physical function	12-step staircase	ascend 12 steps with and without weights equal to 20% of their body weight. Two trials were required.	-time to ascend -stair-climbing power (product of body weight plus weight carried, total stair-rise, divided by ascent time)
Bade MJ et al. 2012 CT, n=118 [46]	Predicting Poor Physical Performance after Total Knee Arthroplasty	functional performance	12 steps steps height 18 cm depth 28 cm	ascend, turn around, and then descend the steps as quickly as possible in a safe manner. Bilateral handrails were available for use if needed, 2tests	-time
Heiberg CE et al. 2012 RCT, n= 68 [16]	Effect of a Walking Skill Training Program in Patients Who Have Undergone Total Hip Arthroplasty	physical functioning	8 steps, step height: 16 cm	ascend and descend 8 steps as fast as possible without running, instructed to use alternate legs, using the stair rail was allowed.	-time to ascend
Hsieh RL et al. 2012 RCT, n= 72 [213]	Short-Term Effects of 890-Nanometer Radiation on Pain, Physical Activity, and Postural Stability in Patients with Knee Osteoarthritis	physical activity	14 steps, step height: 18 cm	ascend and descend a flight of stairs as fast as possible.	-total time
van de Port IGL et al. 2012 RCT, n= 250 [103]	Effects of circuit training as alternative to usual physiotherapy after stroke	functional mobility	5 steps chair placed 0,5 meters from the stairs	modified SCT: combination of the timed up and go test before ascending and descending 5 steps of stairs and then sitting down again. The required time from elevating from the chair to sitting again after the stair climb was measured.	-total time

Alfano LN et al. 2013 CT, n=25 [89]	Correlation of knee strength to functional outcomes in Becker muscular dystrophy	functional performance	4 steps step height: 6 inches = 15,24cm	ascend and descend the stairs as quickly and safely as possible. The use of handrails or other compensatory movement patterns was permitted.	-time
Baert IAC et al. 2013 CT, n=87 [78]	Weak associations between structural changes on MRI and symptoms, function and muscle strength in relation to knee osteoarthritis	physical function	5 steps	ascend, turn around and descend 3 tests, mean value	-time
Baert IAC et al. 2013 CT, n=45+20 [77]	Proprioceptive accuracy in women with early and established knee osteoarthritis and its relation to functional ability, postural control, and muscle strength	physical function	5 steps step height 15 cm	ascend, turn around, and descend five steps. mean value of 3 trials	-time
Basaria A et al. 2013 RCT, n= 76 [162]	The Safety, Pharmacokinetics, and Effects of LGD-4033, a Novel Nonsteroidal Oral, Selective Androgen Receptor Modulator, in Healthy Young Men	physical performance (in terms of safety, tolerability)	12-steps, step height: 17 cm	ascend 12 steps as fast as possible. One test round, then two trials	-total time (best out of two) - SCT- Power (calculated from the time elapsed, body weight, and vertical distance)
Chalé A et al. 2013 RCT, n= 80 [153]	Efficacy of Whey Protein Supplementation on Resistance Exercise-Induced Changes in Lean Mass, Muscle Strength, and Physical Function in Mobility-Limited Older Adults	physical function	10 steps	ascend a 10-rise set of stairs as fast as possible, without hold on to the railing or use assistive devices. They did two trials; average time was recorded.	-average time to ascend
Chung JY et al. 2013 CT, n= 24 [29]	Is bicompartamental knee arthroplasty more favourable to knee muscle strength and physical performance compared to total knee arthroplasty	physical performance	12 steps, step height: 20 cm step depth: 30 cm	ascend and descend 12 steps as fast as possible while still staying safe. Using the handrail was allowed. Two repetitions.	-total time (average of two trials)
van Leeuwen DM et al. 2013 CT, n=22 [30]	Preoperative Strength Training for Elderly Patients Awaiting Total Knee Arthroplasty	physical function	9 steps	ascend, turn around, and descend "walk as quickly and safely" allowed to use the handrail and instructed to	-time
Vincent HK et al. 2013 CT, n =53 [57]	"Functional Pain," Functional Outcomes, and Quality of Life After Hyaluronic Acid Intra-articular Injection for Knee Osteoarthritis	physical function	1 flight, 12 steps	climb up 12 steps as fast as possible. Repeated 3 times, fastest trial time was used for data analysis.	-time to ascend

Akbaba YA et al. 2014 RCT, n=20+20+20 [31]	Intensive supervision of rehabilitation programme improves balance and functionality in the short term after bilateral total knee arthroplasty	physical function	10 steps step height 19 cm depth 27 cm	climb up and down the stairway, as fast as possible	-time
Bieler L et al. 2014 RCT, n= 122 [63]	Intra-rater reliability and agreement of muscle strength, power, and functional performance measures in patients with hip osteoarthritis	functional performance	10 steps, step height: 16,3 cm step depth: 35,8 cm	ascend and descend a flight of 10 steps without using the handrail. The best result of 2 timed trials was used noted.	-total time
Hsieh RL et al. 2014 CT, n=40 [75]	Immediate and medium-term effects of custom-moulded in-soles on pain, physical function, physical activity, and balance control in patients with knee osteoarthritis	physical function	14 steps step height 18 cm	ascend a flight of stairs as quickly as possible.	-time
Judd DL et al. 2014 CT, n=26+18 [151]	Strength and Functional Deficits in Individuals with Hip Osteoarthritis Compared to Healthy, Older Adults	functional performance	12 stairs	climb a flight of 12 stairs, turn around at the top and descend the same flight as quickly and safely as possible. They were permitted to use the handrail for balance but were instructed not to use the handrail to push or pull	-time
Marmon AR et al. 2014 CT, n=24 [47]	Associations between knee extensor power and functional performance in patients after total knee arthroplasty and normal controls without knee pain	Functional performance	12 steps	ascend and descend 12 stairs	-time
Marmon AR et al. 2014 CT, n=84+68 [79]	Perception and Presentation of Function in Patients with Unilateral Versus Bilateral Knee Osteoarthritis	functional ability	12 stairs	ascend and then descend a flight of 12 stairs as quickly and safely as possible. use of the handrail was allowed if necessary.	-time
Tsukagoshi R et al. 2014 RCT, n= 65 [39]	Functional performance of female patients more than 6 months after total hip arthroplasty shows greater improvement with weight-bearing exercise than with non-weight-bearing exercise	functional performance	10 steps step height: 17,5 cm	ascend 10 steps of stairs.	-time to ascend

Winters JD et al. 2014 restro. study, n=35+23 [38]	Preliminary Investigation of Rate of Torque Development Deficits Following Total Knee Arthroplasty	physical function	12 steps height 18 cm depth 28 cm.	ascend, turn around, and then descend the steps as quickly as possible in a safe manner. The handrail was available for use if needed. 2 tests	-time
Zeni, Jr. J et al. 2014 CT, n=56 [80]	Relationship between strength, pain, and different measures of functional ability in patients with end-stage hip osteoarthritis	physical function	12 standard steps	ascend and descend 12 standard steps. A handrail is available during testing	-time
Zhang M et al. 2014 CT, n=72 [159]	Relation Between Anxiety, Depression and Physical Activity and Performance in Maintenance Hemodialysis Patients	physical performance	22 steps	climb the stairs as fast as possible without running, jumping or skipping steps, and were allowed to use the banister for balance if necessary	-time
Altubasi IM et al. 2015 CT, n=21 [134]	Is quadriceps muscle strength a determinant of the physical function of the elderly?	physical function	1 flight 11 steps step height 17 cm	climb up as fast as possible. 1 practice round 2 trials.	-time to ascend
Becker C et al. 2015 CT, n=201 [146]	Myostatin antibody (LY2495655) in older weak fallers: a proof-of-concept	physical performance	4 steps 12 steps step height: 15-18 cm	climb up 4 and 12 steps of stairs while using the handrail.	-time to ascend
Brenneman EC et al. 2015 CT, n=38 [58]	A Yoga Strengthening Program Designed to Minimize the Knee Adduction Moment for Women with Knee Osteoarthritis	mobility	9 steps	ascend and descend a 9-step staircase as fast and safely as possible without running and without skipping stairs. Using the handrail was allowed. Two trials were completed.	-mean time to ascend -mean time to descend
Chikani V et al. 2015 CT, n=13+13 [123]	Impairment of Anaerobic Capacity in Adults With Growth Hormone Deficiency	physical function leg muscle power	4 flights 48 steps step height 17 cm	ascend four flights as fast as possible, one step at a time. 3 tests separated by a rest period of 5 minutes.	-time
Dias CP et al. 2015 RCT, n= 26 [135]	Effects of eccentric-focused and conventional resistance training on strength and functional capacity of older adults	functional capacity	8 steps step height: 17 cm step length: 31 cm	climb eight steps without using the handrail. Two trials with 3 min of rest between	-fastest performance time
Harries N et al. 2015 CT, n= 43 [102]	A stair-climbing test for measuring mechanical efficiency of ambulation in adults with chronic stroke.	motor performance	4 steps, step height: adjustable 10 to 20 cm	climb up and down the stairs for 5 min, self-pacing their speed, keeping it constant, using the handrail for assistance if needed. They did two repetitions of the test.	-time

Pirotta S et al. 2015 RCT, n= 26 [136]	Effects of vitamin D supplementation on neuroplasticity in older adults	physical function	10 steps, step height: 7,8 cm	ascend 10 steps of stairs as fast as possible without using the handrail or walking aids.	-time to ascend -stair climbing power
Akbaba YA et al. 2016 RCT, n= 60 [31]	Intensive supervision of rehabilitation programme improves balance and functionality in the short term after bilateral total knee arthroplasty	physical functionality	10 steps, step height: 19cm step depth: 27 cm	climb up and down ten steps, as fast as possible. Two trials	-total time (mean value of 2 trials)
Bittel TC et al. 2016 CT, n=79 [109]	Adipose tissue content, muscle performance and physical function in obese adults with type 2 diabetes mellitus and peripheral neuropathy	Physical function	1 flight 10 steps	climbing a flight of stairs with 10 steps	-time -power (calculated from the time taken to ascend the stairs, the known vertical height, gravitational force and body mass)
Collado-Mateo D et al. 2016 CT, n = 20 [90]	Performance of women with fibromyalgia in walking up stairs while carrying a load	step-by- step-performance and trunk tilt	10 steps step height: 17 cm depth: 28 cm	climb 10 stairs without carrying a load, rest for 3 min, repeat the test carrying a load of 5 kg in each hand. sensors for motion capture were used.	-kinematic data
Heiberg KE et al. 2016 RCT, n= 60 [42]	Physical Functioning and Prediction of Physical Activity After Total Hip Arthroplasty: Five-Year Follow-up	physical function	8 steps step height: 16 cm	ascend and descend 8 steps as fast as possible without running, allowed to use the handrail, but not a walking aid.	-total time
Hsieh RL et al. 2016 RCT, n=90 [76]	Clinical effects of lateral wedge arch support insoles in knee osteoarthritis	physical activity	1 flight 14 steps step height: 18 cm	ascend and descend 14 steps in the shortest time possible.	-total time
Reddy S et al. 2016 CT, n=264 [161]	Timed Stair Climbing is the Single Strongest Predictor of Perioperative Complications in Patients Undergoing Abdominal Surgery	physical performance	7 steps	walk down and then up one single flight. Vital signs were collected prior to beginning and immediately after completing the task.	-time
Bade M et al. 2017 RCT, n=162 [45]	Early High-Intensity Versus Low-Intensity Rehabilitation after Total Knee Arthroplasty	physical function	1 flight, 12 stairs, step height: 17,1 cm	climb up and down 12 flights of stairs	-total time
Baldwin JN et al. 2017 CT, n=1000 [164]	Reference values and factors associated with musculoskeletal symptoms in healthy adolescents and adults	stair-climbing ability	flight of stairs	timed up and Down Stairs Test by Zaino et al	-time

Baldwin JN et al. 2017 CT, n=1000 [163]	Relationship between physical performance and self-reported function in healthy individuals across the lifespan	stair-climbing ability	flight of stairs	Timed up and Down Stairs Test by Zaino et al	-time
Bieler T et al. 2017 RCT, n= 152 [64]	In hip osteoarthritis, Nordic Walking is superior to strength training and home-based exercise for improving function	functional performance	10 steps, step height: 16,3 cm step depth: 35,8 cm	ascend and descend a flight of 10 steps without using the handrail as fast as possible.	-total time (best out of two trials)
Freisinger GM et al. 2017 CT, n=33 [65]	Relationships Between Varus-Valgus Laxity of the Severely Osteoarthritic Knee and Gait, Instability, Clinical Performance, and Function	physical function	12 steps	ascend and descend a staircase as quickly as possible in a safe manner. Encouraged not to use the handrail unless necessary to complete the test.	-time
Johnen B et al. 2017 Pilot study, n=45 [137]	Feasibility of a machine vs free weight strength training program and its effects on physical performance in nursing home residents: a pilot study	physical performance	11 steps	climb 11 risers of stairs as fast as possible)	-time
Loyd BJ et al. 2017 retro. study, n=162 [32]	Influence of Hip Abductor Strength on Functional Outcomes Before and After Total Knee Arthroplasty: Post Hoc Analysis of a Randomized Controlled Trial	physical performance	12 steps	ascend and descend a set of 12 steps with or without the use of a handrail for balance. This task was timed and the faster of the 2 trials used for analysis.	-time
Maffiuletti NA et al. 2017 CT, n=40 [110]	Reproducibility of clinician-friendly physical performance measures in individuals with obesity	physical function	13 stairs height 15 cm depth 32 cm	referred to Perron et al. stand up from a chair; walk 3 m, ascend at a comfortable pace, turn around and descend stairs, walk back to the chair; turn and sit down.	-time
Nilsen TS et al. 2017 CT, n=58 [117]	Effects of strength training on body composition, physical functioning, and quality of life in prostate cancer patients during androgen deprivation therapy	physical function	stairs	SCT referred to literature with and without additional 20kg	-time
Romine PE et al. 2017 CT, n=430 [138]	Task-Specific Fatigue Among Older Primary Care Patients	physical function	2 m high flight	climbing a flight of stairs twice, using a handrail and/or cane if needed. 2tests, average used	-time
Sions JM et al. 2017 CT, n=106 [139]	Multifidi Muscle Characteristics and Physical Function among Older Adults with and without Chronic Low Back Pain	physical function	2 steps depth: 28cm, height: 17cm	fast stair descent, "as quickly and as safely as possible". The average of two trials was calculated	-time

Storer TW et al. 2017 RCT, n= 308 [118]	Effects of Testosterone Supplementation for 3 years on Muscle Performance and Physical Function in Older Men	physical function	12 steps	four repetitions of ascending the stairs as fast as possible without running, first two tests unloaded, second two test whilst carrying weights equivalent to 20 % of their body weight.	-time to ascend -stair-climb Power (=product of the total rise of the 12 steps, body weight plus load carried, and acceleration of gravity, all divided by time)
Suh MJ et al. 2017 CT, n=34 [49]	Effects of Early Combined Eccentric-Concentric Versus Concentric Resistance Training Following Total Knee Arthroplasty	physical function	1 flight 12 steps step height: 17cm step wide: 25 cm	ascend and descend a flight of stairs as fast as possible on the word "go". 5-minute rest interval between tests, the best score was recorded	-time
Gagliano-Juca T et al. 2018 RCT, n=99 [119]	Testosterone does not affect agrin cleavage in mobility-limited older men despite improvement in physical function	physical function	12 steps	climb up and down 12 steps of stairs as fast as possible without running while carrying weight equal to 20 % of their body weight.	-total time -stair-climbing power (=product of body weight plus weight carried, total stair-rise, and acceleration of gravity all divided by ascent time)
Mulla DM et al. 2018 RCT, n= 43 [165]	The Effects of Lower Extremity Strengthening Delivered in the Workplace on Physical Function and Work-Related Outcomes Among Desk-Based Workers	mobility	1 flight 11 steps	ascend and descend a flight of stairs with 11 steps as fast as possible without running and without skipping steps. It was permitted to use the handrail. Two trials were completed.	-average time to ascend -average time to descend
Mustafaoğlu R et al. 2018 Pilot study, n=45 [95]	The effects of body weight-supported treadmill training on static and dynamic balance in stroke patients: A pilot, single-blind, randomized trial	physical function	10 steps step height 17 cm	ascending and descending using handrails or assistive devices were allowed, if necessary	-time
Iijima H et al. 2019 CT, n=57 [69]	Stair climbing ability in patients with early knee osteoarthritis: Defining the T clinical hallmarks of early disease	physical function	11 steps step height 17 cm step width 135 cm step tread 29 cm	wearing the standardized shoes with pressure sensor-mounted insoles, descend and ascend the flight	-time -vertical ground reaction force (GRF) was calculated from sensor-mounted shoe data -power of 11-SCT for identifying early knee OA (Kellgren and Lawrence grade 1)
Judd DL et al. 2019 CT, n=79 [52]	Trajectories of functional performance and muscle strength recovery differ after total knee and total hip replacement: A performance-based, longitudinal study.	physical function	12 stairs	ascend and descend 12 stairs	-time

Kim JH et al. 2019 Retro. Study, n=184 [48]	Functional Outcomes After Critical Pathway for Inpatient Rehabilitation of Total Knee Arthroplasty	functional mobility lower extremity muscles forces	1 flight 12 steps step height: 17 cm step wide: 25 cm	ascend and descend a flight of stairs	-time
Lange-Maia BS et al. 2019 CT, n=829 [140]	Factors influencing longitudinal stair climb performance from midlife to early late life: The Study of Women's Health Across the Nation Chicago and Michigan Sites	physical function	4 standard stairs	ascend and descend the stairs for three consecutive cycles. Using the handrail was allowed if needed.	-time
Moukarzel M et al. 2019 RCT, n=24 [33]	The therapeutic role of motor imagery during the chronic phase after total knee arthroplasty	strength and functional mobility	12 steps step height: 18 cm step depth: 28 cm	ascend and descend 12 steps of stairs as fast as possible. One test round and two trials.	-total time
Nunes GS et al. 2019 CT, n = 32 [88]	People with patellofemoral pain have impaired functional performance, that is correlated to hip muscle capacity	physical function	9 steps step height: 17 cm	ascend and descend 9 steps as fast as possible step by step. Using the handrail was allowed.	-total time
Orange ST et al. 2019 CCT, n= 36 [141]	Short- term training and detraining effects of supervised vs. unsupervised resistance exercise in aging adults	physical performance	1 free standing flight of stairs with 5 steps, step height 20 cm	ascend and descend as quickly possible while staying safe. Using the handrails was permitted if needed.	-total time
Shimoura K et al. 2019 RCT, n=50 [70]	Immediate Effects of Transcutaneous Electrical Nerve Stimulation on Pain and Physical Performance in Individuals with Preradiographic Knee Osteoarthritis	physical function	11 steps step height: 17 cm	ascend and descend the stairway as fast as possible, started with both feet on the bottom landing and ended in the same position. 2 trials. Using the handrail for support was allowed.	-total time
Sousa-Gonçalves CR et al. 2019 CT, n=8 [111]	Acute Effects of Whole-Body Vibration Alone or in Combination With Maximal Voluntary Contractions on Cardiorespiratory, Musculoskeletal, and Neuromotor Fitness in Obese Male Adolescents	Musculoskeletal and neuromotor fitness	13 steps step height: 15.3 cm total height: 1.99 m	climb up ordinary stairs at the highest possible speed, according to their capabilities	-time -power: (calculated from the time taken to ascend the stairs, the known vertical height, and body mass)
Suh et al. 2019 CT, n= 195 [74]	Bilateral Quadriceps Muscle Strength and Pain Correlate With Gait Speed and Gait Endurance Early After Unilateral Total Knee Arthroplasty	physical function	1 flight 12 steps step height 17 cm wide: 25 cm	ascend and descend a flight of stairs as quickly as possible. 3trial with five minutes break in between.	-total time (best out of three)

Unhjem R et al. 2019 CT, n=41 [142]	Functional Performance With Age: The Role of Long-Term Strength Training	functional performance	12 steps step height: 17 cm	climbing as fast as possible. Maximum step length was set to 2 stair steps at a time, no handrail support was allowed.	-power (W)
Bade MJ et al. 2010 CCT, n=24 [44]	Outcomes Before and After Total Knee Arthroplasty Compared to Healthy Adults	functional performance	10 steps step height 17.1-cm 12-steps step height 17.1-cm	9 patients were tested on a 10-step staircase, 15 patients and all healthy adults were tested on a 12-step staircase	-time
Daly RM et al. 2020, RCT, n= 216 [154]	Effects of a multivitamin-fortified milk drink combined with exercise on functional performance, muscle strength etc. in middle-aged women	functional muscle power	10 stairs, step height: 17 cm	ascend or descend a flight of 10 stairs as fast as possible without missing a step. Using the handrail if necessary was allowed. They did 1 practice trial and then 3 test trials.	-time to ascend -time to descend -stair climb power on ascend (power (Watts) = $9.81 \times \text{body mass (kg)} \times \text{vertical step height (m)} \times \text{number of steps / time (s)}$)
Gränicher P et al. 2020 Pilot RCT, n=20 [34]	Preoperative exercise in patients undergoing total knee arthroplasty: a pilot randomized controlled trial	functional performance	8 steps height 16 cm depth 30 cm	ascend and descend a flight at usual walking speed, feeling safe and comfortable. handrail or assistive devices were allowed but not encouraged	-time
Larsen JB et al. 2020 Retro. Study, n=217 [40]	Intensive, personalized multimodal rehabilitation in patients with primary or revision total knee arthroplasty: a retrospective cohort study	lower body strength balance	22 stairs	ascend and descend stairs as quickly and as safely as possible. Use of a handrail and walking aid were permitted if needed.	-time
Lee SJ et al. 2020 CT, n=84 [51]	Preoperative physical factors that predict stair-climbing ability at one month after total knee arthroplasty	physical function	12 steps step height 17 cm 25 cm wide	ascend or descend the stairs as quickly as possible on the word "go". 3 trial with a 5-min rest interval Using a handrail was allowed	-time
Mustafaoglu R et al. 2020 RCT, n= 51 [96]	Does robot-assisted gait training improve mobility, activities of daily living and quality of life in stroke	mobility	10 steps step height: 18 cm	climb up and down ten steps without skipping any steps, using one foot for each step and descend without stopping. Use of handrail and/or assistive devices was allowed.	-total time
Onodera CMK et al. 2020 N=153 [60]	The importance of objectively measuring functional tests in complement to self-report assessments in patients with knee osteoarthritis	physical function	9 steps and 9 landings step height 18 cm	reach up and down the stairs	-time

Pozzi F et al. 2020 RCT, n=293 [37]	Restoring physical function after knee replacement: a cross-sectional comparison of progressive strengthening vs standard physical therapy	physical function	12 stairs, (15cm rise, 20cm run)	ascend and descend the set of stairs as fast as possible without skipping steps. Using the handrail was allowed. Time started on command and ended when both feet touched the bottom again.	-total time
Rigamonti AE et al. 2020 CT, n=595 [113]	Impact of a Three-Week in-Hospital Multidisciplinary Body Weight Reduction Program on Body Composition, Muscle Performance and Fatigue in a Pediatric Obese Population with or without Metabolic Syndrome	muscle performance	13 steps step height: 15.3 cm total height: 1.99 m	climb up ordinary stairs at the highest possible speed. 2–3 test trials.	-time
Rigamonti AE et al. 2020 CT, n= 1922 [214]	Effects of a 3-Week In-Hospital Body Weight Reduction Program on Cardiovascular Risk Factors, Muscle Performance, and Fatigue: A Retrospective Study in a Population of Obese Adults with or without Metabolic Syndrome	muscle function	13 steps step height 15.3 cm total height 1.99 m	climb up ordinary stairs at the highest possible speed	-time
Tamini S et al. 2020 CT, n=16 [112]	Acute Effects of Whole-Body Vibration Exercises at 2 Different Frequencies Versus an Aerobic Exercise on Some Cardiovascular, Neuro-motor and Musculoskeletal Parameters in Adult Patients With Obesity	maximal lower limb muscle power motor control	13 steps step height: 15.3 cm total height: 1.99 m	climb up ordinary stairs at their highest speed, in accordance to their capabilities.	-time -power (=product of 1/4patient's weight, gravitational force, and vertical height /time)
Vongsirinavarat M et al. 2020 CT, n=250 [59]	Identification of knee osteoarthritis disability phenotypes regarding activity limitation: a cluster analysis	physical function	Not specified	Timed stair climbing referred to literature	-time
Ahmed Burq HSI et al. 2021, RCT, n= 64 [97]	Effect of whole-body vibration on obstacle clearance and stair negotiation time in chronic stroke patients	mobility function	3 steps step height: 18 cm step depth: 30 cm	ascend and descend 3 steps of stairs in a comfortable speed, accompanied by a therapist. The handrails were used.	-total time
Beckmann M et al. 2021 CT, n=207 [81]	Recovery and prediction of physical function 1 year following hip fracture	physical function	8 steps	ascended and descended 8steps as fast as able without running, using alternate legs and support by stair rail if needed.	-time

Choi JH et al. 2021 CT, n=149 [36]	Performance-based physical function correlates with walking speed and distance at 3 months post unilateral total knee arthroplasty	physical function	12 steps height 17 cm wide 25 cm	ascend and descend a flight of 12 steps, as fast as possible on the word "go". 3 trials, with the 5 min rest in between	-time
Jacksteit R et al. 2021 RCT, n=85 [35]	Low-Load Unilateral and Bilateral Resistance Training to Restore Lower Limb Function in the Early Rehabilitation After Total Knee Arthroplasty: A Randomized Active-Controlled Clinical Trial	physical performance postural control strength of the lower extremities	8 steps step height: 17.5 cm	climb a staircase in a safely and quickly manner using a railing and regular footwear.	-time
Katsoulis K et al. 2021 Retro. study, n=18 [143]	Reliability of Lower Extremity Muscle Power and Functional Performance in Healthy, Older Women	physical function	13 steps step height 18 cm step width 28 cm height 2.34 m	stand at the base of the stairs with feet together and to grab the handrail if necessary during ascent. Upon the instruction "ready, set, go," stair ascent was measured during a "usual" pace (SCUP) and during a "fast" pace (SCFP) with instructions to ascend "as quickly and as safely as possible."	-time -power = ((body mass in kg) × (9.8 m/s ²) × (stair height in meter))/(time in seconds)
Kim BS et al. 2021 CT, n=562 [73]	Associations Between Obesity With Low Muscle Mass and Physical Function in Patients With End-Stage Knee Osteoarthritis	Physical function	1 flight 12 steps step height: 17 cm step length: 25 cm	ascend or descend the stairs as fast as possible upon hearing the word "go". 3 trials, with a 5-minute rest interval between. The fastest time was recorded for each patient	-ascend time -descend time
Khruakhorn S et al. 2021 RCT, n=34 [61]	Effects of hydrotherapy and land-based exercise on mobility and quality of life in patients with knee osteoarthritis: a randomized control trial	physical function	4 steps step width: 26.5 cm step length: 76 cm step height: 15.2 cm	"Walk up – turn back – and go down the stairs as soon as possible but safely" and subsequently started the assessment with the word "Start". 2 trials with a 5-minute rest between were evaluated.	-time
Bayartai ME et al. 2022 CT, n=160 [108]	Changes in the Oswestry Disability Index after a 3-Week In-Patient Multidisciplinary Body Weight Reduction Program in Adults with Obesity	lower limb muscle power functional ability	13 steps step height: 15.3 cm total height: 1.99 m	climb a staircase at maximum speed. 1-2 test trials	-time
De Zwart AH et al. 2022 CT, n=177 [62]	Association Between Measures of Muscle Strength and Performance of Daily Activities in Patients with Knee Osteoarthritis	functional performance	12 steps	ascending and descending (independently) a staircase as fast as possible without running	-time

Ihl T et al. 2022 RCT, n=1771 [98]	Patient-Centered Outcomes in a Randomized Trial Investigating a Multimodal Prevention Program After Transient Ischemic Attack or Minor Stroke: The INSPIRE-TMS Trial	physical activity	first landing of a stairway	ascend as quickly as possible to the first landing of a stairway. Patients were allowed to use the banisters, if necessary	-power (power=body weight×9.8×height of the staircase/mean time of 2 runs)
Ke D et al. 2022 CT, n = 103 [155]	Study of the Reliability of Field Test Methods for Physical Fitness in Children Aged 2–3 Years	physical fitness coordination ability	5 steps step height 12 cm	climb up the stairs with using both feet alternately and without using the handrail. 2 trials.	-time to ascend (best out of two)
Lee SH et al. 2022 CCT, n=30+30 [72]	Validity of the Osteoarthritis Research Society International (OARSI) recommended performance-based tests of physical function in individuals with symptomatic Kellgren and Lawrence grade 0–2 knee osteoarthritis	physical function	9 stairs step height 20 cm	ascended and descended as quickly as possible but in a safe manner.	-time
Lindberg K et al. 2022, RCT, n= 49 [144]	Effectiveness of individualized training based on force– velocity profiling on physical function in older men	physical function	15-steps, Step height: 16 cm	climb up 15 steps of stairs as fast as possible with and without a weight vest of 20 kg.	-time to ascend
Lazzer S et al. 2022 CT, n=139 [114]	Effects of a 3-Week Inpatient Multidisciplinary Body Weight Reduction Program on Body Composition and Physical Capabilities in Adolescents and Adults With Obesity	muscle function	13 steps step height 15.3 cm total height 1.99 m	climb an ordinary stair at the highest possible speed	-time
Petersson N et al. 2022 CT, n=14 [66]	Blood Flow Restricted Walking in Elderly Individuals with Knee Osteoarthritis: A Feasibility Study	Functional performance	11 steps	11step stair climb test referring to another trial	-time
Usubini AG et al. 2022 CT, n=237 [115]	A three-week in-hospital multidisciplinary body weight reduction program exerts beneficial effects on physical and mental health and fatiguability of elderly patients with obesity	muscle function	13 steps step height 15.3 cm to- tal height 1.99 m	climb up ordinary stairs at the highest possible speed	-time -power (kg × 9.81 × 1.99) /s
Carvalho C et al. 2023 CT, n=39 [68]	Association between ankle torque and performance-based tests, self-reported pain, and physical function in patients with knee osteoarthritis	Functional performance	1 flight 11 steps step height: 17 cm step width: 202 cm step tread: 31 cm	go up and down a flight of stairs as quickly and safely as possible.	-time

Fosstveit SH et al. 2023 CT, n=49 [145]	Associations between Power Training-Induced Changes in Body Composition and Physical Function in Older Men: A Pre-Test-Post-Test Experimental Study	Physical function	15 steps Step height: 16 cm	climb 15 steps as fast as possible. The time was recorded using photocells placed at the bottom and top of the stairs at 85 cm height. Warm-up, 2 unloaded tests and 2 with a 20 kg vest	-time
Jankaew A et al. 2023, RCT, n=47 [71]	The effects of low-level laser therapy on muscle strength and functional outcomes in individuals with knee osteoarthritis	functional performance	13 steps, step height: 18 cm	climb up the stairs, turn around, and climb down as quickly as possible while staying safe.	-total time
Kirschner N et al. 2023 CT, n=24 [50]	Determination of Relationships between Symmetry-Based, Performance-Based, and Functional Outcome Measures in Patients Undergoing Total Hip Arthroplasty	physical performance	14 steps	14-step stair-climbing test Preoperatively with no walking aids. Postoperatively, walking support and the stair railing were used.	-time
Lanzi S et al. 2023 CT, n=90 [156]	Time-course evolution of functional performance during a 3-month supervised exercise training program in patients with symptomatic peripheral artery disease	functional performance	12 stairs	climb as quickly as possible using a handrail was allowed. 2 tests	-time
Suslov VM et al. 2023 CT, n=28 [92]	Efficacy and safety of hydrokinesitherapy in patients with dystrophinopathy	physical function	4 steps	descent 4-stairs	-time

Table 2. 37 Studies evaluating cardiorespiratory fitness/exercise capacity

Author, Study Design, Number of Participants	Study Background	SCT-outcome	SCT-technical background	SCT-implementation	SCT- parameters surveyed
0	Effects of Resistance Training on Strength, Power, and Selected Functional Abilities of Women Aged 75 and Older	functional ability	6 flights 12 steps Flight height: 1,885 m	climb up a staircase as far as possible without stopping at a comfortable pace without using the handrail.	-flights per second -heart rate (mean of the final 15 seconds)
Schwid SR et al. 1997 CT, n= 10 [91]	Quantitative assessment of sustained-release 4-aminopyridine for symptomatic treatment of multiple sclerosis	motor function	4 steps step height: 15,24 cm	climb four steps of stairs as quickly as possible.	-time to ascend
Sharp SA et al. 1997 CT, n= 15 [99]	Isokinetic Strength Training of the Hemiparetic Knee: Effects on Function and Spasticity	physical function	1 set, 4 stairs, step height: 17,7 cm	climbed up at a comfortable speed, using their normal pattern of foot placement and hand support.	-time to ascend (average of 3 trials) -calculated cadence (stairs/minute)

Teixeira-Salmela LF et al. 1999 CT, n= 13 [100]	Muscle Strengthening and Physical Conditioning to Reduce Impairment and Disability in Chronic Stroke Survivors	functional performance	5 steps, Step height: 17,7 cm	climb up 5 steps of stairs at a comfortable speed, using their usual patterns of foot placement and hand support. Two trials were completed.	-average time and cadence (stairs per minute)
Kovacs I et al. 2001 CT, n=58 [53]	The therapeutic effects of Cserkeszo "lo" thermal water in osteoarthritis of the knee: a double blind, controlled, follow-up study	physical function	20 steps	ascend and then to descend 20 steps	-time
Sartorio A et al. 2001 CT, n=230 [106]	Changes in motor control and muscle performance after a short-term body mass reduction program in obese subjects	motor control maximal lower limb muscle power	13 steps Step height: 15.3 cm total height: 1.99 m	climb up ordinary stairs at the highest possible speed. Instructor classified the performance	-time -calculated power output
Rutkove SB et al. 2002 RCT, n= 16 [211]	A pilot randomized trial of oxandrolone in inclusion body myositis	physical function	steps unlimited	had to climb as many stairs as possible in 15 seconds.	-number of steps climbed in 15 seconds (best out of two)
Hiroyuki S et al. 2003 CT, n= 34 [125]	Specific effects of balance and gait exercises on physical function among the frail elderly	walking assessment	5 steps step height: 15 cm	climb and descend 5 steps of stairs as fast as possible.	-total time
Katz-Leurer M et al. 2003 RCT, n= 92 [93]	The Influence of Early Aerobic Training on the Functional Capacity in Patients With Cerebrovascular Accident at the Subacute Stage	functional walking	steps unlimited	climb as many stairs as possible at a comfortable speed. Any assisted device was allowed. The test ended when the patient felt fatigue.	-number of stairs climbed
Ljungquist T et al. 2003 CT, n= 186 [83]	Physical performance tests for people with spinal pain—sensitivity to change	physical performance	1 flight, different heights	ascend and descend 1 flight of stairs at a self-selected speed.	-total time
Bonan IV et al. 2004 RCT, n= 20 [101]	Reliance on Visual Information After Stroke. Part II: Effectiveness of a Balance Rehabilitation Program With Visual Cue Deprivation After Stroke	gait	1 set 10 steps	ascend and descend a set of 10 stairs.	-total time
Grant S et al. 2004 CT n= 26 [107]	The effects of a 12-week group exercise programme on physiological and psychological variables and function in overweight women	physical performance	12 steps, step height: 16 cm	climb up 12 stairs, turn on the landing and descend as fast as possible without using the handrail.	-time to ascend -time to descend -total time

Kraemer WJ et al. 2004 CT, n=40 [54]	Effect of a Cetylated Fatty Acid Topical Cream on Functional Mobility and Quality of Life of Patients with Osteoarthritis	functional Mobility	1 flight 11 steps Step height: 13,5 cm	ascend and descend a flight of eleven steps as quickly as possible. 3 to 5 trials were performed, the best times were recorded. Use of the handrails was allowed; 2 members of the research staff accompanied the patient to assure maximal safety.	-total time -time to ascend -time to descend
Mengshoel AM et al. 2004 CT, n=31+26 [86]	Associations between walking time, quadriceps muscle strength and cardiovascular capacity in patients with rheumatoid arthritis and ankylosing spondylitis	physical function	22 steps step height 20 cm	walk as rapidly as possible without running up and down a staircase without a railing.	-time
Molsted S et al. 2004 CT, n= 33 [157]	Five Months of Physical Exercise in Hemodialysis Patients: Effects on Aerobic Capacity, Physical Function and Self-Rated Health	physical function	2 flights	ascend and descend two flights of stairs as quickly as possible for 2 minutes.	-number of steps (ascending and descending)
Seynnes O et al. 2004 CT, n= 22 [126]	Physiological and Functional Responses to Low-Moderate Versus High-Intensity Progressive Resistance Training in Frail Elders	functional limitation	4 steps, Step height: 0,15 m Step length: 0.30 m	climb 4 risers of stairs as fast as possible without using a hand-rail. They did 3 repetitions with 2 minutes break in between.	- Stair-climbing power (force 3distance/time), defined as body weight 3 vertical height climbed/time to ascend steps, expressed in watts
Zaino et al. 2004 CT, n= 47 [104]	Timed Up and Down Stairs Test: Preliminary Reliability and Validity of a New Measure of Functional Mobility	functional mobility musculoskeletal and neuromuscular systems	14-steps step height 19.5-cm	stand 30 cm from the bottom of a flight of stairs "Quickly, but safely go up the stairs, turn around on the top step (landing) and come all the way down until both feet land on the bottom step (landing)." The subjects were allowed to choose any method of traversing the stairs.	-time
Capodaglio D et al. 2005 CT, n=60 [129]	Muscle function and functional ability improves more in community-dwelling older women with a mixed-strength training programme	functional ability	2 flights 12 steps	climb up a staircase as quickly as possible without stopping and without using the handrail as support, to turn around on the top platform and then walk down.	-time
Galvao DA et al. 2005 RCT, n= 28 [148]	Resistance Exercise Dosage in Older Adults: Single- Versus Multiset Effects on Physical Performance and Body Composition	physical Performance	1 flight, 11 stairs, step height: 16 cm	climb a flight of stairs as fast as possible while staying safe and without use of the handrails.	-time to ascend

Laufer Y et al. 2005 CT, n= 103 [55]	Effect of pulsed short-wave diathermy on pain and function of subjects with osteoarthritis of the knee	functional mobility	1 flight 15 stairs step height: 15 cm	First trial: climb 15 steps, using the handrail or a cane was allowed Second trial: descend the same 15 steps of stairs.	-time to ascend -time to descend
Mizner RL et al. 2005 CT, n=40 [18]	Preoperative Quadriceps Strength Predicts Functional Ability One Year After Total Knee Arthroplasty	functional performance	12 steps step height: 18 cm step depth: 28 cm	climb as quickly as they felt safe and comfortable. use of one handrail was allowed if necessary, but encouraged to minimize their use of the handrail. 1 practice test 2 tests Assistive devices were allowed only if the subject was unsafe or could not complete the test without the assistance of a cane or walker	- time (average)
Mizner RL et al. 2005 CT, n=40 [17]	Quadriceps Strength and the Time Course of Functional Recovery After Total Knee Arthroplasty	functional performance	1 flight 12 steps step height: 18cm depth 28 cm	ascend and descend a flight as quickly as it feels safe and comfortable. handrail was allowed if required	-time
Storer TW et al. 2005 CT, n= 12 [158]	Endurance exercise training during haemodialysis improves strength, power, fatigue and physical performance in maintenance haemodialysis patients	physical performance	1 staircase, 4-steps Total height: 0,625 m	ascend a 4-step staircase as fast as possible. One test round, three trials, best time out of the three was taken as stair-climb score.	-time to ascend (stair-climb score) - Power (calculated from subject's body weight, vertical ascent and ascent time)
Eyyigor et al. 2006 CT, n= 20 [150]	Effects of a group-based exercise program on the physical performance, muscle strength and quality of life in older women	Physical performance	10 steps step height 20cm	ascend a staircase and turn on the landing and descend the stairs without stopping and without using the handrail for support.	-time to ascend -time to descend
Galvao DA et al. 2006 CT, n= 10 [122]	Resistance Training and Reduction of Treatment Side Effects in Prostate Cancer Patients	physical performance	1 flight, 13 stairs, step height: 17 cm	climb the flight of stairs as fast as possible while staying safe.	-total time
Henwood TR et al. 2006 RCT, n= 67 [127]	Short-term resistance training and the older adult: the effect of varied programmes for the enhancement of muscle strength and functional performance	functional performance	1 flight 11 steps step height: 16 cm	ascend 11 stairs; without the use of a handrail.	-time to ascend -stairclimbing power (using time to ascend, body weight, gravity, step height, number of steps)

Bar-Haim et al. 2007 CT, n=36 [105]	Prediction of mechanical efficiency from heart rate during stair-climbing in children with cerebral palsy	mechanical efficiency	4 steps step height adjustable 1-17 cm,	walk up and down 4 steps for 4 minutes at a pace of their choice. stopping time and number of ascents recorded.	-number of ascents -stopping time -work - breath-by-breath V _{O2} and HR during exercise
Capodaglio P et al. 2007 CCT, n= 58 [128]	Long-term strength training for community-dwelling people over 75: impact on muscle function, functional ability, and lifestyle	functional abilities	2 flights 12 steps each	climb up as fast as possible without stopping and without using the handrail as support, to turn on the top and walk down again.	-total time
Nyland J et al. 2007 Retro. study n=31 [19]	Self-reported chair-rise ability relates to stair-climbing readiness of total knee arthroplasty patients: A pilot study	functional performance	1 flight 10 steps step height 7-inch (17.8 cm) Step depth 11-inch (27.9 cm)	ascend a flight as quickly as possible without compromising safety. Following a 30-second rest period at the top of the steps, subjects were instructed to descend the steps as quickly as possible without compromising safety. Using a handrail was allowed	-time
Smeets RJEM et al. 2007 RCT, n= 221 [84]	Physical capacity tasks in chronic low back pain	physical capacity	five steps circuit, shaped like an eight	walk a stair up and down for 1 min.	-number of steps climbed
Westlake KP et al. 2007 CT, n=46+24 [130]	Velocity discrimination: Reliability and construct validity in older adults	Physical performance	13 standard steps	ascend and descend steps using one rail at a "quick, but safe speed". 1-min rest period was allowed at the top and bottom landing. 2 tests	-ascend time -descend time
Bruun-Olsen V et al. 2008 RCT, n= 57 [20]	The immediate and long-term effects of a walking-skill program compared to usual physiotherapy care in patients who have undergone total knee arthroplasty (TKA)	physical function	16 stairs, step height: 16 cm	ascend and descend 16 steps using alternate legs, it was allowed to support themselves by holding onto the rail.	-total time
Dreher M et al. 2008 RCT n= 16 [160]	Exercise in severe COPD: Is walking different from stair-climbing	physical function	44 steps step height: 0.16 m	climb up 44 steps. One group with supplemental oxygen during exercise, one without.	-total time -vitals -blood gas -blood lactate

Galea MP et al. 2008 RCT, n= 23 [43] [116]	A Targeted Home- and Center-Based Exercise Program for People After Total Hip Replacement Effects of a supra-physiological dose of testosterone on physical function, muscle performance, mood, and fatigue in men with HIV-associated weight loss	physical function physical performance	1 set, 4 stairs, step height: 17,5 cm step depth: 29,8 cm adjustable rail 12 steps, 4 middle steps recorded (0,66m height)	climb 4 steps as fast as possible. Using the handrail was allowed, without pulling themselves up. Patients did one test trial and completed 2 trials. ascend 12 steps as rapidly as possible. ascent time recorded over the middle four steps (0.66 m). Handrail holding was allowed for balance.	-time to ascend -lower-limb power (using weight, height of stairs and time to calculate) -ascent time over the middle four steps -stairclimbing power (the product of body mass, total step rise, and the acceleration of gravity all divided by time to traverse the middle four steps)
Kortebein P et al. 2008 CT, n=11 [131]	Functional Impact of 10 Days of Bed Rest in Healthy Older Adults	lower extremity strength and power	stairs	climb stairs as fast as comfortably possible with one hand near, but not on, the handrail.	-stair ascent power participant's weight (N) and the time in seconds to ascend 10 steps (Power 1/4 (Distance/Time) 3 Weight)
Lee MJ et al. 2008 RCT, n= 52 [94]	Comparison of Effect of Aerobic Cycle Training and Progressive Resistance Training on Walking Ability After Stroke	walking ability	10 steps	ascend a standardized flight of 10 stairs as quickly and safely as possible, using a hand-rail for support was allowed.	-time to ascend - Stair climb power (calculated from the time taken to ascend the stairs, the known vertical height, and body mass)
Schmitt LC et al. 2008 CT n=52 [56]	Instability, Laxity, and Physical Function in Patients With Medial Knee Osteoarthritis	physical function	12 stairs step height: 18 cm	ascend and descend a set of 12 stairs as quickly as they felt safe and comfortable. They were encouraged not to use the handrail, but were not prohibited from doing so for safety.	-time
Storer TW et al. 2008 RCT n=44 [152]	Changes in Muscle Mass, Muscle Strength, and Power, but not Physical Function are Related to Testosterone Dose in Healthy Older Men	physical function	SCT 1: 4 steps, total height: 0,66 m SCT 2: 12 steps, total recorded height: 0,69 m	SCT 1: climb up 4 steps of stairs as fast as possible without using the handrail. SCT 2: climb up 12 steps of stairs, but only from step 4-8 the time was recorded.	-time to ascend 4 steps of stairs -stair climbing power (fastest time achieved, subjects' body weight, total rise height, and the acceleration of gravity)
Vogt L et al. 2008 CT, n= 179 [147]	Cognitive status and ambulatory rehabilitation outcome in geriatric patients	ambulatory status	1 flight step height: 19 cm step depth: 28 cm	ascend and descend a flight of steps step by step as fast and comfortably as possible.	-performance (rated by using a 4-point ordinal scale scoring system based on the subject's difficulty in performing the task and the use of the handrail for support and balance)
Bruun-Olsen V et al. 2009 RCT, n= 63 [212]	Continuous passive motion as an adjunct to active exercises in early rehabilitation following total knee arthroplasty	physical function and mobility	8 steps, step height: 16 cm	walk up and down eight steps, using alternate legs no handrail or use of a walking aid.	-total time

Eyigor S et al. 2009 RCT, n= 40 [149]	A trial of Turkish folklore dance on the physical performance, balance, depression, and quality of life in older women	physical performance	10 steps, step height: 20 cm	climb up 10 steps of stairs, turn on the landing and descend, as fast as possible. Using the handrail for support was not allowed.	-total time
Farquhar S et al. 2009 CT, n=183 [22]	The Chitranjan Ranawat Award The Nonoperated Knee Predicts Function 3 Years after Unilateral Total Knee Arthroplasty	physical function	1 flight 12 steps step height 18 cm	ascend and descend a flight as safely and as quickly as possible; use of one handrail is allowed if required. One practice test, two tests used for	-time
Harmer AR et al. 2009 RCT, n=102 [21]	Land-Based Versus Water-Based Rehabilitation Following Total Knee Replacement	functional mobility	18 steps 1 landing in between	ascend 18 stairs as rapidly as possible. Using handrails and walking aids as required was allowed.	-time to ascend -stair climbing power (=calculated by using body mass, total stair height and ascent time)
LeBrasseur KL et al. 2009 RCT, n=252 [132]	Effects of testosterone therapy on muscle performance and physical function in older men with mobility limitations (The TOM Trial): design and methods	lower extremity function mobility	12 steps	ascend a flight of stairs as fast as possible and allowed to use the handrail only if needed. using a switch mat timing system (Lafayette Instrument Company, Lafayette, IN).	-time
Petterson SC et al. 2009 RCT, n= 200 [23]	Improved Function from Progressive Strengthening Interventions After Total Knee Arthroplasty	functional performance	12 steps step height: 7,9 cm	ascend and descend 12 steps. Two trials were completed.	-average total time of two trials
Pua YH et al. 2009 CT, n=92 [67]	Self-Report and Physical Performance Measures of Physical Function in Hip Osteoarthritis: Relationship to Isometric Quadriceps Torque Development	Physical performance	6 steps step height: 18 cm step depth: 30 cm	climb up and down 6 stairs in their usual manner. Handrails were on the right side of the stairs, and participants held them loosely for safety if necessary.	-time
Yoshida Y et al. 2009 CT, n=12 [24]	Examining outcomes from total knee arthroplasty and the relationship between quadriceps strength and knee function over time	Physical function	1 flight	go up and down a flight of stairs as quickly as an individual feels safe and comfortably referred to literature	-time
Andersson et al. 2010 CT, n= 198 [85]	Performance Tests in People with Chronic Low Back Pain	physical performance	5 stairs unlimited	climb up and down 5 stairs for 1 minute. Using the handrail was allowed.	-number of steps climbed in 1min
Heiberg KE et al. 2010 CT, n=63 [41]	Pain and recovery of physical functioning nine months after total knee arthroplasty	physical function	8 steps step height 16 cm	walking up and down a flight of stairs, using alternate legs, with no support from a rail or walking aid.	-time

Hirota et al. 2010 CT, n = 493 [133]	Association between the Trail Making Test and physical performance in elderly Japanese	movement parameters	4 steps, step height: 10 cm	ascend and descend 4 steps as fast as possible, using the handrail if needed.	-total time
Stevens-Lapsley JE et al. 2010 CT, n=140 [25]	Impact of Body Mass Index on Functional Performance After Total Knee Arthroplasty	functional performance	12 steps step height 20.1 cm	ascend and descend 12 steps	-time
Wetzel JL et al. 2010 CT, n=64 [87]	Six-Minute Walk Test for Persons with Mild or Moderate Disability from Multiple Sclerosis: Performance and Explanatory Factors	functional lower-extremity strength power	4 steps	ascend, turn, and descend, using a handrail as necessary.	-power: calculated using the following equation: (number of steps * step height [m] * body weight) -time
Zeni JA Jr et al. 2010 CT, n=40 [27]	Clinical predictors of elective total joint replacement in persons with end-stage knee osteoarthritis	functional mobility	12 stairs	ascend the steps on the investigators command, turn around and descend the stairs. Light handrail use was permitted for balance.	-time
Zeni Jr JA et al. 2010 CT, n=155 [26]	Early Postoperative Measures Predict 1- and 2-Year Outcomes After Unilateral Total Knee Arthroplasty: Importance of Contralateral Limb Strength MK-0677 (ibutamoren mesylate) for the treatment of patients recovering from hip fracture	functional performance lower extremity power	1 flight 12 steps 4 steps	ascend and descend a flight of 12 steps as quickly as possible in a safe manner. Use of the handrail allowed if needed for balance. 1 test and then 2 timed trials, (average time was recorded). ascend 4 steps of stairs as fast as possible.	-time -time to ascend -stairclimbing power (=product of patient's weight, gravitational force, and vertical velocity (staircase height/time))
Christiansen CL et al. 2011 CT, n=36+17 [28]	Weight-Bearing Asymmetry During Sit-Stand Transitions Related to Impairment and Functional Mobility After Total Knee Arthroplasty	functional performance	12-steps	ascend, turn around, and descend a flight of stairs. 2 trials. There was a handrail on the staircase, which participants were encouraged to not use during the test.	-time
Sattler F et al. 2011 RCT, n= 112 [120]	Testosterone Threshold Levels and Lean Tissue Mass Targets Needed to Enhance Skeletal Muscle Strength and Function	muscle performance and physical function	12 steps	ascend 12 steps of stairs, time to ascend the middle four steps in the 12-step staircase was measured. photocells used to measure time for stair climbing power	-time to ascend the middle 4 steps in 12-step staircase -stair climbing power
Travison TG et al. 2011 RCT, n=165 [121]	Clinical Meaningfulness of the Changes in Muscle Performance and Physical Function Associated with Testosterone Administration in Older Men With Mobility Limitation	physical function	12-step staircase	ascend 12 steps with and without weights equal to 20% of their body weight. Two trials were required.	-time to ascend -stair-climbing power (product of body weight plus weight carried, total stair-rise, divided by ascent time)

Bade MJ et al. 2012 CT, n=118 [46]	Predicting Poor Physical Performance after Total Knee Arthroplasty	functional performance	12 steps steps height 18 cm depth 28 cm	ascend, turn around, and then descend the steps as quickly as possible in a safe manner. Bilateral handrails were available for use if needed, 2 tests	-time
Heiberg CE et al. 2012 RCT, n= 68 [16]	Effect of a Walking Skill Training Program in Patients Who Have Undergone Total Hip Arthroplasty	physical functioning	8 steps, step height: 16 cm	ascend and descend 8 steps as fast as possible without running, instructed to use alternate legs, using the stair rail was allowed.	-time to ascend
Hsieh RL et al. 2012 RCT, n= 72 [213]	Short-Term Effects of 890-Nanometer Radiation on Pain, Physical Activity, and Postural Stability in Patients with Knee Osteoarthritis	physical activity	14 steps, step height: 18 cm	ascend and descend a flight of stairs as fast as possible.	-total time
van de Port IGL et al. 2012 RCT, n= 250 [103]	Effects of circuit training as alternative to usual physiotherapy after stroke	functional mobility	5 steps chair placed 0,5 meters from the stairs	modified SCT: combination of the timed up and go test before ascending and descending 5 steps of stairs and then sitting down again. The required time from elevating from the chair to sitting again after the stair climb was measured.	-total time
Alfano LN et al. 2013 CT, n=25 [89]	Correlation of knee strength to functional outcomes in Becker muscular dystrophy	functional performance	4 steps step height: 6 inches = 15,24cm	ascend and descend the stairs as quickly and safely as possible. The use of handrails or other compensatory movement patterns was permitted.	-time
Baert IAC et al. 2013 CT, n=87 [78]	Weak associations between structural changes on MRI and symptoms, function and muscle strength in relation to knee osteoarthritis	physical function	5 steps	ascend, turn around and descend 3 tests, mean value	-time
Baert IAC et al. 2013 CT, n=45+20 [77]	Proprioceptive accuracy in women with early and established knee osteoarthritis and its relation to functional ability, postural control, and muscle strength	physical function	5 steps step height 15 cm	ascend, turn around, and descend five steps. mean value of 3 trials	-time
Basaria A et al. 2013 RCT, n= 76 [162]	The Safety, Pharmacokinetics, and Effects of LGD-4033, a Novel Nonsteroidal Oral, Selective Androgen Receptor Modulator, in Healthy Young Men	physical performance (in terms of safety, tolerability)	12-steps, step height: 17 cm	ascend 12 steps as fast as possible. One test round, then two trials	-total time (best out of two) - SCT- Power (calculated from the time elapsed, body weight, and vertical distance)

Chalé A et al. 2013 RCT, n= 80 [153]	Efficacy of Whey Protein Supplementation on Resistance Exercise-Induced Changes in Lean Mass, Muscle Strength, and Physical Function in Mobility-Limited Older Adults	physical function	10 steps	ascend a 10-rise set of stairs as fast as possible, without hold on to the railing or use assistive devices. They did two trials; average time was recorded.	-average time to ascend
Chung JY et al. 2013 CT, n= 24 [29]	Is bicompartamental knee arthroplasty more favourable to knee muscle strength and physical performance compared to total knee arthroplasty	physical performance	12 steps, step height: 20 cm step depth: 30 cm	ascend and descend 12 steps as fast as possible while still staying safe. Using the handrail was allowed. Two repetitions.	-total time (average of two trials)
van Leeuwen DM et al. 2013 CT, n=22 [30]	Preoperative Strength Training for Elderly Patients Awaiting Total Knee Arthroplasty	physical function	9 steps	ascend, turn around, and descend "walk as quickly and safely" allowed to use the handrail and instructed to	-time
Vincent HK et al. 2013 CT, n =53 [57]	"Functional Pain," Functional Outcomes, and Quality of Life After Hyaluronic Acid Intra-articular Injection for Knee Osteoarthritis	physical function	1 flight, 12 steps	climb up 12 steps as fast as possible. Repeated 3 times, fastest trial time was used for data analysis.	-time to ascend
Akbaba YA et al. 2014 RCT, n=20+20+20 [31]	Intensive supervision of rehabilitation programme improves balance and functionality in the short term after bilateral total knee arthroplasty	physical function	10 steps step height 19 cm depth 27 cm	climb up and down the stairway, as fast as possible	-time
Bieler L et al. 2014 RCT, n= 122 [63]	Intra-rater reliability and agreement of muscle strength, power, and functional performance measures in patients with hip osteoarthritis	functional performance	10 steps, step height: 16,3 cm step depth: 35,8 cm	ascend and descend a flight of 10 steps without using the handrail. The best result of 2 timed trials was used noted.	-total time
Hsieh RL et al. 2014 CT, n=40 [75]	Immediate and medium-term effects of custom-moulded insoles on pain, physical function, physical activity, and balance control in patients with knee osteoarthritis	physical function	14 steps step height 18 cm	ascend a flight of stairs as quickly as possible.	-time
Judd DL et al. 2014 CT, n=26+18 [151]	Strength and Functional Deficits in Individuals with Hip Osteoarthritis Compared to Healthy, Older Adults	functional performance	12 stairs	climb a flight of 12 stairs, turn around at the top and descend the same flight as quickly and safely as possible. They were permitted to use the handrail for balance but were instructed not to use the handrail to push or pull	-time

Marmon AR et al. 2014 CT, n=24 [47]	Associations between knee extensor power and functional performance in patients after total knee arthroplasty and normal controls without knee pain	Functional performance	12 steps	ascend and descend 12 stairs	-time
Marmon AR et al. 2014 CT, n=84+68 [79]	Perception and Presentation of Function in Patients with Unilateral Versus Bilateral Knee Osteoarthritis	functional ability	12 stairs	ascend and then descend a flight of 12 stairs as quickly and safely as possible. use of the handrail was allowed if necessary.	-time
Tsukagoshi R et al. 2014 RCT, n= 65 [39]	Functional performance of female patients more than 6 months after total hip arthroplasty shows greater improvement with weight-bearing exercise than with non-weight-bearing exercise	functional performance	10 steps step height: 17,5 cm	ascend 10 steps of stairs.	-time to ascend
Winters JD et al. 2014 restro. study, n=35+23 [38]	Preliminary Investigation of Rate of Torque Development Deficits Following Total Knee Arthroplasty	physical function	12 steps height 18 cm depth 28 cm.	ascend, turn around, and then descend the steps as quickly as possible in a safe manner. The handrail was available for use if needed. 2tests	-time
Zeni, Jr. J et al. 2014 CT, n=56 [80]	Relationship between strength, pain, and different measures of functional ability in patients with end-stage hip osteoarthritis	physical function	12 standard steps	ascend and descend 12 standard steps. A handrail is available during testing	-time
Zhang M et al. 2014 CT, n=72 [159]	Relation Between Anxiety, Depression and Physical Activity and Performance in Maintenance Hemodialysis Patients	physical performance	22 steps	climb the stairs as fast as possible without running, jumping or skipping steps, and were allowed to use the banister for balance if necessary	-time
Altubasi IM et al. 2015 CT, n=21 [134]	Is quadriceps muscle strength a determinant of the physical function of the elderly?	physical function	1 flight 11 steps step height 17 cm	climb up as fast as possible. 1 practice round 2 trials.	-time to ascend
Becker C et al. 2015 CT, n=201 [146]	Myostatin antibody (LY2495655) in older weak fallers: a proof-of-concept	physical performance	4 steps 12 steps step height: 15-18 cm	climb up 4 and 12 steps of stairs while using the handrail.	-time to ascend
Brenneman EC et al. 2015 CT, n=38 [58]	A Yoga Strengthening Program Designed to Minimize the Knee Adduction Moment for Women with Knee Osteoarthritis	mobility	9 steps	ascend and descend a 9-step staircase as fast and safely as possible without running and without skipping stairs. Using the handrail was allowed. Two trials were completed.	-mean time to ascend -mean time to descend

Chikani V et al. 2015 CT, n=13+13 [123]	Impairment of Anaerobic Capacity in Adults With Growth Hormone Deficiency	physical function leg muscle power	4 flights 48 steps step height 17 cm	ascend four flights as fast as possible, one step at a time. 3 tests separated by a rest period of 5 minutes.	-time
Dias CP et al. 2015 RCT, n= 26 [135]	Effects of eccentric-focused and conventional resistance training on strength and functional capacity of older adults	functional capacity	8 steps step height: 17 cm step length: 31 cm	climb eight steps without using the handrail. Two trials with 3 min of rest between	-fastest performance time
Harries N et al. 2015 CT, n= 43 [102]	A stair-climbing test for measuring mechanical efficiency of ambulation in adults with chronic stroke.	motor performance	4 steps, step height: adjustable 10 to 20 cm	climb up and down the stairs for 5 min, self-pacing their speed, keeping it constant, using the handrail for assistance if needed. They did two repetitions of the test.	-time
Pirotta S et al. 2015 RCT, n= 26 [136]	Effects of vitamin D supplementation on neuroplasticity in older adults	physical function	10 steps, step height: 7,8 cm	ascend 10 steps of stairs as fast as possible without using the handrail or walking aids.	-time to ascend -stair climbing power
Akbaba YA et al. 2016 RCT, n= 60 [31]	Intensive supervision of rehabilitation programme improves balance and functionality in the short term after bilateral total knee arthroplasty	physical functionality	10 steps, step height: 19cm step depth: 27 cm	climb up and down ten steps, as fast as possible. Two trials	-total time (mean value of 2 trials)
Bittel TC et al. 2016 CT, n=79 [109]	Adipose tissue content, muscle performance and physical function in obese adults with type 2 diabetes mellitus and peripheral neuropathy	Physical function	1 flight 10 steps	climbing a flight of stairs with 10 steps	-time -power (calculated from the time taken to ascend the stairs, the known vertical height, gravitational force and body mass)
Collado-Mateo D et al. 2016 CT, n = 20 [90]	Performance of women with fibromyalgia in walking up stairs while carrying a load	step-by- step-performance and trunk tilt	10 steps step height: 17 cm depth: 28 cm	climb 10 stairs without carrying a load, rest for 3 min, repeat the test carrying a load of 5 kg in each hand. sensors for motion capture were used.	-kinematic data
Heiberg KE et al. 2016 RCT, n= 60 [42]	Physical Functioning and Prediction of Physical Activity After Total Hip Arthroplasty: Five-Year Follow-up	physical function	8 steps step height: 16 cm	ascend and descend 8 steps as fast as possible without running, allowed to use the handrail, but not a walking aid.	-total time
Hsieh RL et al. 2016 RCT, n=90 [76]	Clinical effects of lateral wedge arch support insoles in knee osteoarthritis	physical activity	1 flight 14 steps step height: 18 cm	ascend and descend 14 steps in the shortest time possible.	-total time

Reddy S et al. 2016 CT, n=264 [161]	Timed Stair Climbing is the Single Strongest Predictor of Perioperative Complications in Patients Undergoing Abdominal Surgery	physical performance	7 steps	walk down and then up one single flight. Vital signs were collected prior to beginning and immediately after completing the task.	-time
Bade M et al. 2017 RCT, n=162 [45]	Early High-Intensity Versus Low-Intensity Rehabilitation after Total Knee Arthroplasty	physical function	1 flight, 12 stairs, step height: 17,1 cm	climb up and down 12 flights of stairs	-total time
Baldwin JN et al. 2017 CT, n=1000 [164]	Reference values and factors associated with musculoskeletal symptoms in healthy adolescents and adults	stair-climbing ability	flight of stairs	timed up and Down Stairs Test by Zaino et al	-time
Baldwin JN et al. 2017 CT, n=1000 [163]	Relationship between physical performance and self-reported function in healthy individuals across the lifespan	stair-climbing ability	flight of stairs	Timed up and Down Stairs Test by Zaino et al	-time
Bieler T et al. 2017 RCT, n= 152 [64]	In hip osteoarthritis, Nordic Walking is superior to strength training and home-based exercise for improving function	functional performance	10 steps, step height: 16,3 cm step depth: 35,8 cm	ascend and descend a flight of 10 steps without using the handrail as fast as possible.	-total time (best out of two trials)
Freisinger GM et al. 2017 CT, n=33 [65]	Relationships Between Varus-Valgus Laxity of the Severely Osteoarthritic Knee and Gait, Instability, Clinical Performance, and Function	physical function	12 steps	ascend and descend a staircase as quickly as possible in a safe manner. Encouraged not to use the handrail unless necessary to complete the test.	-time
Johnen B et al. 2017 Pilot study, n=45 [137]	Feasibility of a machine vs free weight strength training program and its effects on physical performance in nursing home residents: a pilot study	physical performance	11 steps	climb 11 risers of stairs as fast as possible)	-time
Loyd BJ et al. 2017 retro. study, n=162 [32]	Influence of Hip Abductor Strength on Functional Outcomes Before and After Total Knee Arthroplasty: Post Hoc Analysis of a Randomized Controlled Trial	physical performance	12 steps	ascend and descend a set of 12 steps with or without the use of a handrail for balance. This task was timed and the faster of the 2 trials used for analysis.	-time
Maffiuletti NA et al. 2017 CT, n=40 [110]	Reproducibility of clinician-friendly physical performance measures in individuals with obesity	physical function	13 stairs height 15 cm depth 32 cm	referred to Perron et al. stand up from a chair; walk 3 m, ascend at a comfortable pace, turn around and descend stairs, walk back to the chair, turn and sit down.	-time

Nilsen TS et al. 2017 CT, n=58 [117]	Effects of strength training on body composition, physical functioning, and quality of life in prostate cancer patients during androgen deprivation therapy	physical function	stairs	SCT referred to literature with and without additional 20kg	-time
Romine PE et al. 2017 CT, n=430 [138]	Task-Specific Fatigue Among Older Primary Care Patients	physical function	2 m high flight	climbing a flight of stairs twice, using a handrail and/or cane if needed. 2tests, average used	-time
Sions JM et al. 2017 CT, n=106 [139]	Multifidi Muscle Characteristics and Physical Function among Older Adults with and without Chronic Low Back Pain	physical function	2 steps depth: 28cm, height: 17cm	fast stair descent, "as quickly and as safely as possible". The average of two trials was calculated	-time
Storer TW et al. 2017 RCT, n= 308 [118]	Effects of Testosterone Supplementation for 3 years on Muscle Performance and Physical Function in Older Men	physical function	12 steps	four repetitions of ascending the stairs as fast as possible without running, first two tests unloaded, second two test whilst carrying weights equivalent to 20 % of their body weight.	-time to ascend -stair-climb Power (=product of the total rise of the 12 steps, body weight plus load carried, and acceleration of gravity, all divided by time)
Suh MJ et al. 2017 CT, n=34 [49]	Effects of Early Combined Eccentric-Concentric Versus Concentric Resistance Training Following Total Knee Arthroplasty	physical function	1 flight 12 steps step height: 17cm step wide: 25 cm	ascend and descend a flight of stairs as fast as possible on the word "go". 5-minute rest interval between tests, the best score was recorded	-time
Gagliano-Juca T et al. 2018 RCT, n=99 [119]	Testosterone does not affect agrin cleavage in mobility-limited older men despite improvement in physical function	physical function	12 steps	climb up and down 12 steps of stairs as fast as possible without running while carrying weight equal to 20 % of their body weight.	-total time -stair-climbing power (=product of body weight plus weight carried, total stair-rise, and acceleration of gravity all divided by ascent time)
Mulla DM et al. 2018 RCT, n= 43 [165]	The Effects of Lower Extremity Strengthening Delivered in the Workplace on Physical Function and Work-Related Outcomes Among Desk-Based Workers	mobility	1 flight 11 steps	ascend and descend a flight of stairs with 11 steps as fast as possible without running and without skipping steps. It was permitted to use the handrail. Two trials were completed.	-average time to ascend -average time to descend
Mustafaoğlu R et al. 2018 Pilot study, n=45 [95]	The effects of body weight-supported treadmill training on static and dynamic balance in stroke patients: A pilot, single-blind, randomized trial	physical function	10 steps step height 17 cm	ascending and descending using handrails or assistive devices were allowed, if necessary	-time

Iijima H et al. 2019 CT, n=57 [69]	Stair climbing ability in patients with early knee osteoarthritis: Defining the T clinical hallmarks of early disease	physical function	11 steps step height 17 cm step width 135 cm step tread 29 cm	wearing the standardized shoes with pressure sensor-mounted insoles, descend and ascend the flight	-time -vertical ground reaction force (GRF) was calculated from sensor-mounted shoe data -power of 11-SCT for identifying early knee OA (Kellgren and Lawrence grade 1)
Judd DL et al. 2019 CT, n=79 [52]	Trajectories of functional performance and muscle strength recovery differ after total knee and total hip replacement: A performance-based, longitudinal study.	physical function	12 stairs	ascend and descend 12 stairs	-time
Kim JH et al. 2019 Retro. Study, n=184 [48]	Functional Outcomes After Critical Pathway for Inpatient Rehabilitation of Total Knee Arthroplasty	functional mobility lower extremity muscles forces	1 flight 12 steps step height: 17 cm step wide: 25 cm	ascend and descend a flight of stairs	-time
Lange-Maia BS et al. 2019 CT, n=829 [140]	Factors influencing longitudinal stair climb performance from midlife to early late life: The Study of Women's Health Across the Nation Chicago and Michigan Sites	physical function	4 standard stairs	ascend and descend the stairs for three consecutive cycles. Using the handrail was allowed if needed.	-time
Moukarzel M et al. 2019 RCT, n=24 [33]	The therapeutic role of motor imagery during the chronic phase after total knee arthroplasty	strength and functional mobility	12 steps step height: 18 cm step depth: 28 cm	ascend and descend 12 steps of stairs as fast as possible. One test round and two trials.	-total time
Nunes GS et al. 2019 CT, n = 32 [88]	People with patellofemoral pain have impaired functional performance, that is correlated to hip muscle capacity	physical function	9 steps step height: 17 cm	ascend and descend 9 steps as fast as possible step by step. Using the handrail was allowed.	-total time
Orange ST et al. 2019 CCT, n= 36 [141]	Short- term training and detraining effects of supervised vs. unsupervised resistance exercise in aging adults	physical performance	1 free standing flight of stairs with 5 steps, step height 20 cm	ascend and descend as quickly possible while staying safe. Using the handrails was permitted if needed.	-total time

Shimoura K et al. 2019 RCT, n=50 [70]	Immediate Effects of Transcutaneous Electrical Nerve Stimulation on Pain and Physical Performance in Individuals with Preradiographic Knee Osteoarthritis	physical function	11 steps step height: 17 cm	ascend and descend the stairway as fast as possible, started with both feet on the bottom landing and ended in the same position. 2 trials. Using the handrail for support was allowed.	-total time
Sousa-Gonçalves CR et al. 2019 CT, n=8 [111]	Acute Effects of Whole-Body Vibration Alone or in Combination With Maximal Voluntary Contractions on Cardiorespiratory, Musculoskeletal, and Neuromotor Fitness in Obese Male Adolescents	Musculoskeletal and neuromotor fitness	13 steps step height: 15.3 cm total height: 1.99 m	climb up ordinary stairs at the highest possible speed, according to their capabilities	-time -power: (calculated from the time taken to ascend the stairs, the known vertical height, and body mass)
Suh et al. 2019 CT, n= 195 [74]	Bilateral Quadriceps Muscle Strength and Pain Correlate With Gait Speed and Gait Endurance Early After Unilateral Total Knee Arthroplasty	physical function	1 flight 12 steps step height 17 cm wide: 25 cm	ascend and descend a flight of stairs as quickly as possible. 3trial with five minutes break in between.	-total time (best out of three)
Unhjem R et al. 2019 CT, n=41 [142]	Functional Performance With Age: The Role of Long-Term Strength Training	functional performance	12 steps step height: 17 cm	climbing as fast as possible. Maximum step length was set to 2 stair steps at a time, no handrail support was allowed.	-power (W)
Bade MJ et al. 2010 CCT, n=24 [44]	Outcomes Before and After Total Knee Arthroplasty Compared to Healthy Adults	functional performance	10 steps step height 17.1-cm 12-steps step height 17.1-cm	9 patients were tested on a 10-step staircase, 15 patients and all healthy adults were tested on a 12-step staircase	-time
Daly RM et al. 2020, RCT, n= 216 [154]	Effects of a multivitamin-fortified milk drink combined with exercise on functional performance, muscle strength etc. in middle-aged women	functional muscle power	10 stairs, step height: 17 cm	ascend or descend a flight of 10 stairs as fast as possible without missing a step. Using the handrail if necessary was allowed. They did 1 practice trial and then 3 test trails.	-time to ascend -time to descend -stair climb power on ascend (power (Watts) = $9.81 \times \text{body mass (kg)} \times \text{vertical step height (m)} \times \text{number of steps / time (s)}$)
Gränicher P et al. 2020 Pilot RCT, n=20 [34]	Preoperative exercise in patients undergoing total knee arthroplasty: a pilot randomized controlled trial	functional performance	8 steps height 16 cm depth 30 cm	ascend and descend a flight at usual walking speed, feeling safe and comfortable. handrail or assistive devices were allowed but not encouraged	-time
Larsen JB et al. 2020 Retro. Study, n=217 [40]	Intensive, personalized multimodal rehabilitation in patients with primary or revision total knee arthroplasty: a retrospective cohort study	lower body strength balance	22 stairs	ascend and descend stairs as quickly and as safely as possible. Use of a handrail and walking aid were permitted if needed.	-time

Lee SJ et al. 2020 CT, n=84 [51]	Preoperative physical factors that predict stair-climbing ability at one month after total knee arthroplasty	physical function	12 steps step height 17 cm 25 cm wide	ascend or descend the stairs as quickly as possible on the word "go". 3 trial with a 5-min rest interval Using a handrail was allowed	-time
Mustafaoglu R et al. 2020 RCT, n= 51 [96]	Does robot-assisted gait training improve mobility, activities of daily living and quality of life in stroke	mobility	10 steps step height: 18 cm	climb up and down ten steps without skipping any steps, using one foot for each step and descend without stopping. Use of handrail and/or assistive devices was allowed.	-total time
Onodera CMK et al. 2020 N=153 [60]	The importance of objectively measuring functional tests in complement to self-report assessments in patients with knee osteoarthritis	physical function	9 steps and 9 landings step height 18 cm	reach up and down the stairs	-time
Pozzi F et al. 2020 RCT, n=293 [37]	Restoring physical function after knee replacement: a cross-sectional comparison of progressive strengthening vs standard physical therapy	physical function	12 stairs, (15cm rise, 20cm run)	ascend and descend the set of stairs as fast as possible without skipping steps. Using the handrail was allowed. Time started on command and ended when both feet touched the bottom again.	-total time
Rigamonti AE et al. 2020 CT, n=595 [113]	Impact of a Three-Week in-Hospital Multidisciplinary Body Weight Reduction Program on Body Composition, Muscle Performance and Fatigue in a Pediatric Obese Population with or without Metabolic Syndrome	muscle performance	13 steps step height: 15.3 cm total height: 1.99 m	climb up ordinary stairs at the highest possible speed. 2–3 test trials.	-time
Rigamonti AE et al. 2020 CT, n= 1922 [214]	Effects of a 3-Week In-Hospital Body Weight Reduction Program on Cardiovascular Risk Factors, Muscle Performance, and Fatigue: A Retrospective Study in a Population of Obese Adults with or without Metabolic Syndrome	muscle function	13 steps step height 15.3 cm total height 1.99 m	climb up ordinary stairs at the highest possible speed	-time

Tamini S et al. 2020 CT, n=16 [112]	Acute Effects of Whole-Body Vibration Exercises at 2 Different Frequencies Versus an Aerobic Exercise on Some Cardiovascular, Neuro-motor and Musculo-skeletal Parameters in Adult Patients With Obesity	maximal lower limb muscle power motor control	13 steps step height: 15.3 cm total height: 1.99 m	climb up ordinary stairs at their highest speed, in accordance to their capabilities.	-time -power (=product of 1/4patient's weight, gravitational force, and vertical height /time)
Vongsirinavarat M et al. 2020 CT, n=250 [59]	Identification of knee osteoarthritis disability phenotypes regarding activity limitation: a cluster analysis	physical function	Not specified	Timed stair climbing referred to literature	-time
Ahmed Burq HSI et al. 2021, RCT, n= 64 [97]	Effect of whole-body vibration on obstacle clearance and stair negotiation time in chronic stroke patients	mobility function	3 steps step height: 18 cm step depth: 30 cm	ascend and descend 3 steps of stairs in a comfortable speed, accompanied by a therapist. The handrails were used.	-total time
Beckmann M et al. 2021 CT, n=207 [81]	Recovery and prediction of physical function 1 year following hip fracture	physical function	8 steps	ascended and descended 8steps as fast as able without running, using alternate legs and support by stair rail if needed.	-time
Choi JH et al. 2021 CT, n=149 [36]	Performance-based physical function correlates with walking speed and distance at 3 months post unilateral total knee arthroplasty	physical function	12 steps height 17 cm wide 25 cm	ascend and descend a flight of 12 steps, as fast as possible on the word "go". 3trials, with the 5 min rest in between	-time
Jacksteit R et al. 2021 RCT, n=85 [35]	Low-Load Unilateral and Bilateral Resistance Training to Restore Lower Limb Function in the Early Rehabilitation After Total Knee Arthroplasty: A Randomized Active-Controlled Clinical Trial	physical performance postural control strength of the lower extremities	8 steps step height: 17.5 cm	climb a staircase in a safely and quickly manner using a railing and regular footwear.	-time
Katsoulis K et al. 2021 Retro. study, n=18 [143]	Reliability of Lower Extremity Muscle Power and Functional Performance in Healthy, Older Women	physical function	13 steps step height 18 cm step width 28 cm height 2.34 m	stand at the base of the stairs with feet together and to grab the handrail if necessary during ascent. Upon the instruction "ready, set, go," stair ascent was measured during a "usual" pace (SCUP) and during a "fast" pace (SCFP) with instructions to ascend "as quickly and as safely as possible."	-time -power = ((body mass in kg) × (9.8 m/s ²) × (stair height in meter))/(time in seconds)

Kim BS et al. 2021 CT, n=562 [73]	Associations Between Obesity With Low Muscle Mass and Physical Function in Patients With End-Stage Knee Osteoarthritis	Physical function	1 flight 12 steps step height: 17 cm step length: 25 cm	ascend or descend the stairs as fast as possible upon hearing the word "go". 3 trials, with a 5-minute rest interval between. The fastest time was recorded for each patient	-ascend time -descend time
Khruakhorn S et al. 2021 RCT, n=34 [61]	Effects of hydrotherapy and land-based exercise on mobility and quality of life in patients with knee osteoarthritis: a randomized control trial	physical function	4 steps step width: 26.5 cm step length: 76 cm step height: 15.2 cm	"Walk up – turn back – and go down the stairs as soon as possible but safely" and subsequently started the assessment with the word "Start". 2 trials with a 5-minute rest between were evaluated.	-time
Bayartai ME et al. 2022 CT, n=160 [108]	Changes in the Oswestry Disability Index after a 3-Week In-Patient Multidisciplinary Body Weight Reduction Program in Adults with Obesity	lower limb muscle power functional ability	13 steps step height: 15.3 cm total height: 1.99 m	climb a staircase at maximum speed. 1-2 test trials	-time
De Zwart AH et al. 2022 CT, n=177 [62]	Association Between Measures of Muscle Strength and Performance of Daily Activities in Patients with Knee Osteoarthritis	functional performance	12 steps	ascending and descending (independently) a staircase as fast as possible without running	-time
Ihl T et al. 2022 RCT, n=1771 [98]	Patient-Centered Outcomes in a Randomized Trial Investigating a Multimodal Prevention Program After Transient Ischemic Attack or Minor Stroke: The INSPIRE-TMS Trial	physical activity	first landing of a stairway	ascend as quickly as possible to the first landing of a stairway. Patients were allowed to use the banisters, if necessary	-power (power=body weight×9.8×height of the staircase/mean time of 2 runs)
Ke D et al. 2022 CT, n = 103 [155]	Study of the Reliability of Field Test Methods for Physical Fitness in Children Aged 2–3 Years	physical fitness coordination ability	5 steps step height 12 cm	climb up the stairs with using both feet alternately and without using the handrail. 2 trials.	-time to ascend (best out of two)
Lee SH et al. 2022 CCT, n=30+30 [72]	Validity of the Osteoarthritis Research Society International (OARSI) recommended performance-based tests of physical function in individuals with symptomatic Kellgren and Lawrence grade 0–2 knee osteoarthritis	physical function	9 stairs step height 20 cm	ascended and descended as quickly as possible but in a safe manner.	-time
Lindberg K et al. 2022, RCT, n= 49 [144]	Effectiveness of individualized training based on force– velocity profiling on physical function in older men	physical function	15-steps, Step height: 16 cm	climb up 15 steps of stairs as fast as possible with and without a weight vest of 20 kg.	-time to ascend

Lazzer S et al. 2022 CT, n=139 [114]	Effects of a 3-Week Inpatient Multidisciplinary Body Weight Reduction Program on Body Composition and Physical Capabilities in Adolescents and Adults With Obesity	muscle function	13 steps step height 15.3 cm total height 1.99 m	climb an ordinary stair at the highest possible speed	-time
Petersson N et al. 2022 CT, n=14 [66]	Blood Flow Restricted Walking in Elderly Individuals with Knee Osteoarthritis: A Feasibility Study	Functional performance	11 steps	11step stair climb test referring to another trial	-time
Usubini AG et al. 2022 CT, n=237 [115]	A three-week in-hospital multidisciplinary body weight reduction program exerts beneficial effects on physical and mental health and fatiguability of elderly patients with obesity	muscle function	13 steps step height 15.3 cm total height 1.99 m	climb up ordinary stairs at the highest possible speed	-time -power (kg × 9.81 × 1.99) /s
Carvalho C et al. 2023 CT, n=39 [68]	Association between ankle torque and performance-based tests, self-reported pain, and physical function in patients with knee osteoarthritis	Functional performance	1 flight 11 steps step height: 17 cm step width: 202 cm step tread: 31 cm	go up and down a flight of stairs as quickly and safely as possible.	-time
Fosstveit SH et al. 2023 CT, n=49 [145]	Associations between Power Training-Induced Changes in Body Composition and Physical Function in Older Men: A Pre-Test–Post-Test Experimental Study	Physical function	15 steps Step height: 16 cm	climb 15 steps as fast as possible. The time was recorded using photocells placed at the bottom and top of the stairs at 85 cm height. Warm-up, 2 unloaded tests and 2 with a 20 kg vest	-time
Jankaew A et al. 2023, RCT, n=47 [71]	The effects of low-level laser therapy on muscle strength and functional outcomes in individuals with knee osteoarthritis	functional performance	13 steps, step height: 18 cm	climb up the stairs, turn around, and climb down as quickly as possible while staying safe.	-total time
Kirschner N et al. 2023 CT, n=24 [50]	Determination of Relationships between Symmetry-Based, Performance-Based, and Functional Outcome Measures in Patients Undergoing Total Hip Arthroplasty	physical performance	14 steps	14-step stair-climbing test Preoperatively with no walking aids. Post-operatively, walking support and the stair railing were used.	-time
Lanzi S et al. 2023 CT, n=90 [156]	Time-course evolution of functional performance during a 3-month supervised exercise training program in patients with symptomatic peripheral artery disease	functional performance	12 stairs	climb as quickly as possible using a handrail was allowed. 2 tests	-time

Suslov VM et al. 2023 CT, n=28 [92]	Efficacy and safety of hydrokinesitherapy in patients with dystrophinopathy	physical function	4 steps	descent 4-stairs	-time
---	---	-------------------	---------	------------------	-------

Table 3: 8 Studies evaluating other aspects.

Author, Study Design, Number of Participants	Study Background	SCT-outcome	SCT-technical background	SCT-implementation	SCT- parameters surveyed
Poppius H et al. 1977 CT, n= 13 [207]	Exercise-induced Asthma and Doxantrazole	to determine if doxantrazole can prevent exercise induced asthma	16 floors of stairs	climb up the stairs in rhythmic pace, given by sounds from a tape recorder. The test ended, when they reached the top of the stairs or when they had to stop because of fatigue or dyspnea.	-pace of climbing -number of flights that induced post-exercise asthma -peak expiratory flow
Aveline C et al. 2014 RCT, n= 69 [208]	Pain and Recovery After Total Knee Arthroplasty-a 12-Month Follow-up	pain after exercise	10 steps height 15 cm	walk up and down 10 stairs	-pain analogue scale (VAS)
Khan A et al. 2015 CT, n= 17 [15]	Musical stairs: the impact of audio feedback during stair-climbing physical therapies for children	impact of audio feedback on the use of reciprocal steps	1 flight of stairs	climb up and down one flight of stairs for 1-5 minutes, with and without audio feedback.	- number of steps
Honda H et al. 2016 CT, n= 16 [202]	Stair climbing/descending exercise for a short time decreases blood glucose levels after a meal in people with type 2 diabetes	postprandial blood glucose in type 2 diabetes	21-step staircase step height: 17 cm	ascend and descend 21 steps 6 times without a break. For the ascend they were instructed to climb at a pace of 80-110 steps per minute, for the descend they chose their pace freely. The test was conducted at 60 minutes after meal and 120 minutes after meal.	-heart rate during test -Borg scale -glucose, lactate, C peptide, and non-esterified fatty acid
Takaishi T et al. 2017 CT, n= 14 [203]	Stair ascending-descending exercise accelerates the decrease in postprandial hyperglycemia more efficiently than bicycle exercise	effect of Stair climbing-descending exercise on lowering postprandial hyperglycemia	1 flight 21 steps step height 18 cm	8-10 repetitions of walking down and up one flight of stairs at an intensity of high-moderate to low-vigorous intensity	-heart rate -blood glucose
Bartholomae E et al. 2018 CT, n=30 [205]	Reducing Glycemic Indicators with Moderate Intensity Stepping of Varied, Short Durations in People with Pre-Diabetes	Blood sugar reduction	21 steps	ascended and descended a stairwell continuously	OGTT
Moore J et al. 2020 RCT, n=30 [204]	A single one-minute, comfortable paced, stair-climbing bout reduces postprandial glucose following a mixed meal	Metabolism/Blood sugar	21 steps	0, 1, 3, and 10 min SCD bouts performed at self-selected pace stairwell was ascended and descended in a continuous fashion.	Blood sugar
Matsumoto K et al. 2022 RCT, n=22 [206]	The Effect of Brief Stair-Climbing on Divergent and Convergent Thinking	Divergent and convergent thinking	4 flights 21 stairs	walk downstairs and back after approaching	Ability to think

Moreover, the SCT was performed in various ways. In some trials the test was conducted with 10 steps or more. Most studies evaluated the ascending as well as the descending, others either or. The measured parameters differed a lot as well. With a total of 122 studies, the majority used the time needed to complete the SCT as the only recorded result, the total distance however varied. In only a few trials, other parameters were collected, including vital signs or the number of steps climbed. Several studies calculated the so-called stair-climbing power or VO₂max of the collected values.

Some trials even showed, that the SCT can be a part of physiotherapy. Khan et al. for instance integrated stair climbing with and without audio feedback into the physiotherapy of 17 children with different diagnoses and then investigated whether the use of audio feedback made a difference to the outcome during SCT. [15] Overall, there is no standardization of the SCT and therefore its clinical use is not clearly recorded. The SCT has been used in completely different ways to evaluate different body functions. The implementation varied in the number of steps that patients had to climb, the parameters collected, the questions to be answered with the SCT, permitted aids during the test and the information that patients received before the test. All these aspects varied significantly and were not standardized.

Outcome

Physical and neuromuscular function (and mobility)

Physical function is one of the commonly examined parameters when using the SCT. 155 of the included studies used the SCT to evaluate physical function of participants or patients. (table1)

a) Patients with orthopedic- or muscular conditions

The examined participants often had an orthopedic intervention and were in various stages of rehabilitation. Heiberg et al. for instance evaluated the effect of a walking skill training program in 68 patients who had undergone a total hip arthroplasty and used the SCT for measuring physical function 3 months post operation. [16] Other working groups also used a form of SCT to evaluate physical or motor function in patients with knee or hip arthroplasty, either to compare capacity before and after surgery, others to evaluate a special treatment. [17-51] Judd et al. compared patients with knee and hip replacement. They showed that functional performance and muscle strength recovery differ after total knee and total hip replacement. [52]

Another patient group in which a form of the SCT was applied repetitively are patients with knee or hip osteoarthritis. The SCT in these patients was also used to evaluate different treatment approaches for instance Kovacs et al. evaluated the effect of thermal water in patients with osteoarthritis. [53] Kraemer et al. examined the effect of a treatment combination in patients with osteoarthritis [54], whereas Laufer et al. assessed the effect of pulsed short-wave diathermy on pain and function of subjects with osteoarthritis of the knee. [55] Other studies also used the SCT in pa-

tients with osteoarthritis to evaluate physical or motor function. [27-56]

Other orthopedic patients were also analyzed using a SCT. For example, Beckman et al. assessed physical function in patients after hip fracture. [81] Moreover Adunsky et al. evaluated MK-0677 (ibutamoren mesylate) for the treatment of patients recovering from hip fracture with a SCT. [82] Ljungquist et al. evaluated physical performance in patients with spinal pain. [83] Smeets et al. used the SCT to test physical capacity tasks in chronic low back pain patients. [84,85] assessed physical function in patients with rheumatoid arthritis and ankylosing spondylitis. [86] Wetzel et al. used the SCT to examine functional lower-extremity strength power in patients with multiple sclerosis. [87] Nunes et al. assessed people with patellofemoral pain regarding their functional performance, in correlation to hip muscle capacity. [88] Alfano et al. used the SCT to evaluate the correlation of knee strength to functional outcomes in Becker muscular dystrophy. [89] Collado-Matteo et al. used the SCT to assess the performance of women with fibromyalgia. [90] Schwid et al. assessed the effect of sustained release of 4=aminopyridine on motor function as a symptomatic treatment of multiple sclerosis. [91] Suslov et al. evaluated the efficacy and safety of hydrokinesiotherapy in patients with dystrophinopathy. [92]

b) Follow-up after Cerebrovascular accidents

Apart from orthopedic and surgical patients the SCT was also used to evaluate physical and motoric function in neuromuscular patients as for instance patients after stroke. Katz-Leurer et al. examined neurological patients with cerebrovascular accident, using a SCT. [93] used the SCT in patients after stroke to assess physical function, Mustafaoglu et al. assessed the effects of body weight-supported treadmill training on static and dynamic balance in stroke patients. [94,95] In another study they also evaluated the mobility of stroke patients using a SCT. [96] Ahmed Burq et al. assessed the effect of whole-body vibration on obstacle clearance and stair negotiation time in chronic stroke patients. [97] evaluated a multimodal prevention program after transient ischemic attack or minor stroke. [98] Other working groups also used the SCT to investigate patients after stroke for instance the effect on physical performance of strength- or other training approaches. [99-102] Van de Port et al. examined the effect of circuit training compared with usual physiotherapy in 126 patients with stroke. The SCT combined with a stand-up and go test was utilized to evaluate functional mobility. [103]

Furthermore, the SCT was also used to examine children with cerebral palsy (CP). Zaino et al. used a timed SCT to assess 47 children aged 8-14 years, 27 of them with CP. [104] Bar-Haim et al. also assessed children with CP with the SCT. [105]

c) Obesity and weight reduction

Apart from the approaches mentioned above, the SCT was also used repeatedly to assess obese individuals, for example

after weight reduction. Sartorio et al. assessed the effect after a short-term body mass reduction program in obese subjects on their motor function. [106] had a similar approach and assessed the effect of exercise training in obese women. [107] evaluated the changes in the Oswestry Disability Index after a 3-week in-patient multidisciplinary body weight reduction program in adults with obesity. [108] Bittel et al. assessed physical function in obese adults with type 2 diabetes mellitus and peripheral neuropathy, using a SCT. [109] Maffiuletti et al. examined the reproducibility of clinician-friendly physical performance measures in individuals with obesity using a SCT. [110] Sousa-Goncalves et al. evaluated the acute effects of whole-body vibration alone or in combination with maximal voluntary contractions on cardiorespiratory, musculoskeletal, and neuromotor fitness in obese male adolescents. [111]

Tamini et al. evaluated the acute effects of whole-body vibration exercises at 2 different frequencies versus an aerobic exercise on some cardiovascular, neuromotor and musculoskeletal parameters in adult patients with obesity. [112] Rigamonti et al. assessed the impact of a three-week in-hospital multidisciplinary body weight reduction program on body composition, muscle performance and fatigue in a pediatric obese population with or without metabolic syndrome. A SCT was used to assess the effect on maximal anaerobic power. The protocol covered a staircase with a total height of 1,99m. The SCT could detect effects of the program. They could show a significant reduction in SCT-time in all groups. Whereas patients with metabolic syndrome profited even more than obese individuals without metabolic syndrome. Younger individuals also profited more than older ones. [113] Lazzer et al. also examined the effect of a 3-week inpatient bodyweight reduction program in obese individuals. The SCT was used to evaluate maximum anaerobic exercise capacity. The protocol also covered a total height of 1,99m. They could show that baseline SCT-time was significantly lower in males, compared to females. After the 3-weeks SCT time decreased significantly solely in females. [114] Usubini et al. used the SCT to assess maximum anaerobic power in obese patients before and after a 3-week multidisciplinary weight reduction program. The SCT protocol used, covered a total height of 1,99m. The test was performed at the fastest possible time and power was calculated according to the following formula: $(\text{body-weight} \times 9.81 \times 1.99) / \text{SCT time}$. They could show a significant reduction in SCT time after the program. [115]

d) Endocrine factors

With gaining interest, the SCT was also used to evaluate the effect of testosterone or androgens on physical function. Knapp et al. examined the effects of a supraphysiological dose of testosterone on physical function, muscle performance, mood, and fatigue in men with HIV-associated weight loss. [116] Nilsen et al. evaluated the effects of strength training on body composition, physical functioning, and quality of life in prostate cancer patients during androgen deprivation therapy. [117] Storer et al. assessed the effects of testosterone supplementation for 3 years on muscle per-

formance and physical function in older men. [118] Gagliano-Juca et al. showed that testosterone does not affect agrin cleavage in mobility-limited older men despite improvement in physical function, using a SCT. [119] assessed the effect of testosterone on physical function, using a SCT. [120,121] used the SCT in patients with prostate cancer to assess the effect of exercise on treatment side effects. [122] evaluated the impairment of anaerobic capacity in adults with growth hormone deficiency. [123]

e) Age and the elderly

Another approach for the SCT was to use it so assess mobility and physical function in otherwise healthy elderly. [118,124,153]. for example investigated the efficacy of whey protein supplementation in 80 mobility-limited older adults, using the SCT among other tests to evaluate physical function. [153] Daly et al. examined the effects of a multivitamin-fortified milk drink combined with exercise on functional performance, muscle strength etc. in middle-aged women. [154] assessed the reliability of a SCT to assess physical function and coordination in children aged 2-3. [155]

f) Patients of internal medicine and general surgery

Lanzi et al. examined functional performance during a 3-month supervised exercise training program in patients with symptomatic peripheral artery disease, using a SCT. [156] used the SCT in patients with hemodialysis to assess the effect of an exercise training on physical function [157-159] Dreher et al. used the SCT in patients with COPD. [160] Reddy et al. used a SCT to predict perioperative complications in patients undergoing abdominal surgery. [161]

g) Healthy adults

Apart from being used in special patient groups, the SCT is also used to assess therapy effects and health preventive approaches in healthy adults. Basaria et al. evaluated the safety, pharmacokinetics, and effects of LGD-4033, a novel nonsteroidal oral, selective androgen receptor modulator, in healthy young men. [162] also examined healthy adults with a SCT regarding musculoskeletal symptoms. [163,164] evaluated the effects of lower extremity strengthening delivered in the workplace on physical function and work-related outcomes among desk-based workers. [165]

Cardiorespiratory fitness / exercise capacity

Cardiorespiratory fitness and exercise tolerance are the second most important function assessed with the SCT. 37 of the included trials examined cardiorespiratory fitness (table 2); It could be outlined that the results in SCT showed a significant correlation to CPET-parameters, maximum heart and respiratory rate e.g. Pollok et al. [166] The participants in the trials, evaluating CRF and exercise capacity, were also orthopedic patients, patients before/after lung transplantation and patients with chronic lung diseases for instance chronic airflow obstruction. For example, Elbasan et al. examined children with cystic fibrosis. [167] More-

over, there were several studies in healthy adults, for instance sedentary young women, or obese females. [168,169] Sartorio et al. examined obese children and adolescents. [170] Devendra et al. examined adults with newly diagnoses persistent foramen ovale with a SCT to assess deoxygenation. [171] Hetzler et al. used a SCT in American football athletes. [172]

a) Pulmonary patient groups

Mc Keon et al. evaluated the effect of inspiratory resistive training on patients with severe chronic airflow limitation. To assess exercise endurance, they used the SCT in addition to other exercise tests to determine the influence of the specific training on different types of exercise. The SCT did not follow a predefined protocol. The number of stairs completed by the patient at a normal rate under the supervision of a physiotherapist were recorded without a time limit and the test ended when symptoms of breathlessness or weakness occurred. [173]

Pollock et al. investigated if a SCT can be used to estimate VO_2max in patients with chronic airway obstruction (CAO). The SCT was conducted as a symptom limited SCT with a maximum of 10 flights, most patients achieved at least 4 flights=13,3m height. The test was conducted twice. A resting gas sample and a gas sample during exercise was collected. VO_2 , CO_2 output, minute ventilation (VE), and tidal volume were calibrated using standard techniques. The subject's blood pressure, pulse, and respiratory rate were measured at rest and immediately after the test. Saturation and heart rate was measured throughout the test. The results in CPET were compared to those achieved in bicycle CPET. The number of steps climbed, correlated well with peak VO_2 . They could show a linear correlation between peak VO_2 for stair climbing and cycle ergometry. Whereas VO_2max , HR, RR, BP were higher during SCT as in CPET. According to their results, the study shows that 1-2 flights of stairs is not exhausting enough. Nevertheless, considering this, the study implies that the SCT can be used to estimate VO_2max . [166] Elbasan et al. assessed the effects of chest physiotherapy and aerobic exercise training on physical fitness in children with cystic fibrosis. 16 patients between 5-13 years were examined. All children were assessed at the beginning and the end of a 6-week training. A 10-step SCT was conducted, as well as bicycle CPET. They could detect a positive effect on anaerobic power and speed, which was reflected in the results of SCT and CPET. [167]

b) Lung resection

Pate et al. assessed patients with non-small lung cancer. In this patient group it is stated to be often difficult to decide whether a surgical resection is possible or not, especially in cases with coexisting chronic airway obstruction or ischemic heart disease. They used a symptom limited protocol for SCT. A stair climb of 3 flights or more was associated with reduced postoperative morbidity and therefore a proof for lobectomy, or 5 flights or more for pneumonectomy. The flight height was not standardized. [174]

The working group Brunelli et al. conducted several studies, using a SCT to assess cardiopulmonary fitness/aerobic capacity and oxygen consumption preoperative in patients considered for surgical lung resection. After a cardiovascular evaluation and a spirometry with assessed carbon monoxide diffusion capacity, a predicted postoperative lung function was calculated. Values below 60% indicated further testing. Therefore, a SCT or 6MWT was used. The SCT protocol covered a total of 16 flights, patients were asked to climb at a pace of their choice and stop for exhaustion or symptoms like dyspnea. An achieved altitude >12m was considered satisfying and surgery was offered. Operative morbidity and mortality using this SCT cut-off values were comparable to the data collected with full CPET. They showed a significant correlation between VO_2max and the climbed altitude. In patients with a predicted postoperative function <30%, CPET was indicated. In another study of this working group, they even stated that a reached altitude of >12m was the only predictor of cardiopulmonary complications. Patients, unable to climb 12m, had a 2.5 to 13fold higher risk for cardiopulmonary complications and mortality, compared to those climbing >22m. An achieved altitude of >18m was significantly correlated with long-term survival. This working group also calculated stair climbing work, VO_2max , oxygen-pulse, using the following formula:

Work = (height of the step-in meters x steps per minute x body weight in kilograms x 0.1635). $\text{VO}_2\text{ max}$ in milliliters per minute = $(5.8 \times \text{body weight in kilograms} + 151 + 10.1 \times \text{work})$. Maximum oxygen pulse = $(\text{VO}_2\text{ max}/\text{maximum heart rate})$. [175-182]

Koegelenberg et al. performed another preoperative study in patients with considered lung resection. The used symptom limited SCT protocol in this study covered a total of 20 m vertical height. Patients were asked to climb as fast as possible. The altitude reached and the average speed was compared to VO_2max during bicycle CPET. They could show that the average climbing speed was an accurate semiquantitative predictor of $\text{VO}_2\text{max}/\text{kg}$. [183] Pancieri et al. used a SCT protocol, covering a total height of 12,16m to examine patients before and after lung resection. As they did before in their working group with Cataneo et al. 40 patients were included. The time needed to complete the stair climb was recorded and the number of functioning lung segments planned for resection, used to predict postoperative test results. The aim was to find an easy exercise test to predict if the patient could tolerate the surgery. [184,185]

Bernasconi et al. compared SCT to treadmill CPET to define cut-off values for lung resection. 56 patients were examined. The SCT protocol included a maximum height of 20m. A portable spirometry was used. The patients reached a mean altitude of 16,9m. 22 patients did not reach 20m. VO_2max differed not significantly between SCT and treadmill CPET. Speed of climbing was significantly associated with $\text{VO}_2\text{ max}$. The group stated that patients climbing 20m with a speed of 15 m/min, are eligible for surgery. [186] Refai et al. investigated the max. inspiratory and expirato-

ry pressure generated in the mouth before and after SCT in 283 patients before lobe- or pneumectomy. 61% of the patients experienced a reduction in PImax. It showed that a predicted loss of >10% was associated with complications. [187] Ito et al. retrospectively analyzed data of 65 old people with non-small cell lung cancer and lobectomy. They used a SCT protocol, covering a total of 18m, 5 flights. They could show that patients without desaturation >4% were eligible for lobectomy. [188]

Kubori et al. compared a SCT with the 6MWT in patients before and after lung resection. The SCT was more sensitive than the 6MWT to detect changes in CRF. The used SCT protocol covered a symptom limited SCT with a maximum of 36 flights. The patients reached a mean altitude of 26.3m before and 18.2 meters 4 weeks after lung resection. The difference in the distance reached in the 6MWT before and after resection was not significant. [189] Nakamura et al. also evaluated patients before and after lung resection, using a SCT. They evaluated if a desaturation of 4% during the SCT could predict complications after lung resection. The symptom limited SCT protocol covered 6 flights with a total height of 22.2m. They could show that patients who underwent the SCT without desaturation had a normal risk for postoperative complications, whereas patients with a desaturation had a higher rate of complications. [190]

Ozeki et al. also evaluated patients with stage I lung cancer before and after lung resection with a SCT to predict whether postoperative exercise capacity will probably be severely reduced or not. The used symptom limited SCT protocol included a total height of 12m. The test was performed at the fastest possible time. Estimated VO2max was calculated, according to Cataneo et al. Patients with complications needed a significant longer time in SCT preoperative. In patients with complications the postoperative exercise capacity was reduced by 4%. [185,191] Dong et al. showed that a SCT could predict postoperative complications in patients with non-small lung cancer. They used a SCT-protocol, covering a maximum height of 18,4m. Desaturation in patients was associated with prognosis. [192]

c) Cardiological and other patient groups of internal medicine

Devendra et al. used a SCT to provoke exercise oxygen desaturation in patients with a persistent foramen ovale. 50 patients were examined. The protocol included climbing and descending 4 flights of stairs. During this procedure oxygen saturation was measured. Provoked exercise desaturation (PED) was defined as a desaturation of at least 8% and a saturation <90%. 17 patients had a PED. 13 of them underwent PFO-disclosure. 10 were followed up. The desaturation improved after closure by a mean of 10%. [171]

Njoeten et al. assessed patients with long Covid. They used a SCT protocol, covering 18 steps, which had to be climbed 3 times up and down as fast as possible. Patients with a normal exercise capacity in CPET were significantly faster as those with reduced

exercise capacity. [193] Hellberg et al. showed that a decline in glomerular filtration rate is associated with a reduced endurance, strength and other symptoms. Endurance was assessed, using a SCT. The SCT covered a total of 12 flights. [194]

d) Patients requiring general surgery

Servio et al. used the SCT to investigate CRF in patients before and after laparoscopic Nissen-funduplicatio. The working group used the SCT protocol of Cataneo et al. It covered a total height of 12,24m, which had to be climbed as fast as possible. Time was recorded and power calculated. Several measurements on different times were collected. They could show that the patients regained their preoperative results shortly (at the 5th postoperative day) after operation. [195] Arruda et al. assessed cardiopulmonary postoperative complications in comparison between upper abdominal and thoracic surgery, using a SCT. [196] Khenafes et al. evaluated CRF in patients before and after cholecystectomy, using a SCT. For the SCT a protocol according to Cataneo et al. was used. A total height of 12,24 meters was covered. They could show a rapid recovery of CRF after surgery. [197]

Cataneo et al. used a standardized protocol in their studies to establish a submaximal exercise test in hospital settings where CPET is not available, to evaluate if the cardiorespiratory system of an individual has the capacity to undergo surgery. They used a SCT protocol, covering 6 flights (12,16m), measured time and calculated the individuals work to climb the stairs. To calculate the work needed to climb the stairs, the body mass, the covered height and gravity was multiplied. The calculated work was divided by the time needed to climb, to get the stair climbing power in Watt. The results were compared to VO2max during CPET. They could show a significant correlation between VO2max and SCT as well as SCT power. As a conclusion they stated that the SCT is a simple cost effective and widely available test to assess an individual's cardiorespiratory capacity, as patients with an impaired cardiac or pulmonary function have difficulty climbing stairs. The working group stated that a SCT should cover a height of at least 12m to get convincing results. [185,198]

e) Normal-weight and overweight healthy study participants

Coll et al. used a modified Chester step test in healthy adults. They used a 20cm step, which had to be stepped on and off on a standardized cadence, which was increased every 2 min. The test was stopped, either symptom limited, or when the individuals reached a heart rate equal to 80% of their maximum HR (220-age). A modification was added in individuals, who reached a cadence of 35 steps/min without reaching any of the mentioned criteria. They were handed 2kg dumbbells. The test was performed twice, and the results were almost equal in both rounds. They found the SCT a reliable test to assess aerobic capacity. [199]

Hetzler et al. developed a football stair climb protocol, including 20 steps with a total height of 3,12m. As football is described

as a highly explosive sport, the group was looking for a test to assess peak anaerobic power. 58 football players were included. They performed 25 trials with 30-40sec rest in between. Time was recorded and power calculated. Power was calculated according to the following formula: $\text{Power} = \text{body mass (kg)} \times 9.81 \text{ ms}^{-2} \times \text{vertical distance} \times \text{time}^{-1}$. The players were divided in 3 groups, according to their position in the game. The skill-group was significantly less powerful. They could show that the test is reliable for measuring peak anaerobic power. To assess the reliability of the test 34 football players repeated the test within a week. [172]

Calavalle et al. used the SCT as a simple method to analyze overall individual physical fitness in firefighters. [200] Boreham et al. performed a „training program“ including stair-climbing in a group of sedentary female students. The effect of this program on CRF was assessed, using a SCT before and after the program. During a 135sec SCT VO₂ and HR was measured with a portable spirometry. After the program a reduction in VO₂ and HR was measured, which suggested a cardiovascular health benefit. [168] The working group Lafortuna and Sartorio et al. examined obese individuals with a SCT. In their first study they assessed the anaerobic power output in adult obese individuals (BMI 30-60kg/m²). A modification of the Margaria stair climbing test, covering a flight with 13 steps was used. The individuals were asked to climb at their highest speed possible. With the collected data, they calculated the average mechanical power output: $W = (M_b \times g \times h) / t$ (M_b =body mass, g =gravity, h =vertical height, t =time). They showed a significant correlation between mechanical power output and BMI. A higher BMI was associated with a higher mechanical power output. The power output also correlated significantly with the amount of fat free mass, whereas men had a higher amount of fat free mass. With aging the fat free mass decreases in both genders. In their second study the group used the same SCT. In this population of obese women, not all were able to perform the test. In this study a <40 BMI had a significant effect on power output, values >40 had not. In the third study severely obese children and adolescents were examined with the same protocol as the obese adults before. They also showed a significant correlation between power output and fat free body mass. Up to the age of 13 they could not detect a significant difference between boys and girls. Due to age a significant increase in power output could be shown. From the age of 13 boys further gained power output, whereas girls started to build a plateau. [106,169,170] Oesch et al. examined patients with lower back pain. They used a SCT protocol, covering 100 steps. [201]

other parameters

Apart from the approaches mentioned above, there are a few trials, which investigated the effect of stair climbing regarding blood sugar [202-205], or cognitive function. [206] Another working group investigated the effect of a musical prompt on the pace of stair climbing. [15] Poppius et al. used a SCT to simulate exercise and evaluated whether doxantrazole can prevent exercise-induced asthma. [207] Aveline et al. assessed pain and recovery after knee arthroplasty. [208]

ery after knee arthroplasty. [208]

SCT-protocols and implementations

In most trials the test was conducted with 10 steps or more. Almost all studies evaluated the ascending as well as the descending. Other studies focused solely on ascending, as for example Lee et al. who used the SCT for evaluating the walking ability of stroke patients. [94] Step height was mentioned in the majority of the included trials (table 1-3). The used stairs varied in step height between 7,8 and 20 cm. (table 1-3) Some of the studies specified the total amount of steps to be covered and thus the total vertical height. A trial by Novoa et al. showed an adequate exertion level from 12 meters and more, comparable to an unlimited approach. [209] Several studies used a variation of the Margaria-SCT protocol, covering a total vertical height of 1,99m. [210] Other groups orientated towards the protocol used by Cataneo et al., covering a total vertical height of 12m. [185] Various trials used one staircase repetitively and instructed their patients to climb up and down for a certain amount of time. (table 1-3) This approach was for instance used by Smeets et al., who investigated the physical capacity of patients with lower back pain. [84]

measured SCT performance parameters

With a total of 122, most of the included trials used the total time needed to complete the SCT as the only measured parameter. 15 other trials counted the number of steps completed in a fixed time. Some trials calculated the cadence.

Other parameters collected included vital signs, the number of steps climbed and the so-called stair-climbing power (table 1-3). Capturing stair-climbing power was mentioned in 41 articles. To calculate the stair-climbing power, the height climbed, the body weight, the acceleration of gravity and the time required to complete the SCT were used in most of the cases. (table 1-3) The Cadence of stair-climbing was assessed in 6 trials. Predicted VO₂max was calculated in 7 studies.

Discussion

This overview shows how broadly the SCT can be used. Varying from age 2 to the elderly, healthy or ill. However, in most of these studies, the focus was solely set on the achieved time during SCT, or the number of steps climbed. Therefore, a lot of significant information was not collected. Even if the maximum oxygen collection during SCT may not always be directly the same as in CPET, a lot of trials could show a significant correlation, depending on the protocol that was used to perform the SCT.

Nevertheless, the information that can be gathered while performing an SCT are often not used detailed enough.

While almost all the included studies used time as the main recorded parameter, vital signs were not gathered in all trials, although this should be possible in significantly more studies and could provide important additional information regarding the physical condition of the participants without major effort. A lot

of important information seems to unremarkably fade away. For instance, a direct comparison of parameters gathered with CPET would be interesting. Brunelli et al. even showed that it is possible to obtain conclusive VO₂-peak values during SCT with a portable gas analyzer. They were also able to demonstrate that there is a correlation between the altitude climbed and the VO₂-peak recorded, which could also be interesting in terms of using the SCT as a maximal exercise test, for example. [182]

Moreover, it must be mentioned that the design and evaluation of the SCT differs a lot in between the trials. A guiding thread lacks, even as a basic protocol. For example, the number of steps ranges from a minimum of 1 step [199] to an unlimited number or not even a detailed description. Some used 1 flight repetitively, others used an unlimited staircase. The total height was not always entitled. Some used a given time frame and assessed the number of steps climbed, others used an unlimited time, and subjects were asked to climb until reaching fatigue, others used a given staircase or number of steps/flights and assessed the time needed to complete. There is also a wide variation regarding the instruction, some trials allowed the use of a handrail or other walking aids.

The given instructions concerning the use of aids can of course influence the test results. For instance, if mobility is examined with the SCT and the participants are for example stroke patients with gait instability, as conducted by Mustafaoglu et al. [96] it obviously makes a difference whether the participants are allowed to use walking aids or not. Another example is the use of a handrail if leg strength is investigated and, in some studies, e.g. Daly et al. the participants are allowed to support themselves with the handrail, while in others they are not, which again could lead to different results. [154]

The different numbers of steps used seem to be due to the different research questions and patient groups. For example, Pate et al. used a staircase with 21 steps per flight and unlimited steps taken to evaluate the cardiorespiratory fitness of their patients. [174] They wanted to use the SCT as a maximal exercise test. Whereas in trials, using only a few steps, as is the trials of Brun-Olsen et al. who evaluated physical function and mobility in patients during early rehabilitation after knee arthroplasty [212], the SCT was used as a submaximal exercise test for most individuals. This difference is also obvious in the different trials, focusing on anaerobic capacity and therefore using the Margaria protocol [210], which only covers 1,99m vertical height in comparison to the trials, focusing on aerobic fitness and aiming on maximum exertion, using the protocol of Cataneo et al. [185]

Furthermore, it is interesting to note that among the studies included, only a limited number have attempted to standardize the SCT or build upon previously collected data. One notable approach within this context is the utilization of stair-climbing power, as described earlier. This is for instance used by Storer et al. and Harmer et al. [21,115,118,170,185,195,197,215] It is an important beneficial information gain, as some trials which for

example used the SCT to determine the cardiorespiratory fitness of their patients [174] could show a significant correlation to VO₂max in CPET, as it has been shown in studies before. [185] Considering this correlation, especially SCT power, defined as body weight x total vertical height divided by SCT time could be a crucial information regarding the CRF assessment with a simple SCT and vice versa a huge lack of information when not gathered while performing SCT.

Almost all the included studies recorded the time required to complete the test whereas only a few recorded other parameters such as vital signs or even blood lactate. (table 1-3) In general, there seems to be a lot of potential in performing a SCT. Nevertheless, the approaches should be focused on a standardized protocol to gather as much information as possible with this simple test. One possible limitation of the SCT is that the patients may need to be accompanied by staff. This could be necessary for safety reasons, as mentioned for instance by Kraemer et al. [54] but also, for example, to collect additional data during the test. This may make the SCT more staff-intensive. However, compared to other tests such as CPET, it would still probably require less know-how, equipment and preparation time and is therefore easier to include in less time-consuming examinations.

Limitations

This review has limitations. There is only a limited number of trials, using a SCT. The reviewed studies also have limitations, such as small sample sizes, varying levels of evidence, and different approaches regarding protocols and recorded results. Moreover, the search was conducted by only two researchers; therefore, the false exclusion of articles cannot be fully dismissed. Nevertheless, the methods and results were approved by the other researchers.

Conclusions

The significance of CRF as a prognostic indicator for long-term health outcomes cannot be overstated. Despite its pivotal role, a thorough assessment of CRF is not yet part of routine medical evaluations. While established methods like CPET provide comprehensive insights into CRF, their extensive resource requirements limit their widespread application in routine check-ups.

The SCT presents a promising alternative due to its simplicity and resource-efficiency as well as its wide applicability during age groups. However, the diverse unstandardized protocols and the wide distributed applications across various studies underscore the need for standardization in its implementation. With the significant correlations shown between SCT results and established clinical tests, such as spirometry and the 6-minute walking test, the SCT holds potential for assessing CRF and functional exercise capacity in diverse patient groups. However, it would again be necessary to obtain comparative data, for example from healthy test subjects, in order to be able to use this test also in patients and even pediatric patients.

Establishing standardized SCT protocols could pave the way

for its integration into routine medical assessments, benefiting both general populations and specific patient cohorts, particularly children and young adults with congenital diseases such as congenital heart disease. Nonetheless, further research with larger sample sizes and consensus on standardized SCT protocols are imperative to solidify its position as a comprehensive assessment tool within the medical field.

Future Goal – Standardization

Studying all those different trials, we developed a standardized protocol, covering all important points of the previous trials.

We aimed for an easy exercise test, which can be used as sub-maximal and/or maximum test as a supplement to CPET. Our protocol therefore covered 4 flights with a total vertical height of 13,14m. As previous trials showed that a total of at least 12meters vertical height was needed to achieve maximum exertion. The test should be conducted at an individual's maximum pace, without the use of a handrail and without taking 2steps or more. At the top of the 4 flights the individual should turn around and climb down the stairs. Vital signs before and after the test should be recorded. The time needed to complete the test was stopped and the SCT power calculated using the following formula: $SCT\text{-}Index = (\text{body weight} \times \text{staircase height}) / (SCT\text{time})$. Moreover, we compared the results in SCT (SCTtime and SCTpower) with the results in 6MWT (distance) and CPET (VO₂max(ml/min); VO₂max (ml/min/kg); oxygen pulse). We could show a significant correlation in different study groups as previously described in the studies above.

We assessed the clinical validity of the SCT in healthy adults as well as in obese adults. Furthermore, in healthy children and adolescents as well as children and adolescents with congenital heart disease. Overall, we could show promising results for the clinical use of the SCT with this standardized protocol in both, healthy and ill patients, as long as they are able to climb stairs. The SCT is an easy tool to assess CRF, either as a tool in preventive medicine or in the assessment of clinical treatment or condition control.

References

- Mandsager K, Harb S, Cremer P, Phelan D, Nissen SE, et al. (2018) Association of Cardiorespiratory Fitness With Long-term Mortality Among Adults Undergoing Exercise Treadmill Testing. *JAMA Network Open* 1(6): e183605-e183605.
- Al-Mallah MH, Sakr S, Al-Qunaibet A (2018) Cardiorespiratory Fitness and Cardiovascular Disease Prevention: an Update. *Curr Atheroscler Rep* 20(1): 1.
- Vainshelboim B, Lima RM, Myers J (2019) Cardiorespiratory fitness and cancer in women: A prospective pilot study. *J Sport Health Sci* 8(5): 457-462.
- WHO guidelines on physical activity and sedentary behaviour. Geneva: World Health Organization 2020.
- Dold SK, Haas NA, Apitz C (2023) Effects of Sports, Exercise Training, and Physical Activity in Children with Congenital Heart Disease-A Review of the Published Evidence. *Children (Basel)* 10(2).
- Tran D (2018) Cardiopulmonary Exercise Testing. *Methods Mol Biol* 1735: 285-295.
- Arena R, Canada JM, Popovic D, Trankle CR, Buono MGD, et al. (2020) Cardiopulmonary exercise testing - refining the clinical perspective by combining assessments. *Expert Rev Cardiovasc Ther* 18(9): 563-576.
- Ortega FB, Ruiz JR, Castillo MJ, Sjöström M (2008) Physical fitness in childhood and adolescence: a powerful marker of health. *Int J Obes (Lond)* 32(1): 1-11.
- (2003) ATS/ACCP Statement on cardiopulmonary exercise testing. *Am J Respir Crit Care Med* 167(2): 211-77.
- Herdy AH, Herdy AH, Milani M, Hossri C, et al. (2016) Cardiopulmonary Exercise Test: Background, Applicability and Interpretation. *Arq Bras Cardiol* 107(5): 467-481.
- McKelvie RS, Jones NL (1989) Cardiopulmonary exercise testing. *Clinics in chest medicine* 10(2): 277-291.
- Malhotra R, Bakken K, Elia ED, Lewis GD (2016) Cardiopulmonary Exercise Testing in Heart Failure. *JACC Heart Fail* 4(8): 607-16.
- Agarwala P, Salzman SH (2020) Six-Minute Walk Test: Clinical Role, Technique, Coding, and Reimbursement. *Chest* 157(3): 603-611.
- Giannitsi S, Bougiakli M, Bechlioulis A, Kotsia A, Michalis LK, et al. (2019) 6-minute walking test: a useful tool in the management of heart failure patients. *Ther Adv Cardiovasc Dis* 13: 1753944719870084.
- Khan A, Biddiss E (2015) Musical stairs: the impact of audio feedback during stair-climbing physical therapies for children. *Disabil Rehabil Assist Technol* 10(3): 231-5.
- Heiberg KE, Bruun-Olsen V, Ekeland A, Mengshoel AM (2012) Effect of a walking skill training program in patients who have undergone total hip arthroplasty: Followup one year after surgery. *Arthritis Care Res (Hoboken)* 64(3): 415-23.
- Mizner RL, Petterson SC, Snyder-Mackler L (2005) Quadriceps strength and the time course of functional recovery after total knee arthroplasty. *J Orthop Sports Phys Ther* 35(7): 424-36.
- Mizner RL, Petterson SC, Stevens JE, Axe MJ, Snyder-Mackler L (2005) Preoperative quadriceps strength predicts functional ability one year after total knee arthroplasty. *J Rheumatol* 32(8): 1533-9.
- Nyland J, Frost K, Quesada P, Angeli C, Swank A, et al. (2007) Self-reported chair-rise ability relates to stair-climbing readiness of total knee arthroplasty patients: a pilot study. *J Rehabil Res Dev* 44(5): 751-9.
- Bruun-Olsen V, Heiberg KE, Wahl AK, Mengshoel AM (2013) The immediate and long-term effects of a walking-skill program compared to usual physiotherapy care in patients who have undergone total knee arthroplasty (TKA): a randomized controlled trial. *Disabil Rehabil* 35(23): 2008-15.
- Harmer AR, Naylor JM, Crosbie J, Russell T (2009) Land-based versus water-based rehabilitation following total knee replacement: a randomized, single-blind trial. *Arthritis Rheum* 61(2): 184-91.
- Farquhar S, Snyder-Mackler L (2010) The Chitranjan Ranawat Award: The nonoperated knee predicts function 3 years after unilateral total knee arthroplasty. *Clin Orthop Relat Res* 468(1): 37-44.
- Petterson SC, Mizner RL, Stevens JE, Rasis L, Bodenstab A, et al. (2009) Improved function from progressive strengthening interventions after total knee arthroplasty: a randomized clinical trial with an imbedded prospective cohort. *Arthritis Rheum* 61(2): 174-83.
- Yoshida Y, Mizner RL, Ramsey DK, Snyder-Mackler L (2008) Examining outcomes from total knee arthroplasty and the relationship between quadriceps strength and knee function over time. *Clin Biomech (Bristol, Avon)* 23(3): 320-8.

25. Stevens-Lapsley JE, Petterson SC, Mizner RL, Snyder-Mackler L (2010) Impact of body mass index on functional performance after total knee arthroplasty. *J Arthroplasty* 25(7): 1104-9.
26. Zeni JA, Snyder-Mackler L (2010) Early postoperative measures predict 1- and 2-year outcomes after unilateral total knee arthroplasty: importance of contralateral limb strength. *Phys Ther* 90(1): 43-54.
27. Zeni JA, Axe MJ, Snyder-Mackler L (2010) Clinical predictors of elective total joint replacement in persons with end-stage knee osteoarthritis. *BMC Musculoskelet Disord* 11: 86.
28. Christiansen CL, Bade MJ, Judd DL, Stevens-Lapsley JE (2011) Weight-bearing asymmetry during sit-stand transitions related to impairment and functional mobility after total knee arthroplasty. *Arch Phys Med Rehabil* 92(10): 1624-9.
29. Chung JY, Min BH (2013) Is bicompartamental knee arthroplasty more favourable to knee muscle strength and physical performance compared to total knee arthroplasty? *Knee Surg Sports Traumatol Arthrosc* 21(11): 2532-41.
30. van Leeuwen DM, de Ruiter CJ, Nolte PA, Haan AD (2014) Preoperative strength training for elderly patients awaiting total knee arthroplasty. *Rehabil Res Pract* 2014: 462750.
31. Akbaba YA, Yeldan I, Guney N, Ozdinciler AR (2016) Intensive supervision of rehabilitation programme improves balance and functionality in the short term after bilateral total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc* 24(1): 26-33.
32. Loyd BJ, Jennings JM, Judd DL, Kim RH, Wolfe P, et al. (2017) Influence of Hip Abductor Strength on Functional Outcomes Before and After Total Knee Arthroplasty: Post Hoc Analysis of a Randomized Controlled Trial. *Phys Ther* 97(9): 896-903.
33. Moukarzel M, Guillot A, Rienzo FD, Hoyek N (2019) The therapeutic role of motor imagery during the chronic phase after total knee arthroplasty: a pilot randomized controlled trial. *Eur J Phys Rehabil Med* 55(6): 806-815.
34. Gränicher P, Stöggli T, Fucetese SF, Adelsberger R, Swanenburg J (2020) Preoperative exercise in patients undergoing total knee arthroplasty: a pilot randomized controlled trial. *Arch Physiother* 10: 13.
35. Jacksteit R, Stöckel T, Behrens M, Feldhege F, Bergschmidt P, et al. (2021) Low-Load Unilateral and Bilateral Resistance Training to Restore Lower Limb Function in the Early Rehabilitation After Total Knee Arthroplasty: A Randomized Active-Controlled Clinical Trial. *Front Med (Lausanne)* 8: 628021.
36. Choi JH, Kim BR, Rim Kim S, Nam KW, Lee SY, et al. (2021) Performance-based physical function correlates with walking speed and distance at 3 months post unilateral total knee arthroplasty. *Gait Posture* 87: 163-169.
37. Pozzi F, White DK, Snyder-Mackler L, Zeni JA (2020) Restoring physical function after knee replacement: a cross sectional comparison of progressive strengthening vs standard physical therapy. *Physiother Theory Pract* 36(1): 122-133.
38. Winters JD, Christiansen CL, Stevens-Lapsley JE (2014) Preliminary investigation of rate of torque development deficits following total knee arthroplasty. *Knee* 21(2): 382-6.
39. Tsukagoshi R, Tateuchi H, Fukumoto Y, Ibuki S, Akiyama H, et al. (2014) Functional performance of female patients more than 6 months after total hip arthroplasty shows greater improvement with weight-bearing exercise than with non-weight-bearing exercise. Randomized controlled trial. *Eur J Phys Rehabil Med* 50(6): 665-75.
40. Larsen JB, Mogensen L, Arendt-Nielsen L, Madeleine P (2020) Intensive, personalized multimodal rehabilitation in patients with primary or revision total knee arthroplasty: a retrospective cohort study. *BMC Sports Sci Med Rehabil* 12: 5.
41. Heiberg K, Bruun-Olsen V, Mengshoel AM (2010) Pain and recovery of physical functioning nine months after total knee arthroplasty. *J Rehabil Med* 42(7): 614-9.
42. Heiberg KE, Figved W (2016) Physical Functioning and Prediction of Physical Activity After Total Hip Arthroplasty: Five-Year Followup of a Randomized Controlled Trial. *Arthritis Care Res (Hoboken)* 68(4): 454-62.
43. Galea MP, Westh R, McMeeken J, Tully E, Weller R, et al. (2008) A targeted home- and center-based exercise program for people after total hip replacement: a randomized clinical trial. *Arch Phys Med Rehabil* 89(8): 1442-7.
44. Bade MJ, Kohrt WM, Stevens-Lapsley JE (2010) Outcomes before and after total knee arthroplasty compared to healthy adults. *J Orthop Sports Phys Ther* 40(9): 559-67.
45. Bade MJ, Struessel T, Dayton M, Foran J, Kim RH, et al. (2017) Early High-Intensity Versus Low-Intensity Rehabilitation After Total Knee Arthroplasty: A Randomized Controlled Trial. *Arthritis Care Res (Hoboken)* 69(9): 1360-1368.
46. Bade MJ, Wolfe P, Zeni JA, Stevens-Lapsley JE, Snyder-Mackler L (2012) Predicting poor physical performance after total knee arthroplasty. *J Orthop Res* 30(11): 1805-10.
47. Marmon AR, Milcarek BI, Snyder-Mackler L (2014) Associations between knee extensor power and functional performance in patients after total knee arthroplasty and normal controls without knee pain. *Int J Sports Phys Ther* 9(2): 168-78.
48. Kim JH, Kim WB, Lee SY, Nam KW, Han EY, et al. (2019) Functional Outcomes After Critical Pathway for Inpatient Rehabilitation of Total Knee Arthroplasty. *Ann Rehabil Med* 43(6): 650-661.
49. Suh MJ, Kim BR, Kim SR, Han EY, Lee SY (2017) Effects of Early Combined Eccentric-Concentric Versus Concentric Resistance Training Following Total Knee Arthroplasty. *Ann Rehabil Med* 41(5): 816-827.
50. Kirschner J, Michel S, Becker R, Stiebitz O, Hommel H, et al. (2023) Determination of Relationships between Symmetry-Based, Performance-Based, and Functional Outcome Measures in Patients Undergoing Total Hip Arthroplasty. *J Pers Med* 13(7).
51. Lee SJ, Han EY, Kim JH, Park YG, Lee SY, et al. (2020) Preoperative physical factors that predict stair-climbing ability at one month after total knee arthroplasty. *J Rehabil Med* 52(5): jrm00064.
52. Judd DL, Wolfe P, LeDoux CV, Hogan C, Dayton MR, et al. (2019) Trajectories of functional performance and muscle strength recovery differ after total knee and total hip replacement: a performance-based, longitudinal study. *Int J Rehabil Res* 42(3): 211-216.
53. Kovács I, Bender T (2002) The therapeutic effects of Cserkeszőlő thermal water in osteoarthritis of the knee: a double blind, controlled, follow-up study. *Rheumatol Int* 21(6): 218-21.
54. Kraemer WJ, Ratamess NR, Anderson JM, Maresh CM, Tiberio DP, et al. (2004) Effect of a cetylated fatty acid topical cream on functional mobility and quality of life of patients with osteoarthritis. *J Rheumatol* 31(4): 767-74.
55. Laufer Y, Nahir AM, Zilberman R, Porat R (2005) Effect of pulsed short-wave diathermy on pain and function of subjects with osteoarthritis of the knee: a placebo-controlled double-blind clinical trial. *Clin Rehabil* 19(3): 255-63.
56. Schmitt LC, Fitzgerald GK, Reisman AS, Rudolph KS (2008) Instability, laxity, and physical function in patients with medial knee osteoarthritis. *Phys Ther* 88(12): 1506-16.
57. Vincent HK, Montero C, Conrad BP, Horodyski M, Connelly J, et al. (2013) "Functional pain," functional outcomes, and quality of life after hyaluronic acid intra-articular injection for knee osteoarthritis. *Pm r* 5(4): 310-8.

58. Brennehan EC, Kuntz AB, Wiebenga EG, Maly MR (2015) A Yoga Strengthening Program Designed to Minimize the Knee Adduction Moment for Women with Knee Osteoarthritis: A Proof-Of-Principle Cohort Study. *PLoS One* 10(9): e0136854.
59. Vongsirinavarat M, Nilmarat P, Somprasong S, Apinonkul B (2020) Identification of knee osteoarthritis disability phenotypes regarding activity limitation: a cluster analysis. *BMC Musculoskelet Disord* 21(1): 237.
60. Onodera CMK, Coelho-Júnior HJ, Carvalho Sampaio RA, Duarte Lana JFS, Uchida MC, et al. (2020) The importance of objectively measuring functional tests in complement to self-report assessments in patients with knee osteoarthritis. *Gait Posture* 82: 33-37.
61. Khruakhorn S, Chiwarakranon S (2021) Effects of hydrotherapy and land-based exercise on mobility and quality of life in patients with knee osteoarthritis: a randomized control trial. *J Phys Ther Sci* 33(4): 375-383.
62. De Zwart A, Dekker J, Verberne S, Roorda LD, Lems WF, et al. (2022) Association Between Measures of Muscle Strength and Performance of Daily Activities in Patients with Knee Osteoarthritis. *J Rehabil Med* 54: jrm00316.
63. Bieler T, Magnusson SP, Kjaer M, Beyer N (2014) Intra-rater reliability and agreement of muscle strength, power and functional performance measures in patients with hip osteoarthritis. *J Rehabil Med* 46(10): 997-1005.
64. Bieler T, Siersma V, Magnusson SP, Kjaer M, Beyer N, et al. (2017) In hip osteoarthritis, Nordic Walking is superior to strength training and home-based exercise for improving function. *Scand J Med Sci Sports* 27(8): 873-886.
65. Freisinger GM, Hutter EE, Lewis J, Granger JF, Pan X, et al. (2017) Relationships between varus-valgus laxity of the severely osteoarthritic knee and gait, instability, clinical performance, and function. *J Orthop Res* 35(8): 1644-1652.
66. Petersson N, Jørgensen SL, Kjeldsen T, Mechlenburg I, Aagaard P (2022) Blood Flow Restricted Walking in Elderly Individuals with Knee Osteoarthritis: A Feasibility Study. *J Rehabil Med* 54: jrm00282.
67. Pua YH, Wrigley TV, Collins M, Cowan SM, Bennell KL (2009) Self-report and physical performance measures of physical function in hip osteoarthritis: relationship to isometric quadriceps torque development. *Arthritis Rheum* 61(2): 201-8.
68. Carvalho C, Gonçalves GH, Petrella M, Mattiello SM, Oliveira Sato TD, et al. (2023) Association between ankle torque and performance-based tests, self-reported pain, and physical function in patients with knee osteoarthritis. *Arch Rheumatol* 38(3): 387-396.
69. Iijima H, Eguchi R, Shimoura K, Aoyama T, Takahashi M (2019) Stair climbing ability in patients with early knee osteoarthritis: Defining the clinical hallmarks of early disease. *Gait Posture* 72: 148-153.
70. Shimoura K, Iijima H, Suzuki Y, Aoyama T (2019) Immediate Effects of Transcutaneous Electrical Nerve Stimulation on Pain and Physical Performance in Individuals with Preradiographic Knee Osteoarthritis: A Randomized Controlled Trial. *Arch Phys Med Rehabil* 100(2): 300-306.e1.
71. Jankaew A, Lin You Y, Yang TH, Chang YW, Lin CF (2023) The effects of low-level laser therapy on muscle strength and functional outcomes in individuals with knee osteoarthritis: a double-blinded randomized controlled trial. *Sci Rep* 13(1): 165.
72. Lee SH, Kao CC, Liang HW, Wu HT (2022) Validity of the Osteoarthritis Research Society International (OARSI) recommended performance-based tests of physical function in individuals with symptomatic Kellgren and Lawrence grade 0-2 knee osteoarthritis. *BMC Musculoskelet Disord* 23(1): 1040.
73. Kim BS, Lee HJ, Kim SR, Choi JH, Kim BR, et al. (2021) Associations Between Obesity With Low Muscle Mass and Physical Function in Patients With End-Stage Knee Osteoarthritis. *Geriatr Orthop Surg Rehabil* 12: 21514593211020700.
74. Suh MJ, Kim WB, Park YG, Lee SY, Nam KW, et al. (2019) Bilateral Quadriceps Muscle Strength and Pain Correlate with Gait Speed and Gait Endurance Early After Unilateral Total Knee Arthroplasty: A Cross-sectional Study. *Am J Phys Med Rehabil* 98(10): 897-905.
75. Hsieh RL, Lee WC (2014) Immediate and medium-term effects of custom-moulded insoles on pain, physical function, physical activity, and balance control in patients with knee osteoarthritis. *J Rehabil Med* 46(2): 159-65.
76. Hsieh RL, Lee WC (2016) Clinical effects of lateral wedge arch support insoles in knee osteoarthritis: A prospective double-blind randomized study. *Medicine (Baltimore)* 95(27): e3952.
77. Baert IA, Luyten FP, Verschueren SMP, Staes F, Jonkerset I, al. (2023) Proprioceptive accuracy in women with early and established knee osteoarthritis and its relation to functional ability, postural control, and muscle strength. *Clin Rheumatol* 32(9): 1365-74.
78. Baert IA, Staes F, Truijen S, Mahmoudian A, Noppe N, et al. (2014) Weak associations between structural changes on MRI and symptoms, function and muscle strength in relation to knee osteoarthritis. *Knee Surg Sports Traumatol Arthrosc* 22(9): 2013-25.
79. Marmon AR, Zeni JA, Snyder-Mackler L (2013) Perception and presentation of function in patients with unilateral versus bilateral knee osteoarthritis. *Arthritis Care Res (Hoboken)* 65(3): 406-13.
80. Zeni J, Rasis L, Abujaber S, Pozzi F (2014) Relationship between strength, pain, and different measures of functional ability in patients with end-stage hip osteoarthritis. *Arthritis Care Res (Hoboken)* 66(10): 1506-12.
81. Beckmann M, Bruun-Olsen V, Pripp AH, Bergland A, Smith T, et al. (2022) Recovery and prediction of physical function 1 year following hip fracture. *Physiother Res Int* 27(3): e1947.
82. Adunsky A, Papanicolaou DA, Liu N, Chandler J, Heyden N, et al. (2011) MK-0677 (ibuprofen mesylate) for the treatment of patients recovering from hip fracture: a multicenter, randomized, placebo-controlled phase IIb study. *Arch Gerontol Geriatr* 53(2): 183-9.
83. Ljungquist T, Nygren A, Jensen I, Harms-Ringdahl K (2003) Physical performance tests for people with spinal pain--sensitivity to change. *Disabil Rehabil* 25(15): 856-66.
84. Smeets RJ, van Geel AC, Kester AD, Knottnerus JA (2007) Physical capacity tasks in chronic low back pain: what is the contributing role of cardiovascular capacity, pain and psychological factors? *Disabil Rehabil* 29(7): 577-86.
85. Andersson EI, Lin CC, Smeets RJ (2010) Performance tests in people with chronic low back pain: responsiveness and minimal clinically important change. *Spine (Phila Pa 1976)* 35(26): E1559-63.
86. Mengshoel AM, Jokstad K, Bjerkhoel F (2004) Associations between walking time, quadriceps muscle strength and cardiovascular capacity in patients with rheumatoid arthritis and ankylosing spondylitis. *Clin Rheumatol* 23(4): 299-305.
87. Wetzel JL, Fry DK, Pflizer LA (2011) Six-minute walk test for persons with mild or moderate disability from multiple sclerosis: performance and explanatory factors. *Physiother Can* 63(2): 166-80.
88. Nunes GS, Pizzari T, Barton CJ, Crossley KM, Serrão FV, et al. (2019) People with patellofemoral pain have impaired functional performance, that is correlated to hip muscle capacity. *Phys Ther Sport* 40: 85-90.
89. Alfano LN, Lowes LP, Flanigan KM, Mendell JR (2013) Correlation of knee strength to functional outcomes in Becker muscular dystrophy.

- Muscle Nerve 47(4): 550-4.
90. Collado-Mateo D, Adsuar JC, Olivares PR, Gusi N, Maestre-Cascales C, et al. (2016) Performance of women with fibromyalgia in walking up stairs while carrying a load. *PeerJ* 4: e1656.
 91. Schwid SR, Petrie MD, McDermott MP, Tierney DS, Mason DH, et al. (197) Quantitative assessment of sustained-release 4-aminopyridine for symptomatic treatment of multiple sclerosis. *Neurology* 48(4): 817-21.
 92. Suslov VM, Lieberman LN, Carlier PG, Ponomarenko GN, Ivanov DO, et al. (2023) Efficacy and safety of hydrokinesitherapy in patients with dystrophinopathy. *Front Neurol* 14: 1230770.
 93. Katz-Leurer M, Shochina M, Carmeli E, Friedlander Y (2003) The influence of early aerobic training on the functional capacity in patients with cerebrovascular accident at the subacute stage. *Arch Phys Med Rehabil* 84(11): 1609-14.
 94. Lee MJ, Kilbreath SL, Singh MF, Zeman B, Lord SR, et al. (2008) Comparison of effect of aerobic cycle training and progressive resistance training on walking ability after stroke: a randomized sham exercise-controlled study. *J Am Geriatr Soc* 56(6): 976-85.
 95. Mustafaoglu R, Erhan B, Yeldan I, Hüseyinsinoğlu BE, Gündüz B, et al. (2018) The effects of body weight-supported treadmill training on static and dynamic balance in stroke patients: A pilot, single-blind, randomized trial. *Turk J Phys Med Rehabil* 64(4): 344-352.
 96. Mustafaoglu R, Erhan B, Yeldan I, Gunduz B, Tarakci E (2020) Does robot-assisted gait training improve mobility, activities of daily living and quality of life in stroke? A single-blinded, randomized controlled trial. *Acta Neurol Belg* 120(2): 335-344.
 97. Ahmed Burq HSI, Karimi H, Ahmad A, Gilani SA, Hanif A (2021) Effect of whole-body vibration on obstacle clearance and stair negotiation time in chronic stroke patients; A randomized controlled trial. *J Bodyw Mov Ther* 27: 698-704.
 98. Ihl T, Ahmadi M, Laumeier I, Steinicke M, Fers C, et al. (2022) Patient-Centered Outcomes in a Randomized Trial Investigating a Multimodal Prevention Program After Transient Ischemic Attack or Minor Stroke: The INSPiRE-TMS Trial. *Stroke* 53(9): 2730-2738.
 99. Sharp SA, Brouwer BJ (1997) Isokinetic strength training of the hemiparetic knee: effects on function and spasticity. *Arch Phys Med Rehabil* 78(11): 1231-6.
 100. Teixeira-Salmela LF, Olney SJ, Nadeau S, Brouwer B (1999) Muscle strengthening and physical conditioning to reduce impairment and disability in chronic stroke survivors. *Arch Phys Med Rehabil* 80(10): 1211-8.
 101. Bonan IV, Yelnik AP, Colle FM, Michaud C, Normand E, et al. (2004) Reliance on visual information after stroke. Part II: Effectiveness of a balance rehabilitation program with visual cue deprivation after stroke: a randomized controlled trial. *Arch Phys Med Rehabil* 85(2): 274-8.
 102. Harries N, Loeppky JA, Shaheen S, Al-Jarrah M, Molteni F, et al. (2015) A stair-climbing test for measuring mechanical efficiency of ambulation in adults with chronic stroke. *Disabil Rehabil* 37(11): 1004-8.
 103. van de Port IG, Wevers LEG, Lindeman E, Kwakkel G (2012) Effects of circuit training as alternative to usual physiotherapy after stroke: randomised controlled trial. *Bmj* 344: e2672.
 104. Zaino CA, Marchese VG, Westcott SL (2004) Timed up and down stairs test: preliminary reliability and validity of a new measure of functional mobility. *Pediatr Phys Ther* 16(2): 90-8.
 105. Bar-Haim S, Belokopytov M, Harries N, Loeppky JA, Kaplanski J (2008) Prediction of mechanical efficiency from heart rate during stair-climbing in children with cerebral palsy. *Gait Posture* 27(3): 512-7.
 106. Sartorio A, Lafortuna CL, Conte G, Faglia G, Narici MV (2001) Changes in motor control and muscle performance after a short-term body mass reduction program in obese subjects. *J Endocrinol Invest* 24(6): 393-8.
 107. Grant S, Todd K, Aitchison TC, Kelly P, Stoddart D (2004) The effects of a 12-week group exercise programme on physiological and psychological variables and function in overweight women. *Public Health* 118(1): 31-42.
 108. Bayartai ME, Luomajoki H, Micheli RD, Tringali G, Marazzi N, et al. (2022) Changes in the Oswestry Disability Index after a 3-Week In-Patient Multidisciplinary Body Weight Reduction Program in Adults with Obesity. *J Clin Med* 11(11): 3175.
 109. Bittel DC, Bittel AJ, Tuttle LJ, Hastings MK, Commeyan PK, et al. (2015) Adipose tissue content, muscle performance and physical function in obese adults with type 2 diabetes mellitus and peripheral neuropathy. *J Diabetes Complications* 29(2): 250-7.
 110. Maffiuletti NA, Tringali G, Patrizi A, Agosti F, Sartorio A (2017) Reproducibility of clinician-friendly physical performance measures in individuals with obesity. *J Rehabil Med* 49(8): 677-681.
 111. Sousa-Gonçalves CR, Tringali G, Tamini S, Micheli RD, Soranna D, et al. (2019) Acute Effects of Whole-Body Vibration Alone or in Combination with Maximal Voluntary Contractions on Cardiorespiratory, Musculoskeletal, and Neuromotor Fitness in Obese Male Adolescents. *Dose Response* 17(4): 1559325819890492.
 112. Tamini S, Micheli RD, Tringali G, Bernardo-Filho M, Sartorio A (2020) Acute Effects of Whole-Body Vibration Exercises at 2 Different Frequencies Versus an Aerobic Exercise on Some Cardiovascular, Neuromotor and Musculoskeletal Parameters in Adult Patients with Obesity. *Dose Response* 18(4): 1559325820965005.
 113. Rigamonti AE, Tamini S, Saezza A, Cella SG, Sartorio A, et al. (2020) Impact of a Three-Week in-Hospital Multidisciplinary Body Weight Reduction Program on Body Composition, Muscle Performance and Fatigue in a Pediatric Obese Population with or without Metabolic Syndrome. *Nutrients* 12(1).
 114. Lazzar S, D'Alleva M, Vaccari F, Tringali G, Micheli RD, et al. (2022) Effects of a 3-Week Inpatient Multidisciplinary Body Weight Reduction Program on Body Composition and Physical Capabilities in Adolescents and Adults with Obesity. *Front Nutr* 9: 840018.
 115. Guerrini Usubini A, Bottacchi M, Bondesan A, Caroli D, Castelnuovo G, et al. (2022) A three-week in-hospital multidisciplinary body weight reduction program exerts beneficial effects on physical and mental health and fatigability of elderly patients with obesity. *Front Aging Neurosci* 14: 1054941.
 116. Knapp PE, LaValley M, Zhang A, Ulloor J, Bhasin S, et al. (2008) Effects of a supraphysiological dose of testosterone on physical function, muscle performance, mood, and fatigue in men with HIV-associated weight loss. *Am J Physiol Endocrinol Metab* 294(6): E1135-43.
 117. Nilsen TS, Raastad T, Skovlund E, Courneya KS, Langberg CW, et al. (2015) Effects of strength training on body composition, physical functioning, and quality of life in prostate cancer patients during androgen deprivation therapy. *Acta Oncol* 54(10): 1805-13.
 118. Storer TW, Pencina K, Li Z, Travison TG, Miciek R, et al. (2017) Effects of Testosterone Supplementation for 3 Years on Muscle Performance and Physical Function in Older Men. *J Clin Endocrinol Metab* 102(2): 583-593.
 119. Gagliano-Jucá T, Hettwer S, Dahinden P, Bhasin S, Basaria S, et al. (2018) Testosterone does not affect agrin cleavage in mobility-limited older men despite improvement in physical function. *Andrology* 6(1): 29-36.

120. Sattler F, Yarasheski K, Binder E, Schroeder ED, Kawakubo M, et al. (2011) Testosterone threshold levels and lean tissue mass targets needed to enhance skeletal muscle strength and function: the HORMA trial. *J Gerontol A Biol Sci Med Sci* 66(1): 122-9.
121. Travison TG, Lazzari A, Davis S, Ni P, Collins L, et al. (2011) Clinical meaningfulness of the changes in muscle performance and physical function associated with testosterone administration in older men with mobility limitation. *J Gerontol A Biol Sci Med Sci* 66(10): 1090-9.
122. Galvão DA, Nosaka K, Taaffe DR, Spry N, Kristjanson LJ, et al. (2006) Resistance training and reduction of treatment side effects in prostate cancer patients. *Med Sci Sports Exerc* 38(12): 2045-52.
123. Chikani V, Cuneo RC, Hickman I, Ho KKY (2015) Impairment of anaerobic capacity in adults with growth hormone deficiency. *J Clin Endocrinol Metab* 100(5): 1811-8.
124. Skelton DA, Young A, Greig CA, Malbut KE (1995) Effects of resistance training on strength, power, and selected functional abilities of women aged 75 and older. *J Am Geriatr Soc* 43(10): 1081-7.
125. Hiroyuki S, Uchiyama Y, Kakurai S (2003) Specific effects of balance and gait exercises on physical function among the frail elderly. *Clin Rehabil* 17(5): 472-9.
126. Seynnes O, Hue O, Pras P, Legros P, Bernard PL, et al. (2004) Physiological and functional responses to low-moderate versus high-intensity progressive resistance training in frail elders. *J Gerontol A Biol Sci Med Sci* 59(5): 503-9.
127. Henwood TR, Taaffe DR (2006) Short-term resistance training and the older adult: the effect of varied programmes for the enhancement of muscle strength and functional performance. *Clin Physiol Funct Imaging* 26(5): 305-13.
128. Capodaglio P, Edda MC, Facioli M, Saibene F (2007) Long-term strength training for community-dwelling people over 75: impact on muscle function, functional ability and life style. *Eur J Appl Physiol* 100(5): 535-42.
129. Capodaglio P, Capodaglio EM, Ferri A, Scaglioni G, Marchi A, Saibene F (2005) Muscle function and functional ability improves more in community-dwelling older women with a mixed-strength training programme. *Age Ageing* 34(2): 141-7.
130. Westlake KP, Wu Y, Culham EG (2007) Velocity discrimination: reliability and construct validity in older adults. *Hum Mov Sci* 26(3): 443-56.
131. Kortebein P, Protas E, Conger S, Lombeida J, Wolfe R, et al. (2008) Functional impact of 10 days of bed rest in healthy older adults. *J Gerontol A Biol Sci Med Sci* 63(10): 1076-81.
132. LeBrasseur NK, Miciek R, Mazer N, Storer TW, Bhasin S, et al. (2009) Effects of testosterone therapy on muscle performance and physical function in older men with mobility limitations (The TOM Trial): design and methods. *Contemp Clin Trials* 30(2): 133-40.
133. Hirota C, Watanabe M, Sun W, Tanimoto Y, Kono R, et al. (2010) Association between the Trail Making Test and physical performance in elderly Japanese. *Geriatr Gerontol Int* 10(1): 40-7.
134. Altubasi, I.M., Is quadriceps muscle strength a determinant of the physical function of the elderly? *J Phys Ther Sci*, 2015. 27(10): p. 3035-8.
135. Dias, C.P., et al., Effects of eccentric-focused and conventional resistance training on strength and functional capacity of older adults. *Age (Dordr)*, 2015. 37(5): p. 99.
136. Pirotta S, Kidgell DJ, Daly RM (2015) Effects of vitamin D supplementation on neuroplasticity in older adults: a double-blinded, placebo-controlled randomised trial. *Osteoporos Int* 26(1): 131-40.
137. Johnen B, Schott N (2018) Feasibility of a machine vs free weight strength training program and its effects on physical performance in nursing home residents: a pilot study. *Aging Clin Exp Res* 30(7): 819-828.
138. Romine PE, Kiely DK, Holt N, Percac-Lima S, Leveille S, et al. (2017) Task-Specific Fatigue Among Older Primary Care Patients. *J Aging Health* 29(2): 310-323.
139. Sions JM, Coyle PC, Velasco TO, Elliott JM, Hicks GE (2017) Multifidus Muscle Characteristics and Physical Function Among Older Adults With and Without Chronic Low Back Pain. *Arch Phys Med Rehabil* 98(1): 51-57.
140. Lange-Maia BS, Fitzpatrick SL, Janssen I, Dugan SA, Kravitz HM, et al. (2019) Factors Influencing Longitudinal Stair Climb Performance from Midlife to Early Late Life: The Study of Women's Health Across the Nation Chicago and Michigan Sites. *J Nutr Health Aging* 23(9): 821-828.
141. Orange ST, Marshall P, Madden LA, Vince RV (2019) Short-Term Training and Detraining Effects of Supervised vs. Unsupervised Resistance Exercise in Aging Adults. *J Strength Cond Res* 33(10): 2733-2742.
142. Unhjem R, den Hoven LTV, Nygård M, Hoff J, Wang E (2019) Functional Performance with Age: The Role of Long-Term Strength Training. *J Geriatr Phys Ther* 42(3): 115-122.
143. Katsoulis K, Mathur S, Amara CE (2021) Reliability of Lower Extremity Muscle Power and Functional Performance in Healthy, Older Women. *J Aging Res* 2021: 8817231.
144. Lindberg K, Lohne-Seiler H, Fosstveit SH, Sibayan EE, Fjeller JS, et al. (2022) Effectiveness of individualized training based on force-velocity profiling on physical function in older men. *Scand J Med Sci Sports* 32(6): 1013-1025.
145. Fosstveit SH, Løvold S, Kolnes T, Vårvik FT, Berntsen S, et al. (2023) Associations between Power Training-Induced Changes in Body Composition and Physical Function in Older Men: A Pre-Test-Post-Test Experimental Study. *Int J Environ Res Public Health* 20(22).
146. Becker C, Lord SR, Studenski SA, Warden SJ, Fielding RA, et al. (2015) Myostatin antibody (LY2495655) in older weak fallers: a proof-of-concept, randomised, phase 2 trial. *Lancet Diabetes Endocrinol* 3(12): 948-57.
147. Vogt L, Wieland K, Bach M, Himmelreich H, Banzer W (2008) Cognitive status and ambulatory rehabilitation outcome in geriatric patients. *J Rehabil Med* 40(10): 876-8.
148. Galvão DA, Taaffe DR (2005) Resistance exercise dosage in older adults: single- versus multiset effects on physical performance and body composition. *J Am Geriatr Soc* 53(12): 2090-7.
149. Eyigor S, Karapolat H, Durmaz B, Ibisoglu U, Cakir S (2009) A randomized controlled trial of Turkish folklore dance on the physical performance, balance, depression and quality of life in older women. *Arch Gerontol Geriatr* 48(1): 84-8.
150. Eyigor S, Karapolat H, Durmaz B (2007) Effects of a group-based exercise program on the physical performance, muscle strength and quality of life in older women. *Arch Gerontol Geriatr* 45(3): 259-71.
151. Judd DL, Thomas AC, Dayton MR, Stevens-Lapsley JE (2014) Strength and functional deficits in individuals with hip osteoarthritis compared to healthy, older adults. *Disabil Rehabil* 36(4): 307-12.
152. Storer TW, Woodhouse L, Magliano L, Singh AB, Dzekov C, et al. (2008) Changes in muscle mass, muscle strength, and power but not physical function are related to testosterone dose in healthy older men. *J Am Geriatr Soc* 56(11): 1991-9.
153. Chalé A, Cloutier GJ, Hau C, Phillips EM, Dallal GE, et al. (2013)

- Efficacy of whey protein supplementation on resistance exercise-induced changes in lean mass, muscle strength, and physical function in mobility-limited older adults. *J Gerontol A Biol Sci Med Sci* 68(6): 682-90.
154. Daly RM, Gianoudis J, Ross BD, O'Connell SL, Kruger M, et al. (2020) Effects of a multivitamin-fortified milk drink combined with exercise on functional performance, muscle strength, body composition, inflammation, and oxidative stress in middle-aged women: a 4-month, double-blind, placebo-controlled, randomized trial. *Am J Clin Nutr* 112(2): 427-446.
 155. Ke D, Wang D, Huang H, Hu X, Sasaki J, et al. (2022) Study of the Reliability of Field Test Methods for Physical Fitness in Children Aged 2-3 Years. *Int J Environ Res Public Health* 19(12).
 156. Lanzi S, Pousaz A, Calanca L, Mazzola L (2023) Time-course evolution of functional performance during a 3-month supervised exercise training program in patients with symptomatic peripheral artery disease. *Vasc Med* 28(5): 404-411.
 157. Molsted S, Eidemak I, Sorensen HT, Kristensen JH (2004) Five months of physical exercise in hemodialysis patients: effects on aerobic capacity, physical function and self-rated health. *Nephron Clin Pract* 96(3): c76-81.
 158. Storer TW, Casaburi R, Sawelson S, Kopple JD (2005) Endurance exercise training during haemodialysis improves strength, power, fatigability and physical performance in maintenance haemodialysis patients. *Nephrol Dial Transplant* 20(7): 1429-37.
 159. Zhang M, Upreti R, Martin D, Kalantar-Zadeh K, Kopple JD, et al. (2014) Relation between anxiety, depression, and physical activity and performance in maintenance hemodialysis patients. *J Ren Nutr* 24(4): 252-60.
 160. Dreher M, Sonntag F, Prettin S, Kabitz HJ, Windisch W, et al. (2008) Exercise in severe COPD: is walking different from stair-climbing? *Respir Med* 102(6): 912-8.
 161. Reddy S, Contreras CM, Singletary B, Bradford TM, Waldrop MG, et al. (2016) Timed Stair Climbing Is the Single Strongest Predictor of Perioperative Complications in Patients Undergoing Abdominal Surgery. *J Am Coll Surg* 222(4): 559-66.
 162. Basaria S, Collins L, Dillon EL, Orwoll K, Storer TW, et al. (2013) The safety, pharmacokinetics, and effects of LGD-4033, a novel nonsteroidal oral, selective androgen receptor modulator, in healthy young men. *J Gerontol A Biol Sci Med Sci* 68(1): 87-95.
 163. Baldwin JN, McKay MJ, Hiller CE, Moloney N, Nightingale EJ, et al. (2017) Relationship between physical performance and self-reported function in healthy individuals across the lifespan. *Musculoskelet Sci Pract* 30: 10-17.
 164. Baldwin JN, McKay MJ, Moloney N, Hiller CE, Nightingale EJ, et al. (2017) Reference values and factors associated with musculoskeletal symptoms in healthy adolescents and adults. *Musculoskelet Sci Pract* 29: 99-107.
 165. Mulla DM, Jarvis RS, Stephens A, Keir PJ, Maly MR, et al. (2018) The Effects of Lower Extremity Strengthening Delivered in the Workplace on Physical Function and Work-Related Outcomes Among Desk-Based Workers: A Randomized Controlled Trial. *J Occup Environ Med* 60(11): 1005-1014.
 166. Pollock M, Roa J, Benditt J, Celli B (1993) Estimation of ventilatory reserve by stair climbing. A study in patients with chronic airflow obstruction. *Chest* 104(5): 1378-83.
 167. Elbasan B, Tunali N, Duzgun I, Ozcelik U (2012) Effects of chest physiotherapy and aerobic exercise training on physical fitness in young children with cystic fibrosis. *Ital J Pediatr* 38: 2.
 168. Boreham CA, Wallace WF, Nevill A (2000) Training effects of accumulated daily stair-climbing exercise in previously sedentary young women. *Prev Med* 30(4): 277-81.
 169. Lafortuna CL, Agosti F, Proietti M, Adorni F, Sartorio A (2006) The combined effect of adiposity, fat distribution and age on cardiovascular risk factors and motor disability in a cohort of obese women (aged 18-83). *J Endocrinol Invest* 29(10): 905-12.
 170. Sartorio A, Agosti F, De Col A, Lafortuna CL (2006) Age- and gender-related variations of leg power output and body composition in severely obese children and adolescents. *J Endocrinol Invest* 29(1): 48-54.
 171. Devendra GP, Rane AA, Krasuski RA (2012) Provoked exercise desaturation in patent foramen ovale and impact of percutaneous closure. *JACC Cardiovasc Interv* 5(4): 416-9.
 172. Hetzler RK, Vogelpohl RE, Stickley CD, Kuramoto AN, Delaura MR, et al. (2010) Development of a modified Margaria-Kalamen anaerobic power test for American football athletes. *J Strength Cond Res* 24(4): 978-84.
 173. McKeon JL, Turner J, Kelly C, Dent A, Zimmerman PV (1986) The effect of inspiratory resistive training on exercise capacity in optimally treated patients with severe chronic airflow limitation. *Aust N Z J Med* 16(5): 648-52.
 174. Pate P, Tenholder ME, Griffin JP, Eastridge CE, Weiman DS (1996) Preoperative assessment of the high-risk patient for lung resection. *Ann Thorac Surg* 61(5): 1494-500.
 175. Brunelli A, Refai MA, Monteverde M, Borri A, Salati M et al. (2003) Predictors of exercise oxygen desaturation following major lung resection. *Eur J Cardiothorac Surg* 24(1): 145-8.
 176. Brunelli A, Monteverde M, Refai MA, Fianchini A (2004) Stair climbing test as a predictor of cardiopulmonary complications after pulmonary lobectomy in the elderly. *Ann Thorac Surg* 77(1): 266-70.
 177. Brunelli A, Monteverde M, Borri A, Salati M, Al Refai M, et al. (2003) Predicted versus observed maximum oxygen consumption early after lung resection. *Ann Thorac Surg* 76(2): 376-80.
 178. Brunelli A, Monteverde M, Salati M, Borri A, Al Refai M, et al. (2001) Stair-climbing test to evaluate maximum aerobic capacity early after lung resection. *Ann Thorac Surg* 72(5): 1705-10.
 179. Brunelli A, Pompili C, Berardi R, Mazzanti P, Onofri A, et al. (2012) Performance at preoperative stair-climbing test is associated with prognosis after pulmonary resection in stage I non-small cell lung cancer. *Ann Thorac Surg* 93(6): 1796-800.
 180. Brunelli A, Refai M, Xiumé F, Salati M, Sciarra V, et al. (2008) Performance at symptom-limited stair-climbing test is associated with increased cardiopulmonary complications, mortality, and costs after major lung resection. *Ann Thorac Surg* 86(1): 240-7; discussion 247-8.
 181. Brunelli A, Socci L, Refai M, Salati M, Xiumé F, et al. (2007) Quality of life before and after major lung resection for lung cancer: a prospective follow-up analysis. *Ann Thorac Surg* 84(2): 410-6.
 182. Brunelli A, Xiumé F, Refai M, Salati M, Sabbatini A, et al. (2010) Peak oxygen consumption measured during the stair-climbing test in lung resection candidates. *Respiration* 80(3): 207-11.
 183. Koegelenberg CF, Diacon AH, Irani S, Bolliger CT (2008) Stair climbing in the functional assessment of lung resection candidates. *Respiration* 75(4): 374-9.
 184. Pancieri MV, Cataneo DC, Montovani JC, Maria Cataneo AJ (2010) Comparison between actual and predicted postoperative stair-climbing test, walk test and spirometric values in patients undergoing lung resection. *Acta Cir Bras* 25(6): 535-40.
 185. Cataneo DC, Cataneo AJ (2007) Accuracy of the stair climbing test

- using maximal oxygen uptake as the gold standard. *J Bras Pneumol* 33(2): 128-33.
186. Bernasconi M, Koegelenberg CFN, Bolliger CT, Barnard BJ, Maree D, et al. (2012) Speed of ascent during stair climbing identifies operable lung resection candidates. *Respiration* 84(2): 117-22.
 187. Refai M, Brunelli A, Sabbatini A, Xiumè F, Pompili C, et al. (2014) Can maximal inspiratory and expiratory pressures during exercise predict complications in patients submitted to major lung resections? A prospective cohort study. *Eur J Cardiothorac Surg* 45(4): 665-69; discussion 669-70.
 188. Ito H, Masuda M, Yokose T, Yamada K, Nakayama H (2015) Outcomes of lobectomy in 'active' octogenarians with clinical stage I non-small-cell lung cancer. *Ann Thorac Cardiovasc Surg* 21(1): 24-30.
 189. Kubori Y, Matsuki R, Hotta A, Morisawa T, Tamaki A (2017) Comparison between stair-climbing test and six-minute walk test after lung resection using video-assisted thoracoscopic surgery lobectomy. *J Phys Ther Sci* 29(5): 902-904.
 190. Nakamura T, Sawabata N, Susaki Y, Muro S (2020) Desaturation during the stair-climbing test for patients who will undergo pulmonary resection: an indicator of postoperative complications. *Gen Thorac Cardiovasc Surg* 68(1): 49-56.
 191. Ozeki N, Nakamura S, Hakiri S, Iwano S, Fukui T, et al. (2021) Factors associated with changes in the 12-m stair-climbing time after lung lobectomy. *Gen Thorac Cardiovasc Surg* 69(2): 282-289.
 192. Dong J, Mao Y, Li J, He J (2017) Stair-Climbing Test Predicts Postoperative Cardiopulmonary Complications and Hospital Stay in Patients with Non-Small Cell Lung Cancer. *Med Sci Monit* 23: 1436-1441.
 193. Njøten KL, Espehaug B, Kvale G, Frisk B, Jørgensen M, et al. (2023) Relationship between exercise capacity and fatigue, dyspnea, and lung function in non-hospitalized patients with long COVID. *Physiol Rep* 11(22): e15850.
 194. Hellberg M, Höglund P, Svensson P, Abdulahi H, Clyne N (2017) Decline in measured glomerular filtration rate is associated with a decrease in endurance, strength, balance and fine motor skills. *Nephrology (Carlton)* 22(7): 513-519.
 195. Sérvio TC, Pereira RS, Cataneo DC (2012) Study on functional cardiorespiratory changes after laparoscopic Nissen fundoplication. *Acta Cir Bras* 27(7): 499-504.
 196. Arruda KA, Cataneo DC, Cataneo AJ (2013) Surgical risk tests related to cardiopulmonary postoperative complications: comparison between upper abdominal and thoracic surgery. *Acta Cir Bras* 28(6): 458-66.
 197. Khenafes TE, de Lima JF, Camargo Pereira RSD, Cataneo DC (2014) Cardiorespiratory evaluation in pre and post operative moments of laparoscopic cholecystectomy. *Acta Cir Bras* 29(6): 394-9.
 198. Cataneo DC, Kobayasi S, de Carvalho LR, Paccanaro RC, Maria Cataneo AJ (2010) Accuracy of six minute walk test, stair test and spirometry using maximal oxygen uptake as gold standard. *Acta Cir Bras* 25(2): 194-200.
 199. Coll F, Hill K, Burrows S, Watson C, Edgar D (2020) Modified Chester Step Test in a Healthy Adult Population: Measurement Properties and Development of a Regression Equation to Estimate Test Duration. *Phys Ther* 100(8): 1411-1418.
 200. Calavalle AR, Sisti D, Stocchi V, Rocchi MBL, Sal MD, et al. (2013) A simple method to analyze overall individual physical fitness in firefighters. *J Strength Cond Res* 27(3): 769-75.
 201. Oesch P, Jan Kool, Jansen B, Meyer K (2015) Functional Capacity Evaluation: Performance of Patients with Chronic Non-specific Low Back Pain Without Waddell Signs. *J Occup Rehabil* 25(2): 257-66.
 202. Honda H, Hayashi T, Miki T, Igaki M, Hatanaka Y, et al. (2016) Stair climbing/descending exercise for a short time decreases blood glucose levels after a meal in people with type 2 diabetes. *BMJ Open Diabetes Res Care* 4(1): e000232.
 203. Takaishi T, Hayashi T (2017) Stair ascending-descending exercise accelerates the decrease in postprandial hyperglycemia more efficiently than bicycle exercise. *BMJ Open Diabetes Res Care* 5(1): e000428.
 204. Moore J, Salmons H, Vinoskey C, Kressler J (2020) A single one-minute, comfortable paced, stair-climbing bout reduces postprandial glucose following a mixed meal. *Nutr Metab Cardiovasc Dis* 30(11): 1967-1972.
 205. Bartholomae E, Johnson Z, Moore J, Ward K, Kressler J (2018) Reducing Glycemic Indicators with Moderate Intensity Stepping of Varied, Short Durations in People with Pre-Diabetes. *J Sports Sci Med* 17(4): 680-685.
 206. Matsumoto K, Chen C, Oda Y, Nakagawa S, Hirotsu M, et al. (2021) The Effect of Brief Stair-Climbing on Divergent and Convergent Thinking. *Front Behav Neurosci* 15: 834097.
 207. Poppius H, Stenius B (1977) Exercise-induced asthma and doxantrazole. *Eur J Clin Pharmacol* 11(2): 107-9.
 208. Aveline C, Roux AL, Hetet HL, Bonnet F, Cognet F, et al. (2014) Pain and recovery after total knee arthroplasty: a 12-month follow-up after a prospective randomized study evaluating Nefopam and Ketamine for early rehabilitation. *Clin J Pain* 30(9): 749-54.
 209. Novoa NM, Rodríguez M, Gómez MT, Jiménez MF, Varela G, (2015) Fixed-altitude stair-climbing test replacing the conventional symptom-limited test. A pilot study. *Arch Bronconeumol* 51(6): 268-72.
 210. Margaria R, Aghemo P, Rovelli E (1966) Measurement of muscular power (anaerobic) in man. *J Appl Physiol* 21(5): 1662-4.
 211. Rutkove SB, Parker RA, Nardin RA, Connolly CE, Raynor EM, et al. (2002) A pilot randomized trial of oxandrolone in inclusion body myositis. *Neurology* 58(7): 1081-7.
 212. Bruun-Olsen V, Heiberg KE, Mengshoel AM (2009) Continuous passive motion as an adjunct to active exercises in early rehabilitation following total knee arthroplasty - a randomized controlled trial. *Disabil Rehabil* 31(4): 277-83.
 213. Hsieh RL, Lee WC, Lo MT, Liao WC (2012) Short-term effects of 890-nanometer radiation on pain, physical activity, and postural stability in patients with knee osteoarthritis: a double-blind, randomized, placebo-controlled study. *Arch Phys Med Rehabil* 93(5): 757-64.
 214. Rigamonti AE, Scacchi M, Cicolini S, Caroli D, De Col A, et al. (2020) Effects of a 3-Week In-Hospital Body Weight Reduction Program on Cardiovascular Risk Factors, Muscle Performance, and Fatigue: A Retrospective Study in a Population of Obese Adults with or without Metabolic Syndrome. *Nutrients* 12(5): 1495.
 215. Lafortuna CL, Sartorio A, Sartorio A, Marinone PG, Agosti F (2004) The relationship between body composition and muscle power output in men and women with obesity. *J Endocrinol Invest* 27(9): 854-61.



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/JPFMTS.2025.11.555810](https://doi.org/10.19080/JPFMTS.2025.11.555810)

**Your next submission with Juniper Publishers
will reach you the below assets**

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission
<https://juniperpublishers.com/online-submission.php>