



Natural Antimicrobials and Antidiabetic Plant Extracts, Singly and in Combination, Potential Source of Novel Antimicrobials and Hypoglycemic Agents



Jagessar RC*

Department of Chemistry, University of Guyana, Turkey

Received Date: January 25, 2019; Published Date: March 15, 2019

*Corresponding author: Jagessar RC, Department of Chemistry, University of Guyana, Guyana, Turkeyen Campus, Greater Georgetown, Turkey

Abstract

Research in the design and synthesis of antimicrobials must be continued on this planet, considering that bacteria develop antimicrobial resistance to antimicrobial drugs. In addition, many synthetic drugs have side effects when administered and the cost of synthesizing drugs is an expensive endeavor and is non realized in developing country like Guyana. Guyana, however, vast and diverse flora have been shown to possess potent and selective antimicrobial activity that can lead to the further development of herbal medicines with new applications. In addition, our plants have been shown to possess striking hypoglycemic effect in guinea pig model experiments. Further research is necessary to continue with the development of novel herbal medicines, singly or in combination. In addition, the isolation and purification of the known and unknown active natural products that may be responsible for the medicinal activity, singly or in combination. In addition, a modulation of their structure/activity relationship. Unknown natural products would form the basis of novel drug discovery

Keywords: Antimicrobials; Hypoglycaemic Agents; Guyana; Flora; Novel Drug Discovery.

Introduction

Research in the design and synthesis of antimicrobials will continue to be problematic on our planet, considering that bacteria and fungi develop resistance to antimicrobials over a period of time [1-5]. This results from indiscriminate use of commercial antimicrobial drugs for the treatment of infectious diseases and the current global antibiotic resistance. Many synthetic drugs have several adverse side effects which are usually irreversible when administered and the cost of synthesizing drugs in most cases is an expensive endeavor [1-5] and is non-realistic for a developing country like Guyana. Plants have a long therapeutic history over thousands of years and still considered to be promising source of medicine in the traditional health care system [6]. Plants also have a wide variety of secondary metabolites some of which are antimicrobial [7-9]. Crude plants extracts have also demonstrated antimicrobial activity [10-12].

Guyana flora is richly bio diversified and it's organic and aqueous extracts have been shown to possess potent and selective antimicrobial activity to date, compared with standard antibiotics: penicillin, nystatin and ampicillin [13-17]. Also, recently antidiabetic activities [18]. With the view of avoiding clinical trials, an expensive endeavor for a Third World country, we have shown that Guyana's flora has strong potent antibacterial and antifungal activ

ity. We have over the years, investigated the antimicrobial activity and selectivity of a wide range of plants and fruits against human pathogens such as *E. coli*, *S. aureus*, *Bacillus subtilis*, *K. pneumonia*, *P. aeruginosa* and *C. albicans*. Extracts include those from *Brassica rapa chinensis*, *Artocarpus altilis*, *Solanum melongena* fruits, leaves of *Moringa oleifera* [19], passion fruit (*Passiflora edulis*) [20], *Phyllanthus acidus*, *Sphagneticola trilobata* and *Dolioscarpus dentatus's* [21] bark amongst others. Antimicrobial potency and selectivity have been comparable to standard antimicrobials such as penicillin and nystatin. against pathogens. Also, the antimicrobial potency and selectivity can be modulated in the presence and absence of transition metal salts such as zinc and copper. For example, a study to investigate the Antimicrobial Activity of the Uncombined and Combined Aqueous Extract of *Phyllanthus acidus*, *Sphagneticola trilobata* Leaves and *Dolioscarpus dentatus* Bark against Selective Pathogenic microorganisms in the absence and presence of Zn^{2+} cations revealed that the highest AZOI of 165.05 mm^2 was induced by *P. acidus* extract against *E. coli*. In the absence of Zn^{2+} , zero AZOI was observed for the combined extract of *P. acidus* + *S. trilobata*, *S. trilobata* + *D. dentatus* and *P. acidus* + *D. dentatus* against *S. aureus* and *K. pneumoniae*. The combined plant extract, without Zn^{2+} , seems to induce a higher AZOI against *E.*

coli, *K. pneumoniae*, *C. albicans* and *P. aeruginosa* in comparison with the individual plant extract. As an example, the combined plant extract of *S. trilobata* and *D. dentatus* induces AZOI of 122.66 mm², whereas *S. trilobata* induces AZOI of 117.79 mm². For the combined plant extract with Zn²⁺, a lower AZOI was induced, compared with the individual plant extract. Selective antimicrobial activity were observed for the uncombined and combined extracts, with and without Zn²⁺ against some of the pathogens. For example, *P. acidus* aqueous extract showed AZOI of 165.1 mm² against *E. coli*, whereas *S. trilobata* showed AZOI of 67.17 mm² against *E. coli* i.e., a selectivity ratio of 2.5 vs *E. coli* with respect to the above two extracts.

Antimicrobial activities of plant extracts at varying concentrations of 0.025g/ml, 0.05g/ml and 0.1g/ml have been investigated mainly using the Disc Diffusion Assay. Here the diameter and the computed Area of Zone of Inhibition have been used as a gauge of the plant's antimicrobial activity. For example, AZOI of 209.34 mm² was induced by the CH₃CH₂OH extract of *Brassica rapa chinensis* against *E. coli* at a concentration of 0.025g/ml whereas an AZOI of 12.56 mm² was induced by *Brassica rapa chinensis* against *Bacillus subtilis* at a concentration of 0.025g/ml. Antidiabetic experiments using plant extracts have been conducted using guinea pigs model experiments.

Conclusion

Plant extracts have indeed shown to possess potent and selective antimicrobial activity against human pathogens. Also, striking hypoglycemic effect. Research should continue in the development of herbal medicines to combat current and new ailments. Also, an isolation of the active principle or natural products on route to novel drug discovery.

References

1. Wilms LR (2009) Guide to Drugs in Canada. Leo Paper Products Ltd. China (3rd edn): 114-117.
2. Macor JE (2010) Annual reports in Medicinal Chemistry, sponsored by the Division of Medicinal Chemistry of the American Chemical Society, Elsevier Inc., 45: 295-311.
3. Wood A (2008) Topics in Drug design and discovery. Annual Reports in Medicinal Chemistry, Elsevier Inc. 41: 353-409.
4. Bonner J (2009) Filling the Antibiotic Gap. Chemistry World, Royal Society of Chemistry 6(8): 16.
5. Kelland K (2013) Antibiotic Resistance Poses Catastrophic Threat To Medicine. Huffington Post 1-3.
6. Shahid W, Durrani R, Iram S, Durrani M, Khan F (2013) Antibacterial activity *in-vitro* of medicinal plants. Sky Journal of Microbiology Research 1(2): 5-21.
7. Rijo P, Faustino C, Simoes MF (2013) Antimicrobial natural products from *Plectranthus* plants. Microbial pathogens and strategies for combating them: science, technology and education 922-931.
8. Arif T, Bhosale JD, Kumar N, Mandal TK, Bendre RS (2009) Natural Products-antifungal agents derived from plants. J Asian Nat Prod Res 11 (7): 621-638.
9. Melgarejo M, Mollined P, Castro JV (2006) Antibacterial activity of four Natural Products from a Bolivian Highland Plant. Revista Boliviana De Quimica 23(1): 40-43.
10. Saswati R, Choudhury M, Paul S (2013) *In-Vitro* Antibacterial Activity of *Alocasia Decipiens* Schott. International Journal of Pharmacy and Pharmaceutical Sciences 5(1).
11. Aarati N, Ranganath N, Soumya G, Kishore B, Mithun K (2011) Evaluation of Antibacterial and Anticandidal Efficiency of Aqueous and Alcoholic Extract of Neem *Azadirachta indica*. International Journal of Research and Pharmaceutical.
12. Meenal K, Khadabadi S, Saboo S, Ghorpade D, Modi A (2010) *In-Vitro* Antimicrobial Activity of The Crude Extracts Of *Colocasia Esculenta* Leaves (Araceae). International Journal of Pharmacy and Pharmaceutical Sciences 1(8).
13. Jagessar RC, Mohamed N (2010) Antimicrobial activity of selected plants extracts from Guyana's flora. Journal of Pure and Applied Microbiology 4(2): 533-540.
14. Jagessar RC, Allen R (2011) Antimicrobial Potency of the Aqueous Extract of leaves of *Terminalia catappa*. Academic Research International 362-371.
15. Jagessar RC, Mars A, Gomathigayam S (2011) Selective Antimicrobial properties of Leaf extract of *Samanea Saman* against *Candida albicans*, *Staphylococcus aureus* and *Escherichia coli* using several microbial techniques. Journal of American Science 7(3): 108-119.
16. Jagessar RC, Mars A, Gomes G (2009) Leaf extract of *Smilax schomburgkiana* exhibit selective antimicrobial properties against pathogenic microorganisms. Life Science Journal 6(1): 76-83.
17. Jagessar RC, Mohammed A, Gomes G (2008) An evaluation of the antibacterial and antifungal activity of leaf extracts of *Momordica Charantia* against *Candida albicans*, *Staphylococcus aureus* and *Escherichia Coli*. Nature and Science 6(1): 1-14.
18. Jagessar RC, Rodriques A, Prasad K, Husain A, Kanhai V, et al. (2018) An investigation of the hypoglycemic effect of the aqueous extract of the fruits of *Psidium Guajava*, *Averrhoa Bilimbi* and the peel of *Tamarindus indica* in Normoglycemic guinea pigs. J Pharmaceutics Drug Delivery Research 7(4): 77-101.
19. Jagessar RC, Ramchartar N, Spencer O (2015) A Review of the Antimicrobial Activity of Various Solvent Type extracts from Fruits and Edible Plants, in Fruit and Pomace Extracts: Biological Activity, Potential Applications and Beneficial Health Effects. Nova Science Publisher.
20. Jagessar RC, Hafeez, Chichester, Crepaul Y (2017) Antimicrobial Activity of the Ethanolic and Aqueous Extract of Passion Fruit (*Passiflora edulis Sims*), in the absence and presence of Zn(OAc)₂.2H₂O. World Journal of Pharmacy and Pharmaceutical Sciences. 6(9): 230-246.
21. Jagessar RC, Hope S (2016) Antimicrobial Activity of the Uncombined and Combined Aqueous Extract of *Phyllanthus acidus*, *Sphagneticola Trilobata* Leaves and *Dolioscarpus Dentatus* Bark against Selective Pathogenic Microorganisms in the absence and presence of Zn²⁺ cations. World Journal of Pharmacy and Pharmaceutical Sciences 5(8): 58-71.



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/NAPDD.2019.05.555652](https://doi.org/10.19080/NAPDD.2019.05.555652)

**Your next submission with Juniper Publishers
will reach you the below assets**

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission

<https://juniperpublishers.com/online-submission.php>