



Accumulation of Heavy Metal and Microbial Contamination of Smoked Catfish (*Clarias gariepinus*)



Karimu Olajubu Ayotunde¹ and Victor- Osanyinlusi Remi^{2*}

¹Principal lecturer, Department of pharmaceutical Technology Rufus Giwa Polytechnic Owo Ondo State, Nigeria

²Technologist, Department of pharmaceutical Technology Rufus Giwa Polytechnic Owo Ondo State Nigeria

Submission: January 06, 2025; **Published:** January 31, 2025

***Corresponding author:** Victor - Osanyinlusi Remi, Technologist Department of pharmaceutical Technology Rufus Giwa Polytechnic Owo Ondo State Nigeria

Abstract

Recently, over 70% of fish are smoked as a means of preservation. Smoking is an ancient processing method that remains widely practiced in Nigeria today. This study investigates the levels of heavy metal accumulation and microbial loads in smoked catfish obtained from two different catfish farms to determine the safety of smoked catfish sold in Owo during the research period. Samples were collected from two farms (Farm 1 and Farm 2) in Owo, located in Owo Local Government Area, Ondo State, Nigeria. The identified microorganisms included *Streptococcus* spp., *Staphylococcus aureus*, *Bacillus* spp., *Klebsiella* spp., *Pseudomonas aeruginosa*, and *Escherichia coli*. The microbial counts for samples A and B were as follows: *Streptococcus* spp. (90.0 and 60.0), *Staphylococcus aureus* (160.0 and 170.0), *Bacillus* spp. (230.0 and 215.0), *Klebsiella* spp. (110.0 and 120.0), *Pseudomonas aeruginosa* (15.0 and 10.0), and *Escherichia coli* (2.0 and 1.0). For heavy metals, the concentrations were Cu (0.001 and 0.000), Cd (0.222 and 0.002), Cr (0.840 and 0.670), Mn (2.33 and 1.99), and Zn (132.020 and 127.001). The highest microbial count was observed in *Bacillus* spp. from Sample A (230.0) and Sample B (215.0), while the lowest was found in *Escherichia coli* from Sample B (1.0) and Sample A (2.0). Among the heavy metals, zinc was the most abundant in both samples, with Sample A (132.020) having a higher concentration than Sample B (127.001). Copper was the least abundant, being almost undetectable in Sample A (0.001) and entirely undetectable in Sample B (0.000). The study revealed variations in microbial and heavy metal contamination levels between catfish farms. It highlights the need for regulatory authorities to enforce moisture control measures and implement strategies to reduce human activities that may lead to bacterial growth and heavy metal contamination in smoked catfish products.

Abbreviations: Bacteria; Microorganisms; Contamination; Heavy metals; Smoked fish; Cat fish and Farms

Introduction

Fish, belonging to the phylum Chordata and class Pisces, is predominantly aquatic [1]. It serves as an excellent source of essential minerals needed for proper body function [2]. Globally, fish is a significant source of protein, complemented by fats and oils beneficial for human nourishment. Catfish (*Clarias gariepinus*), widely consumed across socio-economic groups in Nigeria, serves as a vital source of animal protein, rivaling meat consumption. This freshwater species is highly regarded for its unique flavor, texture, and wide cultivation in ponds across the country. Fish also provides vital proteins, vitamins, and minerals, essential for both infant and adult diets [3].

However, fish requires careful handling due to its susceptibility to spoilage in tropical temperatures, which accelerates bacterial, enzymatic, and oxidative processes [4]. In Nigeria, poor handling

practices lead to significant post-harvest losses (30-50%). These losses can be minimized through appropriate processing and preservation methods, such as salting, drying, canning, and smoking [5].

Smoking, a widely adopted method in Nigeria, involves applying heat to reduce water content and inhibit bacterial and enzymatic activity [6]. Despite its effectiveness, improper smoking methods and lack of hygiene often lead to high microbial loads and exposure to contaminants like dust and heavy metals. Studies [7] show that poorly smoked fish can harbor high microbial levels, and open-flame smoking may introduce cancer-promoting compounds into the body.

Heavy metals, characterized by high densities and atomic weights, include essential elements like iron, cobalt, and zinc,

which are beneficial in small amounts but toxic in excess. Toxic heavy metals like cadmium, mercury, and lead are hazardous and often find their way into aquatic environments through activities like mining, agriculture, and industrial waste disposal. These pollutants are absorbed by catfish, posing potential health risks to consumers [8].

This study aims to evaluate the levels of heavy metals and microbial loads in smoked catfish sourced from two farms in Owo, Ondo State, Nigeria.

Materials And Methods

Sample Collection

Two catfish samples were obtained from two separate fish farms in Owo, Owo Local Government Area, Ondo State, Nigeria. The samples were transported in water to the Aquaculture Department at Rufus Giwa Polytechnic, where they were identified using standard procedures.

Sample Preparation

The catfish samples were washed thoroughly under running water and smoked without adding salt until fully dried. The smoked fish were separately crushed using a mortar and pestle, then milled into powder using a blender. The powdered samples were stored in airtight containers for laboratory analysis.

Materials

The materials used included HCl, H₂SO₄, a weighing balance, filter paper, heating mantle, crucible, thread, beaker, conical flask, distilled water, reagent bottles, chloroform, water bath, acetic acid, pipette, muffle furnace, and other necessary laboratory equipment. All chemicals utilized were of analytical grade.

Determination of Heavy Metal Composition

Ashing

Approximately 20 g of each powdered sample was placed in a crucible and ashed in a muffle furnace at 550 °C for 4 hours. The ashed samples were then removed and stored in a desiccator to prevent moisture absorption.

Digestion

The ashed samples were mixed with Nitric acid (HNO₃) and Hydrochloric acid (HCl) in a 1:3 ratio. The mixture was diluted to 100 ml with distilled water in a measuring cylinder, transferred into a beaker, filtered, and stored in sample bottles at room temperature for Atomic Absorption Spectrophotometer (AAS) analysis.

Atomic Absorption Spectrophotometer (AAS) Analysis

The concentrations of heavy metals, including cadmium, manganese, copper, zinc, and chromium, were determined in

the samples using Atomic Absorption Spectrophotometer (AAS Model: 2000). The ash content and heavy metal analysis followed the AOAC [9] method.

Microbial Analysis

Bacteriological Test

A standard plate count method was employed for the microbial analysis. Ten grams of each catfish sample were aseptically weighed and homogenized in 90 ml of sterile peptone water. Serial dilutions were prepared by mixing 1.0 ml of the suspension with 9.0 ml of sterile peptone water, producing a 10⁻¹ dilution. Further dilutions were made up to 10⁻², and the diluted samples were spread onto nutrient agar plates to determine the total viable count.

Preparation of Agar Plates

Petri dishes were sterilized in a hot air oven at 160 °C for 1 hour. Nutrient agar was prepared by dissolving 0.6 g of agar in 100 ml of distilled water and sterilizing it in a microwave at 121 °C for 15 minutes. The agar was allowed to cool to 45 °C before use. All microbial analyses adhered to the procedures outlined by AOAC [10].

Results and Discussion

Results

Discussion

The heavy metal contamination levels in the smoked catfish samples are summarized in (Table 4) above. Zinc was the most abundant metal in both samples, with Sample A showing the highest concentration (132.020). Similarly, Sample B contained 127.001 of zinc. This aligns with findings by Ukulu et al. [11], who reported a high zinc concentration (135.000) in catfish. The zinc levels in this study exceed the WHO [12] and FAO [13] recommended limit of 99.4 mg/kg (Table 1). Zinc plays a crucial role in cell growth, division, and wound healing, but excessive intake can lead to adverse effects such as nausea, diarrhea, metallic taste, vomiting, kidney damage, and stomach issues.

Manganese (Mn) was also detected in both samples, with Sample A showing a higher concentration (2.33) compared to Sample B (1.99). These values are below the WHO (5.00 mg/kg) and FAO (4.00 mg/kg) limits for manganese in food. This indicates that the manganese levels in smoked catfish are insufficient to meet the body's requirements unless supplemented by other manganese-rich sources. Manganese is vital for iron metabolism, particularly in hemoglobin formation (Table 2). These results can be compared to values reported by Igwemmar et al. [14] for catfish (0.07 mg/kg), croaker (0.05 mg/kg), and tilapia (0.40 mg/kg), as well as those reported by Remi (2024) for roasted and fresh grass-cutter meat (0.111-0.114 mg/kg).

Table 1: Showing the Samples and Location of Collection.

Sample	Sample	Collection Point	Cold Room	State
Local Name	Botanical Name			
Cat fish	Clarias gariepinus	Owo	Farm (1)	Ondo

Table 2: Showing the Samples and Location of Collection.

Sample	Sample	Collection Point	Cold Room	State
Local Name	Botanical Name			
Cat fish	Clarias gariepinus	Owo	Farm (2)	Ondo

Table 3: Showing Length (Cm and Inches) And Weigh (Kg And G) Of The Fish Samples.

Fish Sample	L(Cm)	L (Inches)	Wt(G)	Wt (Kg)	Location
A (Cat fish)	30.5	12.007	1000	1	Farm 1
B (Cat fish)	30.70	12.087	1000	1	Farm 2

Table 4: Results of Heavy Metal Composition of the fish Samples (mg/kg).

Fish samples	Cu	Cd	Cr	Mn	Zn
A (Cat fish 1)	0.001	0.222	0.84	2.33	132.02
B (Cat fish 2)	0	0.002	0.67	1.99	127.001
WHO	4.00	0.05	0.05	5.00	99.4
Limit (mg/kg) (1993)					
FAO limit (mg/kg) (1999)	4.00	0.02	0.02	4.00	99.4

Table 5: Results of microbial load of catfish samples (cfu/g).

Parameter	Sample A	Sample B
Streptococcus spp	90	60
Staphylococcus aureus	160	170
Bacillus subtilis	230	215
Klebsiella spp	110	120
Pseudomonas aeruginosa	15	10
Escherichia coli	2	1

Chromium (Cr) levels in both samples were found to be high, with Sample A at 0.84 and Sample B at 0.67. These values exceed the limits recommended by WHO (0.05 mg/kg) and FAO (0.02 mg/kg). Kuton et al. [15] reported lower chromium levels (0.11 and 0.054) in the liver and intestine of *Malapterurus electricus*. Excessive chromium intake can reduce insulin sensitivity (Table 3), damage the liver and kidneys, and interfere with the efficacy of drugs like anti-inflammatories and paracetamol [16].

Cadmium concentrations were the lowest among the metals studied, with Sample B showing 0.002 and Sample A 0.222. Both values are below the standard limits of WHO (0.05 mg/kg) and FAO (0.02 mg/kg). Chima et al. [17] reported the absence of cadmium in some fish samples, while Ukulu et al. [11] found similar levels of 0.221 and 0.015 in catfish from Ajudabo, with undetectable levels in other samples. Cadmium, though present in low amounts here, is highly toxic and can cause kidney damage and bone demineralization at higher concentrations (Table 4).

Copper (Cu) was detected only in Sample A (0.001), while it was undetectable in Sample B. Alinnor & Obiji [18] reported significantly higher copper levels in fish, ranging from 1.247 to 8.00 ppm. The copper levels in this study can also be compared to the 0.199 and 0.212 ppm values for roasted and fresh grass-cutter meat, as reported by Remi [19]. The copper concentrations here were well below the WHO and FAO limit of 4.00 mg/kg. Copper is an essential trace element that supports enzymatic activity and overall metabolic functions.

Discussion

The microbial loads (CFU/g) of the smoked catfish samples are detailed in (Table 5). The bacterial isolates identified from the samples collected from the two catfish farms included *Streptococcus* spp, *Staphylococcus aureus*, *Bacillus* spp, *Klebsiella* spp, *Pseudomonas aeruginosa*, and *Escherichia coli*. The microbial loads in Sample A and Sample B were as follows: *Streptococcus*

spp (90.0 and 60.0), *Staphylococcus aureus* (160.0 and 170.0), *Bacillus* spp (230.0 and 215.0), *Klebsiella* spp (110.0 and 120.0), *Pseudomonas aeruginosa* (15.0 and 10.0), and *Escherichia coli* (2.0 and 1.0).

Among the isolates, *Bacillus* spp had the highest load (230.0 and 215.0) in both samples, while *Escherichia coli* had the lowest (2.0 and 1.0), followed by *Pseudomonas aeruginosa* (15.0 and 10.0). These findings are consistent with those of Ariyo & Obire [20], who reported low values of *Escherichia coli* and *Pseudomonas aeruginosa* in catfish samples. Similarly, Ayuba et al. [21] identified the presence of *Escherichia coli* and *Staphylococcus aureus* in smoked-dried sardine, corroborating the results of this study.

The detection of *Staphylococcus aureus* and *Escherichia coli* in the smoked catfish samples is consistent with Martin (1994), who noted that these microorganisms are commonly associated with smoked fish. *Staphylococcus aureus*, a bacterium frequently found on human skin and mucous membranes, is a leading cause of various human diseases. As Ayuba et al. [21] also noted, *Staphylococcus aureus* can cause both superficial and systemic infections, including boils, pneumonia, and bacteremia. According to Salan et al. [22] and Eyo [4], *Staphylococcus aureus* is resistant to heat, drying, and radiation, and produces toxins that can lead to food poisoning, septic shock, and toxic shock syndrome.

Escherichia coli, often linked to improper fish handling and fecal contamination [23], can cause diarrhea, kidney damage, and urinary tract infections [22] & [4]. The presence of *Staphylococcus aureus* and *Escherichia coli* in these samples poses risks of food poisoning and foodborne diseases, particularly if the contaminated fish is consumed.

The study also found *Pseudomonas* spp, an opportunistic pathogen associated with food spoilage and infections, in both smoked catfish samples. This contrasts with Olayemi et al. [24], who reported the absence of *Pseudomonas* spp in catfish samples from Ibadan. Similar findings have been reported by Adegunloye & Sanusi [25], Daramola et al. [26], and Oku & Amakoromo [27], who identified *Bacillus* spp and *Pseudomonas* spp in smoked fish samples from Tombia and Swale markets.

The presence of bacteria in the smoked catfish samples can be attributed to residual moisture (despite drying) and elevated temperatures that promote bacterial growth [28]. Additionally, *Bacillus* spp are known to produce endospores, allowing them to survive harsh environmental conditions.

Notably, significant differences were observed in the loads of *Staphylococcus aureus* and *Klebsiella* spp between the two samples, with values of 160.0 and 170.0 for *Staphylococcus aureus* and 110.0 and 120.0 for *Klebsiella* spp in Samples A and B, respectively. The highest load of *Streptococcus* spp was recorded in Sample A (90.0), compared to Sample B (60.0) [29].

Conclusion And Recommendation

Conclusion

In summary, the levels of heavy metal and microbial contamination in smoked catfish products vary between different fish farms. This contamination is attributed to factors such as inadequate smoking processes, poor personal hygiene of handlers, unsanitary smoking environments, improper storage, inadequate packaging, and various human activities [30]. These findings suggest that the two smoked catfish samples examined were contaminated not only during the handling process but also at the fish farm level due to the conditions mentioned.

Recommendation

Dehydration: Ensure complete dehydration of smoked catfish to inhibit bacterial growth caused by residual moisture [31].

Waste Management: Monitor and regulate human activities such as improper waste disposal, industrial discharges, and agricultural practices to reduce or eliminate heavy metal contamination in aquatic environments.

Collaborative Efforts: Regulatory bodies like the National Agency for Food and Drug Administration and Control (NAFDAC), the World Health Organization (WHO), and the Food and Agriculture Organization (FAO) should collaborate to establish and enforce moisture content standards for smoked catfish to minimize microbial growth.

Proper Drying: Catfish should be adequately dried before consumption, as heat can significantly reduce microbial contamination.

Public Awareness: Educate stakeholders, including fish farmers and consumers, on the importance of proper handling, smoking, and storage techniques to ensure food safety.

References

1. Salihu-Lasisi M, Akpabio CJ, Ogunsola MO (2013) Comparative nutritional studies on fresh and smoked *Clarias gariepinus* (Catfish) and *Tilapia nilotica* (Tilapia) fishes. *European Journal of Experimental Biology* 3(5):183-185.
2. Terech-Majewska E, Pajdak J, Siwicki AK (2016) Water As A Source Of Macronutrients And Micronutrients For Fish, With Special Emphasis On The Nutritional Requirements Of Two Fish Species: The Common Carp (*Cyprinus Carpio*) And The Rainbow Trout (*Oncorhynchus Mykiss*).
3. Abdullahi SA, Abolude DS, Ega RA (2001) Nutrient Quality of four Oven Dried Freshwater Catfish in Northern Nigeria. *J Trop Biosci* pp: 70-76.
4. Eyo AA (2001) Fish processing Technology in the Tropics. National Institute for Freshwater Fisheries Research (NIFFR), New Bussa, 37-39.
5. Bate EC, Bendall JR (2010) Changes in fish muscle after death. *British Medical Bulletin* (12): 2305.
6. Kumolu-Johnson CA, Aladetohun NF, Ndimele PE (2010) The effects of smoking on the Nutritional Qualities and Shelf-Life of *Clarias gariepinus* (BURCHELL 1822). *Afr J Biotechnol* 9(1): 73-76.

7. Abolagba OJ, Iyeru OA (1998) Study of Insect Pest Infecting Traditionally Processed Fish Sold in Benin City Metropolis, Nigeria. *Nig J Applied Sci* 16: 25-29.
8. Muradoglu FM, Gundogdu S, Ercisli T, Encu F, Balta HZ, et al. (2015) Cadmium toxicity affects chlorophyll II and b content, antioxidant enzyme activities and mineral nutrient accumulation in strawberry. *Biol Res* 48(1): 11.
9. AOAC (2010) Association of Official Analytical Chemists, (15th edition). Virginia pp.1298.
10. AOAC (2000) Enzymes and Enzyme products as Quality Indices. Seafood Enzymes. Harrod NF and Simpsn BK, (Eds) Marcel Dekker Inc New York, Basel, USA pp 337-363.
11. Ukulu HS, Balogun MA, Sawere BT, Chidi HO (2018) Analysis of Heavy Metals in the Catfish Species: *Clarias gariepinus*. *Journal of Chemical Society of Nigeria* 43(4).
12. WHO (1993) Heavy Metals- Environmental aspect-Environment Health Criteria, No 85, Geneva, Switzerland.
13. FAO Fisheries and Aquaculture (1999) Globalisation and Fisheries: Proceedings of an OECD-FAO Workshop Organization for Economic Co-operation and Development, OECD Publishing pp. 56.
14. Igwemmar NC, Kolawole NC, Odunoku SC (2013) Heavy Metal Concentration in Fish Species Sold In Gwagwalada Market, Abuja. (*IJSR*) 2(11): 7-11.
15. Kuton MP, Ayanda IO, Uzoalu IA, Akhiromen DI, George A, et al. (2021) Studies on heavy metals and fish health indicators in *Malapterurus electricus* from Lekki Lagoon, Lagos, Nigeria. *Vet Anim Sci* 12: 100169.
16. Aslam S, Yousafzai AM (2017) Chromium toxicity in fish: A review article. *Journal of Entomology and Zoology Studies*, 5(3): 1483-1488.
17. Nnodum CF, Yusuf KA, Adeniji CA (2017) Heavy Metals Contamination And Human Health Risk From Selected Fish In Lagos Nigeria. *Journal Of Research And Review In Science* 4: 231-237.
18. Alinnor IJ, Obiji IA (2010) Assessment of trace metal composition in fish samples from Nworie river. *Pakistan Journal of Nutrition* 9(1): 81-85.
19. Remi V-O (2024) Concentration of Nutrients Present in Fresh and Roasted Bush Meat (Grass-cutter) (*Thryonomys Swiderianus*). *Acad Res J Biotech* 12(1): 1-9.
20. Ariyo Adenike Bosede, Obire Omokaro (2021) Microbial Flora of Smoke-Dried Catfish (*Clarias Gariepinus*) Processed and Sold In Some Abattoirs In Bayelsa And Rivers States. *African Journal of Environment and Natural Science Research* 4(4): 123-133.
21. Ayuba VO, Alhassan MI, Jimmy UU (2013) Assessment of the microbial load of smoked sardine (*Sardina pilchardus*, Walbaum, 1792) sold in Makurdi markets. *Int J Curr Microbiol App Sci* 2(9): 277-287.
22. Salán OE, Juliana AG, Marilia O (2006) Use of Smoking to Add Value to Salmoned Trout. *Braz Arch Biol Technol* 49(1): 57-62.
23. Talaro KP (2009) Foundations' in Microbiology, 7th Edition. McGraw-Hill Companies. Avenue of the Americas, New York Pp 595-596.
24. Olayemi FF, Raji AO, Adedayo MR (2012) Microbiological quality of catfish (*Clarias gariepinus*) smoked with Nigerian Stored Products Research Institute (NSPRI) developed smoking kiln. *Int Res J Microbiol* 3(13): 426-430.
25. Adegunloye DV, Sanusi AI (2019) Microbiota of catfish (*Clarias glariiepinus*) tissues harvested from vials polluted with soil from the e-wastes dumpsite. *International Journal of Fisheries and Aquaculture* 11(5): 104-111.
26. Daramola JA, Alao FO, Adeniyi AE (2020) Estimation of Bacteria and Fungi in Smoked Catfish (*Clarias gariepinus*) available In OTA Markets. *Journal of Research in Forestry, Wildlife and Environment* 12(2): 65-73.
27. Oku I, Amakoromo E (2013) Microflora of fresh and smoke-dried fish in Yenogoa metropolis, Nigeria. *African Journal of Microbiological Research* 7(35): 4451-4456.
28. Ibrahim BU, Baba J, Sheshi MS (2014) Isolation and Identification of Bacteria Associated with Fresh and Smoked Fish (*Clarias gariepinus*) In Minna Metropolis, Niger State Nigeria. *Journal of Applied & Environmental Microbiology* 2(3): 81-85.
29. Ames G, Clucas I, Paul SS (1991) Post-Harvest Losses of Fish in the Tropics. *Natural Resources Institute*.
30. Karube I, Marouka H, Suzuki S, Watanabe E, Toyana K (1984) Determination of Fish Freshness with an Enzyme Sensor System. *Journal of Agriculture and Food Chemistry* 32: 314-319.
31. Olayemi FF, Adedayo MR, Bamishaiye EI, Awagu EF (2011) Proximate composition of catfish (*Clarias gariepinus*) smoked in Nigerian stored products research institute (NSPRI): Developed kiln. *Int J Fisheries and Aquaculture* 3(5): 96-98.



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/NFSIJ.2025.14.555879](https://doi.org/10.19080/NFSIJ.2025.14.555879)

**Your next submission with Juniper Publishers
will reach you the below assets**

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission
<https://juniperpublishers.com/online-submission.php>