

Case Report

Volume 5 Issue 2 - July 2017

DOI: 10.19080/OAJNN.2017.05.555659

Open Access J Neurol Neurosurg

Copyright © All rights are reserved by Muffaq lashhab

Extradural Spinal Meningioma in a Nine-Year-Old Girl: a Case Report and Review of the Literature



Hisham Ben Khayal¹, Abdulsalam Abograra² and Muffaq lashhab^{3*}

¹Associate Professor of Neurosurgery, Tripoli University, Libya

²Associate Professor of Radiology, Tripoli Medical Center, Tripoli University, Libya

³Neurosurgeon, Ali Omar Askar Neurosurgery Hospital, Tripoli University, Libya

Submission: July 17, 2017; **Published:** August 10, 2017

***Corresponding author:** Muffaq lashhab, Neurosurgeon, Ali Omar Askar Neurosurgery Hospital, Tripoli University, Libya, Email: muaffaq.mkls@gmail.com

Abstract

The spinal meningioma is rare in children with an annual incidence of 5 case per 1000000 people. In the medical literature, approximately 60 cases of spinal meningioma have been reported in the otherwise normal pediatric population [1]. We report a case of a 9 year old girl who presented with weakness of both lower limbs and unable to walk. Imaging studies demonstrated an extradural spinal tumor at dorsal spine. The patient was operated and tumor was totally removed. The postoperative course was excellent. Histology confirmed the diagnosis of meningioma. After surgery there was improvement in signs and symptoms.

Keywords: Thoracic spine; Pediatric; Fracture

Case Report

Clinical presentation

A nine year-old female patient who has no known disease before, she was referred to our clinic with complaints of sudden progressive spastic paraparesis of lower limbs (over a period of three weeks) with bilateral foot drop and inability to walk. Moreover, no infections had been observed prior to symptoms, neither history of trauma nor relevant epidemiological antecedents or previous exposure to radiation. Clinical

examination revealed bilateral reduction in muscular strength of the legs and thighs (both power grade I), Deep tendon reflexes (DTRs) of the lower extremities were observed to be bilaterally slightly hyperactive with ankle clonus, while no changes in sensitivity to pain or proprioception were observed. An examination of the upper limbs was within normal Limits, The results of dermatologic examination were normal. Her hematological and biochemical investigations were normal.




Figure 1: Thoracic spine preoperative MRI images (a) with contrast (b) T2 sagittal and axial at T5,T6,T7 level show extradural meningioma displacing the cord (white arrow)

Radiological findings

Brain MRI showed no abnormality. A Dorso-lumbar MRI Figure 1. revealed a mass measures 40*18mm, which is Iso-intense on T1 image and hyper -intense on T2 image at the anterior extradural space extending from D5 and D7 vertebrae on left side causing compression and displacing spinal cord to right side and posteriorly, it shows homogenous contrast enhancement with posterior wall vertebral body erosion and destruction of the posterior element, she was considered to have an aggressive hemangioma as the initial radiological diagnosis.

Surgical Intervention

The patient underwent D5-D7 laminectomy (Figure 2 & 3) and exposed a whitish tumor compressing the adjacent dura. The tumor was adherent to the lateral side of dura as well as the posterior wall of dorsal vertebrae (located anterior to the cord). The tumor was well separated from the spinal cord and could be microsurgically resected completely with its dural attachment (Figure 3).




Figure 2: Intraoperative image shows D5-D7 laminectomy and an exposed tumor compressing the adjacent dura.




Figure 3: Dorsal spine post-operative MRI images T 1 (a) and T 2 (b) with contrast (c) sagittal and axial at D5,D6,D7 level show total mass resection.

- A. Pathology: revealed a choroid meningioma.
- B. Postoperatively: the patient evolution was satisfactory, with recovery of muscular strength to power grade 3 in three days. She is currently undergoing motor rehabilitation with good clinical evolution.

Discussion

Spinal meningiomas have rarely been describe in pediatrics age [2]. There are very few case reports in the literature describing intraspinal meningioma in children less than 13 years of age [2]. Spinal meningiomas comprise about 25 to 46% of all spinal tumors [3]. Pediatric spinal cord tumors represent only 5% of tumors of the central nervous system, 25% of which occur in the intradural-extramedullary compartment

[4]. Reports in adult patients describe that most of cases tend to occur in females (about 2:1). However, this association is not replicated in children, in which there are reports of an about 1.2:1 predominance of cases in males [2]. A retrospective analysis done by Greene and colleagues of 20 patients showed a median age of presentation of the tumors of about 13 years old [2]. Case reports in younger age groups included those in 14 and 20 month-old infants, while only a few reports of spinal meningiomas in children of school age were found [5,6].

Known risk factors for developing meningiomas include previous exposure to radiation which is the most environmental risk factor predisposing to meningiomas [7]. And the presence of type 2 neurofibromatosis, in which meningiomas are present in 25% to 40% of children with neoplasia [8]. The link to type

2 neurofibromatosis is becoming clearer, with loss of a tumor suppressor gene in the chromosome 22 (NF2). Other gene mutations may contribute to the progression of the meningiomas, leading to the anaplastic type.

Gezen et al. [9] reviewed the main characteristics of 36 cases of spinal cord meningiomas. Regarding to the symptomatology, pain and sensory loss were the most common symptoms associated with some degree of weakness. A grading system proposed by the World Health Organization (WHO) is the most used in the classification of the meningiomas. There are 3 grades - benign (grade I), atypical (grade II) and anaplastic or malignant (grade III).

Conclusion

MRI imaging is the gold standard for diagnosing meningiomas in children. The most common lesion pattern is the finding of T1 and T2 iso-intense masses with different degrees of peritumoral edema. Surgery is the best treatment in cases of spinal meningiomas. Complete resection is possible in most of the cases and is associated with good postoperative functional improvement. This emphasises the need for an early diagnosis and an early referral to a neurosurgical center. Recurrence of the tumor is rare, and in most series the reported rate of recurrence ranged from 3-14% [9,10], Radiosurgery should be considered for the exceptional cases involving recurrent and symptomatic spinal tumors. Follow up using clinical exams and serial imaging is important to enable early detection and selection of further surgical interventions when needed [11,12].

References

1. Gelabert-González M, García-Allut A, Martínez-Rumbo R (2006) Spinal meningiomas. *Neurocirugia (Astur)* 17(2): 125-131.
2. Ciapetta P, Domenicucci M (1988) Spinal meningiomas: prognosis and recovery factors in 22 cases with severe motor deficits. *Acta Neurol Scand* 77(1): 27-30.
3. Sze G (2002) Neoplastic disease of spinal cord. In: Atlas SW (Ed.), *Magnetic Resonance Imaging of the Brain and Spine*. Lippincott Williams & Wilkins, Philadelphia, USA, pp. 1-2000.
4. Binning M, Klimo Jr P, Gluf W, Goumnerova L (2007) Spinal tumors in children. *Neurosurg Clin N Am* 18(4): 631-658.
5. Riemenschneider MJ, Perry A, Reifenberger G (2006) Histological classification and molecular genetics of meningiomas. *Lancet Neurol* 5: 1045-1054.
6. Baumgartner JE, Sorenson JM (1996) Meningioma in the pediatric population. *J Neurooncol* 29(3): 223-228.
7. Levy WJ Jr, Bay J, Dohn D (1982) Spinal cord meningioma. *J Neurosurg* 57(6): 804-812.
8. Solero CL, Fornari M, Giombini S, Lasio G, Oliveri G, Cimino C, et al. (1989) Spinal meningiomas: Review of 174 operated cases. *Neurosurgery* 25(2): 153-160.
9. Gezen F, Kahraman S, Canakci Z, Bedük A (2000) Review of 36 cases of spinal cord meningioma. *Spine* 25: 727-731.
10. Klekamp J, Samii M (1999) Surgical results for spinal meningiomas. *Surg Neurol* 52(6): 552-562.
11. Ferraz-Filho JRL, Floriano VH, Felipe LF, Rocha-Filho JA (2008) Clinical-radiological aspects of primary extracranial meningioma of the ethmoid sinus in a child. *Arq Neuropsiquiatr* 66(2A): 274-275.
12. Wang XQ, Zeng XW, Zhang BY, Dou YF, Wu JS, Jiang CC, et al. (2012) Spinal meningioma in childhood: clinical features and treatment. *Child Nerv Syst* 28: 129-136.



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/OAJNN.2017.05.555659](https://doi.org/10.19080/OAJNN.2017.05.555659)

Your next submission with Juniper Publishers will reach you the below assets

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats (Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission
<https://juniperpublishers.com/online-submission.php>