



A Review of Recent Status on Stingrays (Chondrichthyes: Dasyatidae) In Indonesian Waters



Yuanita Windusari¹ and Muhammad Iqbal^{2*}

¹Sriwijaya University, Department of Biology, Indonesia

²Sriwijaya University, Faculty of Science, Indonesia

Submission: December 17, 2017; **Published:** March 09, 2018

***Corresponding author:** Muhammad Iqbal, Sriwijaya University, Faculty of Science, Conservation Biology Program, Jalan Padang Selasa 524, Palembang, South Sumatra 30129, Indonesia Email: kpbsos26@yahoo.com

Abstract

In this paper, the status of stingrays (family Dasyatidae) in Indonesian waters are summarized. There are 40 valid species of stingrays recorded here. Based on distribution of stingrays within seven faunal region of Indonesia, Kalimantan of Indonesian Borneo has 29 species, the largest diversity among other islands; following Sumatra (27 species), Java (25 species), Sulawesi (14 species), Lesser Sundas (16), Mollucas (9 species) and Papua (17 species). According to the recent of Red List of Threatened Species of the International Union for the Conservation of Nature's (IUCN 2017), there are 18 species are listed as threatened species (under Endangered and Vulnerable status), 2 as Near Threatened (NT), 3 as Data Deficient (DD), 5 as Least Concern (LC) and 12 species are Not Evaluated (NE). Five species listed as Endangered should have high prioritize for conservation: *Fluvitrygon oxyrhynchus* (Sauvage, 1878); *F. kittipongi* (Vidthayanon & Roberts, 2005); *F. signifer* (Compagno & Roberts, 1982); *P. solocirostris* Last, Manjaji & Yearsley, 2005; and *Urogymnus polylepis* (Bleeker, 1852).

Keywords: Review; Dasyatidae; Indonesian waters; Diversity; Conservation

Introduction

The Indonesian archipelago is one of the richest marine biodiversity areas in the world [1]. The country has 4.720 species of fishes, making one the greatest diversity of fish fauna in the world [2]. Many iconic group of fishes known collectively as the 'ray' occur in Southeast Asia, particularly in Indonesian waters [3-5]. One of 'ray' is stingrays (family Dasyatidae), groups of small to very large myliobatiform fishes (adults from 22cm to 260cm DW) and distinguished by the following combination of characters: body variably depressed with a well-formed oval, circular or rhombic disc that fully incorporates head; snout angular to obtuse and sometimes very elongate; nasal curtain well developed, skirt-shaped, rectangular or bilobed; five gill slits; oral papillae usually present on floor of mouth; tail moderately stout to slender-based and more or less elongated (sometimes very elongate and whip-like); dorsal surface variably covered with dermal denticles, thorns and/or tubercles, smooth to very spiny and often with a median thorn row and/or a median denticle band; no dorsal or caudal fins; 1-4 prominent caudal stings, positioned on tail well posterior to pelvic fins; skin folds variably developed on the ventral and sometimes dorsal midline of tail; dorsal surface plain to strongly patterned, usually

darker than ventral surface [6-9]. Recent phylogenetic studies, supported by morphological data, have provided evidence that the Dasyatidae is monophyletic and consists of four major subgroups, the subfamilies Dasyatinae, Neotrygoninae, Urogymninae and Hypolophinae; and a morphologically based review of 89 currently recognised species [6]. In this review, we summary and update current knowledge of the stingrays in Indonesian waters.

Diversity of Stingrays in Indonesian waters

Table 1 shows 40 valid species of stingrays that occur in Indonesian waters. This checklist is compiled and shortlisted from recent major references [6-8]. The fishes were divided within seven faunal regions of Indonesia, where faunal regions match administrative boundaries and has no political significance [10]. Conservation status covering global threatened species follows recent International Union for Conservation of Nature (IUCN) Redlist 2017 [11].

Two species have been added recently for Sumatran waters [12,13], *Fluvitrygon oxyrhynchus* and *Urogymnus polylepis* (Figure 1 & 2). Based on distribution of stingrays within seven

faunal region of Indonesia, Kalimantan of Indonesian Borneo has 29 species, the largest diversity among other islands; following Sumatra (27 species) and Java (25 species). There are only

nine species of stingrays found in Mollucas, indicate the lowest number compare to other islands in Indonesian waters.



Figure 1: A skin showing dorsal surface of Endangered *Urogymnus polylepis* (Bleeker, 1852) on 2 April 2016 in Bungin river, Banyuasin district, South Sumatra (photograph by Muhammad Iqbal).




Figure 2 : *Fluvitrygon oxyrhynchus* (Sauvage, 1878) caught by local fisherman on 24 August 2006 in Musi drainage, South Sumatra province. This is one of very rare stingray in Indonesia (photograph by Muhammad Iqbal).

Conservation of Stingrays in Indonesian waters

The status of most of the world’s rays have been assessed using a standard international system to characterise the risk of extinction of species: the International Union for the Conservation of Nature’s (IUCN) Red List of Threatened Species [14]. Following recent IUCN (2017), there are 18 species are listed as threatened species (under Endangered and Vulnerable status), 2 species listed as Near Threatened, 3 species listed as Data Deficient, 5 species listed as Least Concern and 12 species as Not Evaluated (Table 1). Five species having high prioritize for conservation because listed as Endangered: *Fluvitrygon*

oxyrhynchus; *F. kittipongi* (Vidthayanon & Roberts, 2005); *F. signifer* (Compagno & Roberts, 1982); *P. solocirostris* Last, Manjaji & Yearsley, 2005; and *Urogymnus polylepis*. Four of these species are inhabit freshwaters. To secure the status of stingrays locally and globally, in addition to the addressing the issue of data deficiency, accurate species identification is needed, the strict enforcement of fishing and protection of measures, a significant increase in scientific observer coverage to monitor catches, and increase research on gear modifications, fishing methods, and habitat identification aimed at itigating bycatch and discard mortality of rays [14].

Table 1: List of stingrays in Indonesian waters.

No.	Species	IUCN Status	Distribution							
			S	K	J	S	M	L	P	
1	<i>Bathytoshia lata</i> (Garman, 1880)	LC			+				+	
2	<i>B. heterura</i> (Bleeker, 1852)	NE	+	+	+					
3	<i>B. javaensis</i> (Last & White, 2013)	NE			+					
4	<i>Fluvitrygon oxyrhynchus</i> (Sauvage, 1878)	EN	+	+						
5	<i>F. kittipongi</i> (Vidthayanon & Roberts, 2005)	EN		+						
6	<i>F. signifer</i> (Compagno & Roberts, 1982)	EN	+	+						
7	<i>Hemitrygon bennettii</i> (Müller & Henle, 1841)	DD	+	+						
8	<i>H. longicauda</i> (Last & White, 2013)	NE								+
9	<i>H. parvonigra</i> (Last & White, 2008)	DD			+				+	
10	<i>Himantura uarnak</i> (Gmelin, 1789)	VU	+	+	+	+	+	+	+	
11	<i>H. australis</i> Last, White & Naylor, 2016	NE								+
12	<i>H. leoparda</i> Manjaji-Matsumoto & Last, 2008	VU	+	+	+	+	+	+	+	+

13	<i>H. undulata</i> (Bleeker, 1852)	VU	+	+	+	+			
14	<i>Maculabatis gerrardi</i> (Gray, 1851)	VU	+	+	+				
15	<i>M. astra</i> (Last, Manjaji-Matsumoto & Pogonoski, 2008)	LC	+	+	+	+		+	
16	<i>M. marucra</i> (Bleeker, 1852)	NE							+
17	<i>M. pastinacoides</i> (Bleeker, 1852)	VU	+	+	+				
18	<i>Megatrygon microps</i> (Annandale, 1908)	DD	+	+		+		+	+
19	<i>Neotrygon annotata</i> (Last, 1987)	NT			+			+	+
20	<i>N. australiae</i> (Last, White & Seret, 2016)	NE						+	+
21	<i>N. caeruleopunctata</i> (Last, White & Seret, 2016)	NE	+		+				
22	<i>N. onectatus</i> (Last, White & Seret, 2016)	NE	+	+	+	+			
23	<i>N. picta</i> Last & White, 2008	LC							+
24	<i>N. varidens</i> (Garman, 1855)	NE		+					
25	<i>Pastinachus ater</i> (Macleay, 1883)	LC	+	+	+	+	+	+	+
26	<i>P. gracilicaudus</i> Last & Manjaji-Matsumoto, 2010	NE	+	+	+				
27	<i>P. solocirostris</i> Last, Manjaji & Yearsley, 2005	EN	+	+	+				
28	<i>P. stellurostris</i> Last, Fahmi & Naylor, 2010	NE		+					
29	<i>Pateobatis fai</i> (Jordan & Seale, 1906)	VU	+	+	+			+	+
30	<i>P. hortlei</i> (Last, Manjaji-Matsumoto & Kailola, 2006)	VU							+
31	<i>P. jenkinsii</i> (Annandale, 1909)	VU	+	+	+	+	+	+	+
32	<i>P. uarnacoides</i> (Bleeker, 1852)	VU	+	+	+	+			
33	<i>Pteroplatytrygon violacea</i> (Bonaparte, 1832)	LC	+	+	+	+	+	+	+
34	<i>Taeniura lymma</i> (Forsskål, 1775)	NT	+	+	+	+	+	+	+
35	<i>Taeniurops meyeni</i> (Müller & Henle, 1841)	VU	+	+	+	+	+	+	+
36	<i>Telatrygon biasa</i> Last, White & Naylor, 2016	NE	+	+	+	+			
37	<i>Urogymnus polylepis</i> (Bleeker, 1852)	EN	+	+					
38	<i>U. asperrimus</i> (Bloch & Schneider, 1801)	VU	+	+	+	+	+	+	+
39	<i>U. granulatus</i> (Macleay, 1883)	VU	+	+	+		+	+	+
40	<i>U. lobistoma</i> (Manjaji-Matsumoto & Last, 2006)	VU	+	+					
Total			27	29	25	14	9	16	17

Note:

I. IUCN global status, CR = Critically Endangered, EN = Endangered, VU = Vulnerable, NT = Near Threatened, DD = Data Deficient, LC = Least Concern, NE = Not Evaluated.

II. Distribution abbreviations, S = Sumatera, K = Kalimantan (Indonesian Borneo), J = Java, S = Sulawesi, M = Mollucas, L = Lesser Sunda, P = Papua (West Papua, Indonesian Papua).

Conclusion

Recent status of stingrays (family Dasyatidae) in Indonesian waters are summarized here. This general review may give obvious diversity and conservation challenges. We suppose that further research and monitoring are needed to clarify local status and distribution of stingrays in Indonesian waters, particularly in area where rarely explored.

Acknowledgment

Few visits to Musi basin have drawn attention to stingrays status in Sumatra. Thanks to Project Aware Foundation who granted MI to conducting aquatic survey in South Sumatran waters. We are very grateful to all researchers, students and volunteers who facilitated us during various fieldworks in South Sumatran waters during 2006-2016.

References

1. Tomascik T, Mah AJ, Nontji A, Moosa ML (1997) The ecology of Indonesian seas, Part II. Periplus Editions, Singapore.
2. Froese R, Pauly D (2017) FishBase, World Wide Web electronic publication.
3. Compagno LJV, Roberts TR (1982) Freshwater stingrays (Dasyatidae) of Southeast Asia and New Guinea, with description of a new species of *Himantura* and reports of unidentified species. *Environmental Biology of Fishes* 7(4): 321–339.
4. Last PR, de Carvalho MR, Naylor GJP, Seret B, Stehmann MFW, et al. (2016) Introduction. In: Last others (Eds.). *Rays of the world*, Cornell University Press, Ithaca, USA, pp. 1-9.
5. Kottelat M (2013) The fishes of inland waters of Southeast Asia: a catalogue and core bibliography of the fishes known to occur in freshwaters, mangroves and estuaries. *Raffles Bulletin of Zoology Supplement* 27: 1-663.
6. Last PR, Naylor GJP, Manjaji-Matsumoto BM (2016) A revised classification of the family Dasyatidae (Chondrichthyes: Myliobatiformes) based on new morphological and molecular insights. *Zootaxa* 4139(3): 345-368.
7. Last PR, Compagno LJV (1999) Dasyatidae. In: Carpenter KE, Niem VH (Eds.). *FAO species identification guide for fishery purposes. The living marine resources of the Western Central Pacific. Volume 3. Batoid fishes, chimaeras and bony fishes part 1 (Elopidae to Linophrynidae)*, Food and Agriculture Organization of the United Nations, Rome, Italy, pp. 1479-1510.
8. Last PR, Manjaji-Matsumoto BM, Naylor GJP, White WT (2016) Stingrays, Family Dasyatidae. In: Last others (Eds.). *Rays of the world*, Cornell University Press, Ithaca, USA, pp. 522-618.
9. Nelson JS, Terry C, Grande TC, Wilson MVH (2016) *Fishes of the World*. (5th edn), John Wiley & Sons, Hoboken, USA.
10. Andrew P (1992) The birds of Indonesia, a checklist (Peter's sequence), Kukila Checklist no. 1, Indonesian Ornithological Society, Jakarta, Indonesia.
11. IUCN (2017) The IUCN Red List of Threatened Species.
12. Iqbal M, Yustian I (2016) Occurrence of the giant freshwater stingray *Urogymnus polylepis* in Sumatra, Indonesia (Chondrichthyes: Dasyatidae). *Ichthyological Exploration of Freshwaters* 27(4): 333-336.
13. Iqbal M, Setiawan D, Ajiman (2017) Presence of *Fluvitrygon oxyrhynchus* in Sumatra, Indonesia (Chondrichthyes: Dasyatidae). *Ichthyological Exploration of Freshwaters* 28(1): 85-87.
14. Kyne PM (2016) Ray Conservation. In: Last others (Eds.). *Rays of the world*, Cornell University Press, Ithaca, USA, pp. 21-24.



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/OFOAJ.2018.06.555690](https://doi.org/10.19080/OFOAJ.2018.06.555690)

**Your next submission with Juniper Publishers
will reach you the below assets**

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission
<https://juniperpublishers.com/online-submission.php>