

Blood Flow Restriction Training Considerations



***Carl Biscontini**

Physical Therapist, Above & Beyond Physical Therapy, USA

Submission: September 28, 2017; **Published:** October 03, 2017

***Corresponding author:** Carl Biscontini, Physical Therapist, Above & Beyond Physical Therapy, Queen Creek, Arizona, USA, Tel: (480)987-1870; Email: carl.biscontini@gmail.com.

What is Blood Flow Restriction Training?

In the rehabilitation world, there is a never ending search for advancements in treatment methodologies, techniques, and regimens to improve effectiveness and efficiency in care. Often times, we are faced with various conditions coming into the clinic with restrictions and/or limitations, which can then alter the way we attempt to successfully execute our plan of care. We are always striving to keep on adding to our clinical toolbox to help ensure that each patient gets the best opportunity for a successful recovery. In the recent years, blood flow restriction training (BFR) has evolved with the advancements in evidence-based practice models, covering a wide population, from professional athletes to NASA.

The purpose of blood flow restriction training is to utilize a pneumatic tourniquet cuff to restrict venous return while maintaining arterial blood flow to the musculature. Once applied, exercises are then performed at a lower intensity, ranging from 20-30% of a subject's 1RM [1,2]. The physiological impacts on growth hormone and IGF-1 help to simulate the effects of higher intensity training to help promote protein synthesis to enhance strength and hypertrophy [3-5]. There has been a wide array of various training strategies that have been utilized with the use of BFR training, most occurring at a higher volume of repetitions with short rest periods between sets.

Loenneke et al. [1] assessed various protocols of utilization, from low level exercises of 3 sets of 15 repetitions with 30 second rest breaks, to various low level walking/cycling protocols of varying times. The researchers concluded in their meta analysis that the resistance training programs performed more favorably to the aerobic protocols, and that the greatest effect size was achieved when performed 2 to 3 days per week [6]. From a clinical perspective, safety of interventions must always be the first factor to take into consideration before performing. There are various contraindications in regards to utilizing blood flow restriction training, so it is vital that we properly assess risk factors prior to application. Some contraindications for the use of blood flow restriction training use include but not limited to:

- i. History of any of the following: DVT, Cancer, Open Fracture, Severe Hypertension, Stroke, Diabetes,
Current Infection, Acidosis, Spinal Cord Injury, Rhabdomyolysis
- ii. Compromised circulation or peripheral vascular systems
- iii. Currently on medications that may alter clotting factor
- iv. Extremities utilizing dialysis ports
- v. Increased intracranial pressure

Clark et al. [7] assessed the safety following a 4 weeks protocol of blood flow restriction training and illustrated gains in strength measures without the alteration of nerve/vascular function, markers of coagulation or inflammation, while increasing fibrinolytic activity.

Practice Implications for Blood Flow Restriction Training

There are several clinical situations where blood flow restriction training can help to preserve current musculature while minimizing further damage to the injured area. In post-operative rehabilitation, especially when immobilization and protocol limitations are set, will provide the clinical with another modality to help facilitate hypertrophy at a lower load, while maintaining the integrity of the surgical intervention. This could provide an avenue to optimize earlier phases in the recovery process, and make clinical treatment more valuable and relevant to progress. An example of this could be in the earlier stages of ACL repairs where maintaining integrity of the graft is most important, and incorporating blood flow restriction training into Phase I exercises, such as quad sets, hamstring sets, hip straight leg raise activities.

Injuries resulting in restrictions of non-weight bearing can allow the condition to treat other portions of the kinetic chain while maintaining restrictions. Examples of this would be maintaining muscular strength of lower extremity following a high grade ankle sprain, where a patient may be placed in an immobilization boot. Performing open chain quadriceps

and hamstring activities can be performed with minimal resistance while achieving gains and reducing risks of kinetic chain dysfunction once weight bearing restrictions are lifted. In patients with osteoporosis, this can provide an avenue to provide high intensity muscular gains and potentially improving bone health [8].

Conclusion

This article is intended to help potentially establish studies to develop accelerated rehabilitation protocols implementing blood flow restriction training to help optimize care and reduce potential length of care. Blood flow restriction training could help to minimize degradation of various tissues and structures due to immobilization process. The potential to provide a high intensity physiological response with a low intensity volume of training can help to add another valuable tool to our clinical toolbox to deliver optimal care.

References

1. Loenneke J, Fahs C, Rossow L, Abe T, Bembem M (2012) The anabolic benefits of venous blood flow restriction training may be induced by muscle cell swelling. *Med Hypotheses* 78(1): 151-154.
2. Yasuda T, Loenneke JP, Thiebaud RS, Abe T (2012) Effects of Blood Flow Restricted Low-Intensity Concentric or Eccentric Training on Muscle Size and Strength. *PLoS ONE* 7(12): 1-12.
3. Madarame H, Neya M, Ochi E, Nakazato K, Sato Y, et al. (2008) Cross-Transfer Effects of Resistance Training with Blood Flow Restriction. *Med Sci Sports Exerc* 40(2): 258-263.
4. Cook SB, Brown KA, Deruisseau K, Kanaley JA, Ploutz-Snyder LL (2010) Skeletal muscle adaptations following blood flow-restricted training during 30 days of muscular unloading. *J Appl Physiol* 109(2): 341-349.
5. Fujita S, Abe T, Drummond MJ, Cadenas JG, Dreyer HC, et al. (2007) Blood flow restriction during low-intensity resistance exercise increases S6K1 phosphorylation and muscle protein synthesis. *J Appl Physiol* 103(3): 903-910.
6. Loenneke JP, Wilson JM, Marín PJ, Zourdos MC, Bembem MG (2011) Low intensity blood flow restriction training: a meta-analysis. *Eur J Appl Physiol* 112(5): 1849-1859.
7. Clark BC, Manini TM, Hoffman RL, Williams PS, Guiler MK, et al. (2010). Relative safety of 4 weeks of blood flow-restricted resistance exercise in young, healthy adults. *Scand J Med Sci Sports* 21(5): 653-662.
8. Karabulut M, Bembem DA, Sherk VD, Anderson MA, Abe T (2011) Effects of high intensity resistance training and low-intensity resistance training with vascular restriction on bone markers in older men. *Eur J Appl Physiol* 111(8): 1659-1667.



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/OROAJ.2017.08.555747](https://doi.org/10.19080/OROAJ.2017.08.555747)

Your next submission with Juniper Publishers will reach you the below assets

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission

<https://juniperpublishers.com/online-submission.php>